

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l47

L48            13 L47

=> d l48 1-13 abs ibib hitstr



L48 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AB A series of bis(ether amines), primarily 1,2-bis(4-aminophenoxy)benzenes but including 1,3- and 1,4-bis(4-aminophenoxy)benzenes, with fluoro and alkyl substituents were synthesized. These diamines were prepared by F displacement with 4-FC6H4NO2 or its derivs., or NO2 displacement with 1,4-C6H4(NO2)2, and various phenylenediols and their alkyl or fluoro derivs. The resulting bis(ether nitro) compds. were reduced to the corresponding bis(ether amines).

ACCESSION NUMBER: 1998:376601 CAPLUS

DOCUMENT NUMBER: 129:135942

TITLE: Methyl- and fluoro-substituted bis(4-aminophenoxy)benzenes. A convenient method of synthesis

AUTHOR(S): Eastmond, G. C.; Paprotny, J.

CORPORATE SOURCE: Donnan Laboratories, University Liverpool, Liverpool, L69 7ZD, UK

SOURCE: Synthesis (1998), (6), 894-898

CODEN: SYNTBF; ISSN: 0039-7881

PUBLISHER: Georg Thieme Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 129:135942

IT 210492-44-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(preparation of methylated and fluorinated

[(aminophenoxy)phenoxy]benzenamin

es)

RN 210492-44-7 CAPLUS

CN Benzene, 1,2-bis(2-fluoro-4-nitrophenoxy)- (9CI) (CA INDEX NAME)



IT 210492-53-8P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of methylated and fluorinated

[(aminophenoxy)phenoxy]benzenamin



es)

RN 210492-53-8 CAPLUS

CN Benzenamine, 4,4'-[1,2-phenylenebis(oxy)]bis[3-fluoro- (9CI) (CA INDEX NAME)

L48 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

(Continued)



L48 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

(Continued)

L48 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AB Mixing a resin and a certain phthalocyanine compound having the ability to absorb near IR rays, provides a resin composition which is useful as a heat

radiation-shielding material which is semi-transparent or transparent for visible light but blocks heat rays. C black may also be added to enhance the heat-shielding effect. Thus, polycarbonate containing 0.003% VOPc(BuNH)878 [Pc = phthalocyanine; substituent octakis(butylamino)octafluorophthalocyanine] was molded into a film having transmittance and heat transmittance (JIS R-3106) 78.9 and 62.4%; vs. 89.2 and 84.6, resp., for polycarbonate without the IR absorber.

ACCESSION NUMBER: 1998:79369 CAPLUS

DOCUMENT NUMBER: 128:141505

TITLE: Compositions for shields for heat radiation

INVENTOR(S): Kaieda, Osamu; Yodoshi, Takashi; Morita, Ken;

Matsuura, Michio

Nippon Shokubai Co., Japan

SOURCE: U.S., 15 pp., Cont.-in-part of U.S. Ser. No. 180,488, abandoned.

CODEN: USXQAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5712332	A	19980127	US 1995-478739	19950607
PRIORITY APPLN. INFO.:			JP 1993-4326	A 19930113
			US 1994-180488	B2 19940112

OTHER SOURCE(S): MARPAT 128:141505

IT 163464-84-4



RL: MOD (Modifier or additive use); USES (Uses)

(polymer compns. containing certain phthalocyanines for shields for

heat radiation)


RN 163464-84-4 CAPLUS

CN Zinc, [N,N',N'',N'''-tetrahexyl-1,3,4,8,10,11,15,17,18,22,24,25-dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-tetraminato(2-)-κN29,κN30,κN31,κN32]-, (SP-4-1)- (9CI) (CA INDEX NAME)




PAGE 2-A

PAGE 3-A



PAGE 4-A



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L48 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Facile synthesis of Cp<sup>+</sup>Ru<sup>+</sup> (Cp<sup>+</sup> = pentamethylcyclopentadienyl)  $\pi$ -complexes of highly electron deficient aroms., including 1,3,5-trichloro-, 1,2,4,5-tetrachloro-, pentachloro-, and hexachlorobenzene, is accomplished by the ligand exchange reaction with (Cp<sup>+</sup>Ru(CH<sub>3</sub>CN)<sub>3</sub>)+SO<sub>3</sub>CF<sub>3</sub><sup>-</sup> in polar solvents under mild reaction conditions. The extraordinary activating ability of the Cp<sup>+</sup>Ru<sup>+</sup> moiety is demonstrated by rapid and quant. nucleophilic substitution reactions of the 1,3,5-tri- and 1,2,4,5-tetrachlorobenzene  $\pi$ -complexes with potassium phenoxide, thiophenoxide, 4-aminophenoxide, 4-chlorophenoxide, and 4-fluorophenoxide.

This methodol. allows syntheses of highly substituted and functionalized aromatic compds.

ACCESSION NUMBER: 1995:701727 CAPLUS  
DOCUMENT NUMBER: 123:169866  
TITLE: Synthesis and Nucleophilic Substitution of Highly Chlorinated Arene ( $\eta$ 5-Pentamethylcyclopentadienyl)ruthenium  $\pi$ -Complexes  
AUTHOR(S): Dembek, Alexa A.; Fagan, Paul J.  
CORPORATE SOURCE: Experimental Station, DuPont Central Research and Development, Wilmington, DE, 19880-0328, USA  
SOURCE: Organometallics (1995), 14(8), 3741-5  
CODEN: ORGM7; ISSN: 0276-7333  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 123:169866  
IT 167227-89-6P


RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 167227-89-6 CAPLUS  
CM Ruthenium(II), [(1,2,3,4,5- $\eta$ )-1,2,3,4,5-pentamethyl-2,4-cyclopentadien-1-yl][(1,2,3,4,5,6- $\eta$ )-1,2,4,5-tetrakis(4-fluorophenoxy)benzene]-, salt with trifluoromethanesulfonic acid (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 167227-88-5  
CHF C40 H33 F4 O4 Ru  
CCI CCS

PAGE 1-A

PAGE 3-A




CM 2

CRN 37181-39-8  
CHF C F3 O3 S



PAGE 2-A





L48 ANSWER 5 OF 13 CAPIUS COPYRIGHT 2005 ACS on STN  
AB The phthalocyanines have the formula (R)AMKPC(OR1)b(OR2)cXd, where Pc is

a phthalocyanine nucleus; M is a metal atom; k is the inverse of 1/2 the valency of M; R is a heterocyclic compound or a tertiary amine; each R1

is aryl or heterocyclyl; each R2 is alkyl or cycloalkyl; each X is H or halogen; a is 1 or 2; b = 1-8; c = 0-4; d = 4-15; and b + c + d = 16. Thus, 4,5-dichlorophthalonitrile was heated 2 h at 120° with 2 equiv p-tert-octylphenol in DMF containing K2CO3 to give 4,5-bis(4-tert-octylphenoxy)phthalonitrile, which was cyclocondensed with FeCl3.6H2O, urea, and NH4 molybdate to give the octakis-substituted FePc, which was heated in pyridine to give the dipyriddy complex as a bright green solid with  $\lambda_{max}$  658 nm in toluene.

ACCESSION NUMBER: 1995:662388 CAPIUS  
DOCUMENT NUMBER: 123:85956  
TITLE: Substituted metal phthalocyanines  
INVENTOR(S): Reynolds, Stephen James; White, Raymond Lesley  
PATENT ASSIGNEE(S): Zeneca Ltd., UK  
SOURCE: Eur. Pat. Appl., 12 pp.  
CODEN: EPOXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 638614	A1	19950215	EP 1994-305325	19940720
R: CH, DE, FR, JP 07070458	A2	19950314	JP 1994-190542	19940812
PRIORITY APPL. INFO.:			GB 1993-16820	A 19930812

OTHER SOURCE(S): MARPAT 123:85956  
IT 164652-85-1P  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(preparation of substituted metal phthalocyanines)  
RN 164652-85-1 CAPIUS  
CN Iron, [2,3,9,10,16,17,23,24-octakis(4-fluorophenoxy)-29H,31H-phthalocyaninato(2-)-N29,N30,N31,N32]bis(pyridine)-, (OC-6-12)- (9CI)  
(CA INDEX NAME)

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*


L48 ANSWER 5 OF 13 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 2-A



PAGE 3-A

L48 ANSWER 6 OF 13 CAPIUS COPYRIGHT 2005 ACS on STN  
GI



AB The mixts. contain a transparent resin [e.g., polycarbonate, poly(Me methacrylate), PVC, or poly(ethylene terephthalate)], a near-IR-absorbing phthalocyanine compound I (Z1-16 = SR1, OR2, H, halo, NHY; Z1 of Z1-16 = NHY; R1-2 = Ph, substituted Ph, C1-20 alkyl; Y = Ph, substituted Ph, C1-8 alkyl; M = VO, Zn, Cu, SnCl2, Co, etc.), and, optionally, carbon black. Replacing part of the I with carbon black does not decrease the heat-radiation-shielding ability of the mixts. The mixts. are useful as moldings which transmit visible light while blocking near-IR radiation (i.e., they absorb the heat from sunlight).


ACCESSION NUMBER: 1995:550904 CAPIUS  
DOCUMENT NUMBER: 122:316019  
TITLE: Heat-radiation-shielding mixtures of polymers and near-IR-absorbing phthalocyanine compounds  
INVENTOR(S): Kaieda, Osamu; Yodoshii, Takashi; Morita, Ken; Matsuura, Michio  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Eur. Pat. Appl., 28 pp.  
CODEN: EPOXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 607031	A1	19940720	EP 1994-300205	19940112
EP 607031	B1	19991006		
R: BE, DE, ES, FR, GB, IT, NL				
JP 06264050	A2	19940920	JP 1993-352128	19931229
ES 2136700	T3	19991201	ES 1994-300205	19940112


L48 ANSWER 6 OF 13 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
PRIORITY APPL. INFO.:

OTHER SOURCE(S): MARPAT 122:316019  
IT 163464-84-4  
RL: MOA (Modifier or additive use); POF (Polymer in formulation); PRP (Properties); TEM (Technical or engineered material use); USES (Uses)  
(near-IR absorber; heat-radiation-shielding compns. containing polymers and)  
RN 163464-84-4 CAPIUS  
CN Zinc, [N,N',N'',N'''-tetrahexyl-1,3,4,8,10,11,15,17,18,22,24,25-dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-tetraminato(2-)-N29,N30,N31,N32]-, (SP-4-1)- (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 2-A



PAGE 3-A



PAGE 4-A



L48 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AB Hexafluorobenzene reacts readily with a variety of trimethylsilyl ethers ROSiMe<sub>3</sub> (R = CF<sub>3</sub>CH<sub>2</sub>, FCH<sub>2</sub>CH<sub>2</sub>, H(CF<sub>2</sub>)<sub>n</sub>CH<sub>2</sub> (n = 2, 4), CF<sub>3</sub>(CF<sub>2</sub>)<sub>6</sub>CH<sub>2</sub>, CF<sub>3</sub>(CF<sub>2</sub>)<sub>5</sub>CH<sub>2</sub>CH<sub>2</sub>, MeSiOCH<sub>2</sub>CH<sub>2</sub>, C<sub>6</sub>F<sub>5</sub>OCH<sub>2</sub>CH<sub>2</sub>, C<sub>6</sub>H<sub>5</sub>, 4-FC<sub>6</sub>H<sub>4</sub>) to give from mono- to hexapolyfluoroalkoxy- and polyfluoroaryloxy-benzenes. The structure of C<sub>6</sub>(OCH<sub>2</sub>CF<sub>3</sub>)<sub>6</sub> has been confirmed by single-crystal X-ray anal.

The perfluorinated ether C<sub>6</sub>F<sub>5</sub>OCF<sub>2</sub>CF<sub>3</sub> may be synthesized from C<sub>6</sub>F<sub>5</sub>OCH<sub>2</sub>CF<sub>3</sub> by chlorination and subsequent fluorination with SbF<sub>3</sub>/SbCl<sub>5</sub>. The chlorination of 5,6,7,8-tetrafluoro-1,4-benzodioxane is also discussed.

ACCESSION NUMBER: 1995:57102 CAPLUS

DOCUMENT NUMBER: 122:31037

TITLE: Reaction of hexafluorobenzene with trimethylsilyl ethers

AUTHOR(S): Zhang, Y. F.; Kirchmeier, Robert L.; Shreeve, Jean'ne M.

CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 83844-2343, USA

SOURCE: Journal of Fluorine Chemistry (1994), 68(3), 287-92

CODEN: JFLCAR; ISSN: 0022-1139

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 122:31037


IT 159752-90-6 159752-93-9 159752-94-0

RL: PRP (Properties)

(Reaction of hexafluorobenzene with trimethylsilyl ethers)

RN 159752-90-6 CAPLUS


CN Benzene, hexakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)



RN 159752-93-9 CAPLUS


CN Benzene, fluoropentakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)

L48 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 159752-94-0 CAPLUS

CN Benzene, 1,4-difluoro-2,3,5,6-tetrakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)



\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I, useful as material for hardeners for fluorinated epoxy resins (no data), is prepared via II [R = cyano, CO<sub>2</sub>H]. Thus, a mixture of

tetrafluorophthalonitrile, tetrafluorohydroquinone, and Et<sub>3</sub>N in DMF was heated at 35° for 30 to give 211 II [R = cyano], which was treated with 60% H<sub>2</sub>SO<sub>4</sub> at 150° for 5 h to give 261 II [R = CO<sub>2</sub>H], which was refluxed with Ac<sub>2</sub>O for 2 h to 521 I.

ACCESSION NUMBER: 1994:630662 CAPIUS

DOCUMENT NUMBER: 121:230662

TITLE: preparation of a perfluorinated hexacarboxylic acid as

material for hardeners for fluorinated epoxy resins

INVENTOR(S): Sasaki, Shigekuni; Matsura, Tooru; Ando, Shinji

PATENT ASSIGNEE(S): Nippon Telegraph &amp; Telephone, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JTOGAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06157501	AZ	19940603	JP 1992-340986	19921130
PRIORITY APPL. INFO.:			JP 1992-340986	19921130

OTHER SOURCE(S): CASREACT 121:230662


IT 158394-12-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and conversion into trianhydride)

RN 158394-12-8 CAPIUS

CN 1,2-Benzenedicarboxylic acid, 4,5-bis[4-(3,4-dicarboxy-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)




IT 158394-11-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and hydrolysis of)

RN 158394-11-7 CAPIUS

CN 1,2-Benzenedicarbonitrile, 4,5-bis[4-(3,4-dicyano-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)




IT 158394-13-9P

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)

(preparation of a perfluorinated hexacarboxylic acid as material)

RN 158394-13-9 CAPIUS

CN 1,3-Isobenzofurandione, 4,7-difluoro-5,6-bis[2,3,5,6-tetrafluoro-4-[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-5-isobenzofuranyl)oxy]phenoxy]- (9CI) (CA INDEX NAME)



AB Metal or nonmetal complexes of 1,4,8,11,15,18,22,25-octafluorophthalocyanines in which the 2-, 3-, 9-, 10-, 16-, 17-, 23-, and

24-positions are substituted with F, substituted phenoxy, substituted phenylthio, (un)substituted anilino, alkylamino, substituted alkoxy, or substituted alkylthio groups, not all being F, are prepared directly from the appropriate phthalonitriles or by substitution reactions on the hexadecafluorophthalocyanine complexes. Thus, reaction of tetrafluorophthalonitrile with PhSH and KF in MeCN at 50° for 12 h gave 3,6-difluoro-4,5-bis(phenylthio)phthalonitrile, which was heated

with ZnI<sub>2</sub> in α-ClC<sub>10</sub>H<sub>7</sub> at 200° to give the ZnPC derivative; this was sulfonated with ClSO<sub>3</sub>H in CHCl<sub>2</sub>CHCl<sub>2</sub> to introduce 15 sulfo groups/mol. on the phenylthio groups. The product had λ<sub>max</sub> 725.0 nm in EtOH and 700 nm in H<sub>2</sub>O and showed solubility of 10% in EtOH, 3.9% in EtOCH<sub>2</sub>CH<sub>2</sub>OH, and 12% in H<sub>2</sub>O; its IR spectrum and those of 30 addnl. phthalocyanines are given.

ACCESSION NUMBER: 1993:429981 CAPIUS

DOCUMENT NUMBER: 119:29981

TITLE: Fluorophthalocyanines, their preparation, and near infrared ray-absorbing materials containing them

INVENTOR(S): Kaieda, Osamu; Yodoshi, Takashi; Itoh, Hideki;

ONOTAKI, Miho

PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan

SOURCE: Eur. Pat. Appl., 78 pp.

CODEN: EPXKDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 523959	AZ	19930120	EP 1992-306443	19920714
EP 523959	A3	19940316		
EP 523959	B1	19991013		
R: CH, DE, FR, GB, IT, LI, NL				
JP 05078364	A2	19930330	JP 1992-42586	19920228
JP 2907624	B2	19990621		
US 5359056	A	19941025	US 1992-913028	19920714
JP 05345861	A2	19931227	JP 1992-274125	19921013
JP 2812624	B2	19981022		
PRIORITY APPL. INFO.:			JP 1991-174019	A 19910715
			JP 1992-42586	A 19920228
			JP 1992-94522	A 19920414
			JP 1991-33276	A1 19910228

OTHER SOURCE(S): MARPAT 119:29981

IT 147712-52-5P


RL: IMF (Industrial manufacture); PREP (Preparation)

(preparation of near-IR-absorbing)



RN 147712-52-5 CAPIUS

CN Vanadate (8-), [1,4,8,11,15,18,22,25-octafluoro-2,3,9,10,16,17,23,24-octakis(2-fluorophenoxy)-29H,31H-phthalocyaninato(10-)-κN29,κN30,κN31,κN32]oxo-, octahydrogen, octasulfo deriv. (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




IT 148131-11-7  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (sulfonation of)  
 RN 148131-11-7 CAPLUS  
 CN Vanadium,  
 [1,4,8,11,15,18,22,25-octafluoro-2,3,9,10,16,17,23,24-octakis(2-fluorophenoxy)-29H,31H-phthalocyaninato(2-)-N29,N30,N31,N32]oxo-,


PAGE 1-A




PAGE 2-A



L48 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI



L48 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 (dye, for sublimation type thermal-transfer recording material)  
 RN 122533-59-9 CAPLUS  
 CN 9,10-Anthracenedione, 1,4-diamino-2,3-bis(2-fluorophenoxy)- (9CI) (CA INDEX NAME)




AB Sublimation type thermal recording transfer sheets contain a dye component comprising  $\geq 1$  dye I (R = H, halo, CF<sub>3</sub>, alkyl, alkoxy) and  $\geq 1$  dye II (R<sub>1</sub> = H, halo, OH, CF<sub>3</sub>, alkyl, alkoxy, aralkyloxy, alkoxyalkoxy, aryloxyalkoxy, acyl, alkylcarbonyloxy, (N-substituted) carbamoyl, alkylsulfonyl, arylsulfonyl, (N-substituted) sulfamoyl, alkylsulfonyloxy, arylsulfonyloxy). The dyes have good solubility or dispersibility in resins, and the transfer sheets exhibit good transferability and provide stable, high color quality images. Thus, a mixture of Et cellulose 6, I (R = H) 1, and II (R<sub>1</sub> = OMe-4) 1 part was coated on a PET film to give a thermal transfer film, which gave clear magenta images with high d. and good lightfastness.

ACCESSION NUMBER: 1989:523963 CAPLUS  
 DOCUMENT NUMBER: 111:123963  
 TITLE: Thermal recording transfer sheets containing two kinds of anthraquinone type dyes  
 INVENTOR(S): Hashimoto, Kyoyasu; Suzuki, Yasuyuki  
 PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
 CODEN: JKOKAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 01077584	A2	19890323	JP 1987-235450	19870918
PRIORITY APPLN. INFO.:			JP 1987-235450	19870918

IT 122533-59-9  
 RL: USES (Uses)



AB Reaction of perfluoroarenes with alcs. and phenols, especially steroids, under phase-transfer catalysis gave perfluoroaryl ethers. These could be selectively cleaved with NaOMe in DMF. E.g., reaction of estrone (I) with F3CC6F5 in CH2Cl2 containing Bu4N+ HSO4- and aqueous NaOH at room temperature gave 95%

3-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]estra-1,3,5-trien-17-one (II). Cleavage of II with NaOMe in DMF at 10° for 2 h gave 87% I.  
ACCESSION NUMBER: 1985:471577 CAPLUS  
DOCUMENT NUMBER: 103:71577  
TITLE: Octafluorotoluene as a reagent for the selective protection of alcoholic and phenolic functions. Synthesis and cleavage of perfluorotolyl and other perfluoroaryl ethers of steroids and other model compounds  
AUTHOR(S): Jarman, Michael; McCague, Raymond  
CORPORATE SOURCE: Cancer Res. Campaign Lab., Inst. Cancer Res., Sutton, SM2 5PX, UK  
SOURCE: Journal of Chemical Research, Synopses (1985), (4), 114-15  
CODEN: JRPSDC; ISSN: 0308-2342  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 103:71577  
IT 97631-77-1P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
RN 97631-77-1 CAPLUS  
CN Benzene, 1,2-bis[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)




L48 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Reaction of phenols with Br2C6H3CF3 at 170-300° in the presence of Cu and alkali gave mono- and diphenoxybenzotrifluorides. Thus, 2,5- and 3,4-Br2C6H3CF3, powdered Cu, KI, and m-CF3C6H4OH at 200-20° gave 10.6 (m-CF3C6H4O)C6H4CF3, 25.4 (m-CF3C6H4O)C6H3BrCF3, and 42.4% (m-CF3C6H4O)2C6H3CF3 isomers.  
ACCESSION NUMBER: 1975:170346 CAPLUS  
DOCUMENT NUMBER: 82:170346  
TITLE: Phenoxyl derivatives of trifluoromethylbenzene  
INVENTOR(S): Schlafke, Rolf; Jenkner, Herbert  
PATENT ASSIGNEE(S): Chemische Fabrik Kalk G.m.b.H.  
SOURCE: Ger. Offen., 21 pp.  
CODEN: GWXXBX  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2325878	A1	19741212	DE 1973-2325878	19730522

PRIORITY APPLN. INFO.: DE 1973-2325878 A 19730522

IT 54846-40-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
RN 54846-40-1 CAPLUS  
CN Benzene, 1,2-bis[4-(4-fluorophenoxy)-4-(trifluoromethyl)- (9CI) (CA INDEX NAME)



L48 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN  
GI For diagram(s), see printed CA Issue  
AB The PbO2 oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).  
ACCESSION NUMBER: 1974:520135 CAPLUS  
DOCUMENT NUMBER: 81:120135  
TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol  
AUTHOR(S): Deniville, Leon; Huynh Anh Hoa  
CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conservatoire Natl. Arts  
SOURCE: Metiers, Paris, Fr.  
Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 487-90  
CODEN: BSCFAS; ISSN: 0037-8968  
DOCUMENT TYPE: Journal  
LANGUAGE: French  
IT 53279-71-3P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
RN 53279-71-3 CAPLUS  
CN 1,4-Benzenediol, 2,3,5,6-tetrakis(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          65.57      3354.90

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)    SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE                        -9.49      -226.30
```

FILE 'REGISTRY' ENTERED AT 20:18:50 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
 DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8


TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
 information enter HELP PROP at an arrow prompt in the file or refer  
 to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=>
Uploading C:\Program Files\Stnexp\Queries\10718532.str
```



```
chain nodes :
1 13 23 25 26 28 29 30 31 32 33 35
ring nodes :
2 3 4 5 6 7 8 9 10 11 12 16 17 18 19 20 21 22
chain bonds :
1-4 8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31
22-35
ring bonds :
2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22
18-19 19-20 20-21 21-22
exact/norm bonds :
8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31
exact bonds :
1-4 22-35
```

normalized bonds :

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
18-19 19-20 20-21 21-22

G1:N,X

G2:X,Ak,H


Match level :

1:CLASS 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:Atom 12:Atom 13:CLASS 14:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom  
21:Atom 22:Atom 23:CLASS 25:CLASS 26:CLASS 28:CLASS 29:CLASS 30:CLASS  
31:CLASS 32:CLASS 33:CLASS 35:CLASS

L49 STRUCTURE UPLOADED

=> d query

L49 STR



G1 N,X

Structure attributes must be viewed using STN Express query preparation.

=> s l49

SAMPLE SEARCH INITIATED 20:19:19 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 127 TO ITERATE

100.0% PROCESSED 127 ITERATIONS  
SEARCH TIME: 00.00.01

14 ANSWERS

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 1864 TO 3216  
PROJECTED ANSWERS: 56 TO 504

L50 14 SEA SSS SAM L49

=> s l49 full

FULL SEARCH INITIATED 20:19:25 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 2836 TO ITERATE

100.0% PROCESSED 2836 ITERATIONS  
SEARCH TIME: 00.00.01

194 ANSWERS

L51 194 SEA SSS FUL L49

=> d l51 1-194 abs ibib hitstr  
'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'  
'IBIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'  
'HITSTR' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN  
SAM - Index Name, MF, and structure - no RN  
FIDE - All substance data, except sequence data  
IDE - FIDE, but only 50 names  
SQIDE - IDE, plus sequence data  
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used  
SQD - Protein sequence data, includes RN  
SQD3 - Same as SQD, but 3-letter amino acid codes are used  
SQN - Protein sequence name information, includes RN  
  
CALC - Table of calculated properties  
EPROP - Table of experimental properties  
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract  
APPS -- Application and Priority Information  
BIB -- CA Accession Number, plus Bibliographic Data  
CAN -- CA Accession Number  
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)  
IND -- Index Data  
IPC -- International Patent Classification  
PATS -- PI, SO  
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels  
IBIB -- BIB, indented, with text labels  
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.



The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.  
HELP FORMATS -- To see detailed descriptions of the predefined formats.  
ENTER DISPLAY FORMAT (IDE):nos  
'NOS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN  
SAM - Index Name, MF, and structure - no RN  
FIDE - All substance data, except sequence data  
IDE - FIDE, but only 50 names  
SQIDE - IDE, plus sequence data  
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used  
SQD - Protein sequence data, includes RN  
SQD3 - Same as SQD, but 3-letter amino acid codes are used  
SQN - Protein sequence name information, includes RN  
  
CALC - Table of calculated properties  
EPROP - Table of experimental properties  
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract  
APPS -- Application and Priority Information  
BIB -- CA Accession Number, plus Bibliographic Data  
CAN -- CA Accession Number  
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)  
IND -- Index Data  
IPC -- International Patent Classification  
PATS -- PI, SO  
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels  
IBIB -- BIB, indented, with text labels  
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.  
HELP FORMATS -- To see detailed descriptions of the predefined formats.  
ENTER DISPLAY FORMAT (IDE):bib.  
'BIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN  
SAM - Index Name, MF, and structure - no RN  
FIDE - All substance data, except sequence data  
IDE - FIDE, but only 50 names  
SQIDE - IDE, plus sequence data  
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used  
SQD - Protein sequence data, includes RN  
SQD3 - Same as SQD, but 3-letter amino acid codes are used  
SQN - Protein sequence name information, includes RN  
  
CALC - Table of calculated properties  
EPROP - Table of experimental properties  
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract  
APPS -- Application and Priority Information  
BIB -- CA Accession Number, plus Bibliographic Data  
CAN -- CA Accession Number  
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)  
IND -- Index Data  
IPC -- International Patent Classification  
PATS -- PI, SO  
STD -- BIB, IPC, and NCL  
  
IABS -- ABS, indented, with text labels  
IBIB -- BIB, indented, with text labels  
ISTD -- STD format, indented  
  
OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels  
  
SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented,

with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.  
HELP FORMATS -- To see detailed descriptions of the predefined formats.  
ENTER DISPLAY FORMAT (IDE):abs  
'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN  
SAM - Index Name, MF, and structure - no RN  
FIDE - All substance data, except sequence data  
IDE - FIDE, but only 50 names  
SQIDE - IDE, plus sequence data  
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used  
SQD - Protein sequence data, includes RN  
SQD3 - Same as SQD, but 3-letter amino acid codes are used  
SQN - Protein sequence name information, includes RN  
  
CALC - Table of calculated properties  
EPROP - Table of experimental properties  
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract  
APPS -- Application and Priority Information  
BIB -- CA Accession Number, plus Bibliographic Data  
CAN -- CA Accession Number  
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)  
IND -- Index Data  
IPC -- International Patent Classification  
PATS -- PI, SO  
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels  
IBIB -- BIB, indented, with text labels  
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.  
HELP FORMATS -- To see detailed descriptions of the predefined formats.  
ENTER DISPLAY FORMAT (IDE):none  
'NONE' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN  
SAM - Index Name, MF, and structure - no RN  
FIDE - All substance data, except sequence data  
IDE - FIDE, but only 50 names  
SQIDE - IDE, plus sequence data  
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used  
SQD - Protein sequence data, includes RN  
SQD3 - Same as SQD, but 3-letter amino acid codes are used  
SQN - Protein sequence name information, includes RN  
  
CALC - Table of calculated properties  
EPROP - Table of experimental properties  
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract  
APPS -- Application and Priority Information  
BIB -- CA Accession Number, plus Bibliographic Data  
CAN -- CA Accession Number  
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)  
IND -- Index Data  
IPC -- International Patent Classification  
PATS -- PI, SO  
STD -- BIB, IPC, and NCL  
  
IABS -- ABS, indented, with text labels  
IBIB -- BIB, indented, with text labels  
ISTD -- STD format, indented  
  
OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels  
  
SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.

HELP FORMATS -- To see detailed descriptions of the predefined formats.

<-----User Break----->

ENTER DISPLAY FORMAT (IDE): rn

L51 ANSWER 1 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 788823-51-8 REGISTRY

L51 ANSWER 2 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 788823-46-1 REGISTRY

L51 ANSWER 3 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 788823-43-8 REGISTRY

L51 ANSWER 4 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 788823-34-7 REGISTRY

L51 ANSWER 5 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 713525-85-0 REGISTRY

L51 ANSWER 6 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 676515-14-3 REGISTRY

L51 ANSWER 7 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 676515-05-2 REGISTRY

L51 ANSWER 8 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 676515-04-1 REGISTRY

L51 ANSWER 9 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 666698-30-2 REGISTRY

L51 ANSWER 10 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 659720-09-9 REGISTRY

L51 ANSWER 11 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 548465-93-6 REGISTRY

L51 ANSWER 12 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 548465-90-3 REGISTRY



L51 ANSWER 13 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 548465-86-7 REGISTRY

L51 ANSWER 14 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 500577-29-7 REGISTRY

L51 ANSWER 15 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 500577-27-5 REGISTRY

L51 ANSWER 16 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 482574-81-2 REGISTRY

L51 ANSWER 17 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 477274-17-2 REGISTRY

L51 ANSWER 18 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 400742-27-0 REGISTRY

L51 ANSWER 19 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 398130-67-1 REGISTRY

L51 ANSWER 20 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 380574-44-7 REGISTRY

L51 ANSWER 21 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 373642-43-4 REGISTRY

L51 ANSWER 22 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 373641-71-5 REGISTRY

L51 ANSWER 23 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 373641-22-6 REGISTRY

L51 ANSWER 24 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 354989-14-3 REGISTRY

L51 ANSWER 25 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 352536-58-4 REGISTRY

L51 ANSWER 26 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 351438-36-3 REGISTRY

L51 ANSWER 27 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 339321-86-7 REGISTRY

L51 ANSWER 28 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 332406-15-2 REGISTRY

L51 ANSWER 29 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 331660-58-3 REGISTRY

L51 ANSWER 30 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 331655-53-9 REGISTRY

L51 ANSWER 31 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 330220-63-8 REGISTRY

L51 ANSWER 32 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 329964-53-6 REGISTRY

L51 ANSWER 33 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 329219-28-5 REGISTRY

L51 ANSWER 34 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 329213-56-1 REGISTRY

L51 ANSWER 35 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 329212-77-3 REGISTRY

L51 ANSWER 36 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 328275-82-7 REGISTRY

L51 ANSWER 37 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 328275-38-3 REGISTRY

L51 ANSWER 38 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 328088-45-5 REGISTRY

L51 ANSWER 39 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 326893-10-1 REGISTRY

L51 ANSWER 40 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 326893-03-2 REGISTRY

L51 ANSWER 41 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 326892-95-9 REGISTRY

L51 ANSWER 42 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 326892-94-8 REGISTRY

L51 ANSWER 43 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 326892-41-5 REGISTRY

L51 ANSWER 44 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 318258-94-5 REGISTRY



L51 ANSWER 45 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 314048-52-7 REGISTRY

L51 ANSWER 46 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 313968-95-5 REGISTRY

L51 ANSWER 47 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 313249-29-5 REGISTRY

L51 ANSWER 48 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 312583-57-6 REGISTRY

L51 ANSWER 49 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 306766-00-7 REGISTRY

L51 ANSWER 50 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 290352-21-5 REGISTRY

L51 ANSWER 51 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 290352-20-4 REGISTRY

L51 ANSWER 52 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 290352-19-1 REGISTRY

L51 ANSWER 53 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 245126-36-7 REGISTRY

L51 ANSWER 54 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 245126-34-5 REGISTRY

L51 ANSWER 55 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN . 245126-33-4 REGISTRY

L51 ANSWER 56 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 245126-32-3 REGISTRY

L51 ANSWER 57 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 245126-31-2 REGISTRY

L51 ANSWER 58 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-46-6 REGISTRY

L51 ANSWER 59 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-45-5 REGISTRY

L51 ANSWER 60 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-44-4 REGISTRY

LS1 ANSWER 61 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-42-2 REGISTRY

LS1 ANSWER 62 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-41-1 REGISTRY

LS1 ANSWER 63 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-40-0 REGISTRY

LS1 ANSWER 64 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 223791-39-7 REGISTRY

L51 ANSWER 65 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 221117-66-4 REGISTRY

L51 ANSWER 66 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 213693-17-5 REGISTRY

L51 ANSWER 67 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 210492-62-9 REGISTRY

L51 ANSWER 68 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 210492-61-8 REGISTRY

L51 ANSWER 69 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 208122-72-9 REGISTRY

L51 ANSWER 70 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 208122-71-8 REGISTRY

L51 ANSWER 71 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 208038-75-9 REGISTRY

L51 ANSWER 72 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 208038-72-6 REGISTRY

LS1 ANSWER 73 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 208038-65-7 REGISTRY

LS1 ANSWER 74 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 207504-84-5 REGISTRY

LS1 ANSWER 75 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-09-7 REGISTRY

LS1 ANSWER 76 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-08-6 REGISTRY



L51 ANSWER 77 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-05-3 REGISTRY

L51 ANSWER 78 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-02-0 REGISTRY

L51 ANSWER 79 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-01-9 REGISTRY

L51 ANSWER 80 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203808-00-8 REGISTRY

LS1 ANSWER 81 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-99-2 REGISTRY

LS1 ANSWER 82 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-98-1 REGISTRY

LS1 ANSWER 83 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-97-0 REGISTRY

LS1 ANSWER 84 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-96-9 REGISTRY

LS1 ANSWER 85 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-95-8 REGISTRY

LS1 ANSWER 86 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-94-7 REGISTRY

LS1 ANSWER 87 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-93-6 REGISTRY

LS1 ANSWER 88 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-92-5 REGISTRY

L51 ANSWER 89 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 203807-91-4 REGISTRY

L51 ANSWER 90 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 193757-00-5 REGISTRY

L51 ANSWER 91 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186981-83-9 REGISTRY

L51 ANSWER 92 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186981-82-8 REGISTRY

LS1 ANSWER 93 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186981-79-3 REGISTRY

LS1 ANSWER 94 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-53-0 REGISTRY

LS1 ANSWER 95 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-52-9 REGISTRY

LS1 ANSWER 96 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-51-8 REGISTRY

L51 ANSWER 97 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-20-1 REGISTRY

L51 ANSWER 98 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-17-6 REGISTRY

L51 ANSWER 99 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186980-15-4 REGISTRY

L51 ANSWER 100 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186979-65-7 REGISTRY

LS1 ANSWER 101 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186979-63-5 REGISTRY

LS1 ANSWER 102 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 186979-62-4 REGISTRY

LS1 ANSWER 103 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-23-8 REGISTRY

LS1 ANSWER 104 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-22-7 REGISTRY

LS1 ANSWER 105 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-21-6 REGISTRY

LS1 ANSWER 106 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-20-5 REGISTRY

LS1 ANSWER 107 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-18-1 REGISTRY

LS1 ANSWER 108 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-17-0 REGISTRY



L51 ANSWER 109 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-15-8 REGISTRY

L51 ANSWER 110 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-14-7 REGISTRY

L51 ANSWER 111 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-13-6 REGISTRY

L51 ANSWER 112 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 185697-11-4 REGISTRY

LS1 ANSWER 113 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 181507-42-6 REGISTRY

LS1 ANSWER 114 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 177032-01-8 REGISTRY

LS1 ANSWER 115 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 177032-00-7 REGISTRY

LS1 ANSWER 116 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 177031-89-9 REGISTRY

L51 ANSWER 117 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 173447-54-6 REGISTRY

L51 ANSWER 118 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 168839-60-9 REGISTRY

L51 ANSWER 119 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 167227-89-6 REGISTRY

L51 ANSWER 120 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 167227-88-5 REGISTRY

LS1 ANSWER 121 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165380-05-2 REGISTRY

LS1 ANSWER 122 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-94-2 REGISTRY

LS1 ANSWER 123 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-93-1 REGISTRY

LS1 ANSWER 124 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-92-0 REGISTRY

LS1 ANSWER 125 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-86-2 REGISTRY

LS1 ANSWER 126 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-83-9 REGISTRY

LS1 ANSWER 127 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-78-2 REGISTRY

LS1 ANSWER 128 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-65-7 REGISTRY

0

L51 ANSWER 129 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-64-6 REGISTRY

L51 ANSWER 130 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-63-5 REGISTRY

L51 ANSWER 131 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-62-4 REGISTRY

L51 ANSWER 132 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-60-2 REGISTRY

L51 ANSWER 133 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-59-9 REGISTRY

L51 ANSWER 134 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-58-8 REGISTRY

L51 ANSWER 135 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-57-7 REGISTRY

L51 ANSWER 136 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-56-6 REGISTRY

LS1 ANSWER 137 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-55-5 REGISTRY

LS1 ANSWER 138 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-54-4 REGISTRY

LS1 ANSWER 139 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-53-3 REGISTRY

LS1 ANSWER 140 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-47-5 REGISTRY



LS1 ANSWER 141 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-46-4 REGISTRY

LS1 ANSWER 142 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-44-2 REGISTRY

LS1 ANSWER 143 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-43-1 REGISTRY

LS1 ANSWER 144 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-42-0 REGISTRY

L51 ANSWER 145 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 165379-41-9 REGISTRY

L51 ANSWER 146 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163709-89-5 REGISTRY

L51 ANSWER 147 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163709-88-4 REGISTRY

L51 ANSWER 148 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163464-84-4 REGISTRY

L51 ANSWER 149 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163302-12-3 REGISTRY

L51 ANSWER 150 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163301-37-9 REGISTRY

L51 ANSWER 151 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163301-22-2 REGISTRY

L51 ANSWER 152 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163301-21-1 REGISTRY

L51 ANSWER 153 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163300-97-8 REGISTRY

L51 ANSWER 154 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 163300-89-8 REGISTRY

L51 ANSWER 155 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 159752-94-0 REGISTRY

L51 ANSWER 156 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 159752-93-9 REGISTRY

L51 ANSWER 157 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 159752-90-6 REGISTRY

L51 ANSWER 158 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 156589-15-0 REGISTRY

L51 ANSWER 159 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141863-03-8 REGISTRY

L51 ANSWER 160 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141228-92-4 REGISTRY

L51 ANSWER 161 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141228-91-3 REGISTRY

L51 ANSWER 162 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141228-90-2 REGISTRY

L51 ANSWER 163 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141216-01-5 REGISTRY

L51 ANSWER 164 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141216-00-4 REGISTRY

L51 ANSWER 165 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141215-99-8 REGISTRY

L51 ANSWER 166 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141215-97-6 REGISTRY

L51 ANSWER 167 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 141215-95-4 REGISTRY

L51 ANSWER 168 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 122297-49-8 REGISTRY

LS1 ANSWER 169 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 122281-58-7 REGISTRY

LS1 ANSWER 170 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 122281-57-6 REGISTRY

LS1 ANSWER 171 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 113397-96-9 REGISTRY

LS1 ANSWER 172 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 88483-66-3 REGISTRY



L51 ANSWER 173 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 88483-51-6 REGISTRY

L51 ANSWER 174 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 88483-48-1 REGISTRY

L51 ANSWER 175 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 60793-43-3 REGISTRY

L51 ANSWER 176 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 53279-71-3 REGISTRY

L51 ANSWER 177 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 28096-26-6 REGISTRY

L51 ANSWER 178 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 19228-15-0 REGISTRY

L51 ANSWER 179 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-42-6 REGISTRY

L51 ANSWER 180 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-40-4 REGISTRY

L51 ANSWER 181 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-37-9 REGISTRY

L51 ANSWER 182 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-36-8 REGISTRY

L51 ANSWER 183 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-35-7 REGISTRY

L51 ANSWER 184 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-30-2 REGISTRY

L51 ANSWER 185 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-29-9 REGISTRY

L51 ANSWER 186 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 17742-25-5 REGISTRY

L51 ANSWER 187 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 16065-60-4 REGISTRY

L51 ANSWER 188 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 16031-36-0 REGISTRY

L51 ANSWER 189 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 15963-72-1 REGISTRY

L51 ANSWER 190 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 15038-90-1 REGISTRY

L51 ANSWER 191 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 14901-49-6 REGISTRY

L51 ANSWER 192 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 14796-04-4 REGISTRY

L51 ANSWER 193 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 5026-82-4 REGISTRY

L51 ANSWER 194 OF 194 REGISTRY COPYRIGHT 2005 ACS on STN  
RN 5026-81-3 REGISTRY

}

=> d his

(FILE 'HOME' ENTERED AT 19:20:11 ON 12 JAN 2005)

FILE 'REGISTRY' ENTERED AT 19:20:22 ON 12 JAN 2005

L1 STRUCTURE UPLOADED

L2 0 S L1

L3 2 S L1 FULL

FILE 'CAPLUS' ENTERED AT 19:20:55 ON 12 JAN 2005

L4 1 S L3

FILE 'REGISTRY' ENTERED AT 19:21:43 ON 12 JAN 2005

L5 STRUCTURE UPLOADED

L6 0 S L5

L7 2 S L5 FULL

FILE 'CAPLUS' ENTERED AT 19:22:28 ON 12 JAN 2005

L8 0 S L7

L9 0 S L7

FILE 'REGISTRY' ENTERED AT 19:22:54 ON 12 JAN 2005

L10 STRUCTURE UPLOADED

L11 2 S L10 FULL

FILE 'CAOLD' ENTERED AT 19:23:44 ON 12 JAN 2005

L12 1 S L11

FILE 'REGISTRY' ENTERED AT 19:24:58 ON 12 JAN 2005

FILE 'REGISTRY' ENTERED AT 19:26:23 ON 12 JAN 2005

L13 STRUCTURE UPLOADED

L14 0 S L13

L15 6 S L13 FULL

FILE 'CAPLUS' ENTERED AT 19:26:57 ON 12 JAN 2005

L16 2 S L15

FILE 'REGISTRY' ENTERED AT 19:27:31 ON 12 JAN 2005

L17 STRUCTURE UPLOADED

L18 0 S L17

L19 2 S L17 FULL

FILE 'CAPLUS' ENTERED AT 19:28:57 ON 12 JAN 2005

L20 0 S L19

FILE 'REGISTRY' ENTERED AT 19:29:05 ON 12 JAN 2005

FILE 'REGISTRY' ENTERED AT 19:29:28 ON 12 JAN 2005

FILE 'REGISTRY' ENTERED AT 19:29:38 ON 12 JAN 2005

SET TERMSET E#

DEL SEL Y

SEL L19 2 RN

L21 1 S E1/RN

SET TERMSET LOGIN

FILE 'CAOLD' ENTERED AT 19:29:42 ON 12 JAN 2005

L22 1 S L21

FILE 'REGISTRY' ENTERED AT 19:30:10 ON 12 JAN 2005  
L23 STRUCTURE UPLOADED  
L24 1 S L23  
L25 56 S L23 FULL

FILE 'CAPLUS' ENTERED AT 19:30:52 ON 12 JAN 2005  
L26 39 S L25

FILE 'REGISTRY' ENTERED AT 19:41:56 ON 12 JAN 2005  
L27 STRUCTURE UPLOADED  
L28 0 S L27  
L29 31 S L27 FULL

FILE 'CAPLUS' ENTERED AT 19:43:02 ON 12 JAN 2005  
L30 14 S L29

FILE 'REGISTRY' ENTERED AT 19:46:27 ON 12 JAN 2005  
L31 STRUCTURE UPLOADED  
L32 0 S L31  
L33 10 S L31 FULL

FILE 'CAPLUS' ENTERED AT 19:47:32 ON 12 JAN 2005  
L34 4 S L33

FILE 'REGISTRY' ENTERED AT 19:48:14 ON 12 JAN 2005  
L35 STRUCTURE UPLOADED  
L36 1 S L35  
L37 9 S L35 FULL

FILE 'CAPLUS' ENTERED AT 19:48:58 ON 12 JAN 2005  
L38 5 S L37

FILE 'REGISTRY' ENTERED AT 19:49:43 ON 12 JAN 2005  
L39 STRUCTURE UPLOADED  
L40 2 S L39  
L41 271 S L39 FULL

FILE 'CAPLUS' ENTERED AT 19:50:32 ON 12 JAN 2005  
L42 231 S L41

FILE 'REGISTRY' ENTERED AT 20:14:18 ON 12 JAN 2005  
L43 STRUCTURE UPLOADED  
L44 1 S L43  
L45 STRUCTURE UPLOADED  
L46 0 S L45  
L47 23 S L45 FULL

FILE 'CAPLUS' ENTERED AT 20:17:16 ON 12 JAN 2005  
L48 13 S L47

FILE 'REGISTRY' ENTERED AT 20:18:50 ON 12 JAN 2005  
L49 STRUCTURE UPLOADED  
L50 14 S L49  
L51 194 S L49 FULL

=> fil caplus  
COST IN U.S. DOLLARS

SINCE FILE TOTAL  
ENTRY SESSION



FULL ESTIMATED COST	201.85	3556.75
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-226.30

FILE 'CAPLUS' ENTERED AT 20:21:48 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.



FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
 FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l51

L52            54 L51

=> d l52 1-54 abs ibib hitstr




AB Thyroid receptor ligands are provided which have the general formula I [wherein: R1 = (un)substituted CONR5R6, CH2NR5R6, NR5COR6, OR7, R8, 4-R9-4,5-dihydrooxazol-2-yl; R2, R3 = H, halo, Cl-4 alkyl or C3-5 cycloalkyl, provided that at least 1 of R2 and R3 = H; R4 = (CH2)nR13 or (CH2)nCONR16CR13R14R15; R5, R6 = H, (hetero)aryl, (cyclo)alkyl, or (hetero)aralkyl; R7 = (hetero)aryl, alkyl, or (hetero)aralkyl; R8 = (hetero)aryl or cycloalkyl; R9 = R7 or H; R10 = H, halo, cyano, or alkyl; R11, R12 = H, halo, alkoxy, OH, cyano, or alkyl; R13 = COOH and esters, phosphonic and phosphinic acid and esters, sulfonic acid, tetrazole, hydroxamic acid, thiazolidinedione, acylsulfonamide, or other carboxylic acid surrogates; R14, R15 = H, alkyl; or R14R15 = (CH2)2-5, forming 3- to 6-membered cycloalkyl rings; R16 = H or Cl-4 alkyl; R17 and R18 = H, halo, or alkyl; n = 0-4; X = O, S, S(O)2, S(O), Se, CO, NH, or CH2]. In addition, a method is provided for preventing, inhibiting or treating diseases or disorders associated with metabolism dysfunction, or which are dependent upon the expression of, a T3 regulated gene, wherein a compound I is administered therapeutically. Claims cover the above, as well as pharmaceutical compns. containing I, and methods of coadministration of I with other compds., particularly certain antidiabetic agents. Compds. I include selective agonists, partial agonists, antagonists, and partial antagonists of thyroid receptors (no data). Approx. 168 compds. were prepared. For instance, Me (3,5-dibromo-4-hydroxyphenyl)acetate underwent O-arylation with (4-MeOC6H4)2I+ BF4-, and the resultant 4-methoxyphenyl ether derivative underwent a sequence of: (1) formylation in the 3-position, (2) O-demethylation, (3) oxidation of the aldehyde to an acid, (4) amidation of the acid, and (5) alkaline saponification of the ester, to give title compound II.

ACCESSION NUMBER: 2004:927006 CAPLUS  
DOCUMENT NUMBER: 141:395288  
TITLE: New [3,5-dihalo-4-(4-hydroxyphenoxy)phenyl]acetic acid derivatives useful as thyroid receptor ligands, and


L52 ANSWER 1 OF 54 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
their preparation, pharmaceutical compositions, and methods of use  
INVENTOR(S): Ryono, Dennis E.; Hangeland, Jon J.; Friends, Todd J.;  
Dejneka, Tamara; Devasthale, Pratik; Caringal, Yolanda  
V.; Zhang, Minsheng; Doweiko, Arthur M. P.; Malm, Johan; Sanin, Andrei  
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
SOURCE: PCT Int. Appl., 94 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004093799	A2	20041104	WO 2004-US11883	20040416
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2005004184	A1	20050106	US 2004-826100	20040415
PRIORITY APPLN. INFO.:		US 2003-463774P P 20030418		


IT 788823-34-7P 788823-43-8P 788823-46-1P  
788823-51-8P  
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(drug candidate; preparation of [dihalo(hydroxyphenoxy)phenyl]acetic acid  
deriv. as thyroid receptor ligands)  
RN 788823-34-7 CAPLUS  
CN Benzenecetic acid, 3,5-dibromo-4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]- (9CI) (CA INDEX NAME)




RN 788823-43-8 CAPLUS  
CN Benzenecetic acid, 3,5-dibromo-4-[3-(3-fluorophenoxy)-4-hydroxyphenoxy]- (9CI) (CA INDEX NAME)



RN 788823-46-1 CAPLUS  
CN Benzenecetic acid, 3,5-dibromo-4-[3-(3-chloro-4-fluorophenoxy)-4-hydroxyphenoxy]- (9CI) (CA INDEX NAME)



RN 788823-51-8 CAPLUS  
CN Benzenecetic acid, 3,5-dichloro-4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]- (9CI) (CA INDEX NAME)




L52 ANSWER 2 OF 54 CAPLUS COPYRIGHT 2005 ACS ON STN  
AB A F compound has perfluorostyrene introduced at a terminal end, characterized in that the introduction of perfluorostyrene results in a facile fabrication of thin films by a UV curing or a thermal curing, high thermal stability and chemical resistance, and low optical propagation loss and birefringence when applied to waveguides. The structure of the F compound is represented by [CH2:CHC6F4Z]yRf(ZArZrF)x[ZC6F4CH:CH2]y', where Z = O or S; Rf = aliphatic or aromatic group; y = 1-10, yr' = 0-1; x = 0-200; Ar = MeC6F4RC6F4Me or C6F3MeX; R = direct single bond, CO, SO2, S and O, and X = F, Cl, Br and I.

ACCESSION NUMBER: 2004:534012 CAPLUS  
DOCUMENT NUMBER: 141:90579  
TITLE: Perfluorostyrene compound, and coating solution and optical waveguide device using the compound  
INVENTOR(S): Kim, Ji-hyang; Kim, Jae-il; Kim, Tae-kyun; Lee, Hyung Jong; Han, Seon Gyu  
PATENT ASSIGNEE(S): Zen Photonics Co., Ltd., S. Korea  
SOURCE: U.S. Pat. Appl. Publ., 13 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004127632	A1	20040701	US 2003-616889	20030710
PRIORITY APPLN. INFO.:		KR 2002-40901 A 20020712		

OTHER SOURCE(S): MARPAT 141:90579  
IT 713525-85-0P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); PACT (Reactant or reagent)  
(preparation and copolym.; perfluorostyrene-terminated photopolymerizable compound for core and cladding layers of optical waveguide device)  
RN 713525-85-0 CAPLUS  
CN Benzene, 1,3,5-tris(3-ethenyl-2,4,5,6-tetrafluorophenoxy)- (9CI) (CA INDEX NAME)



L52 ANSWER 3 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Vinyl aromatic compds. P2C:CPG6X5-nZn, are prepared, in which F = H, D or mixture; Z = -YC6X3, -YU[(o-, m-, p-C6X4CP:CP2)]m, -TV and mixts.; Y and U = O, S, NH, or is not present; V and W = monovalent group and a linking group selected from aromatic, polycyclic aromatic (fused-ring), 5- or 6-member heterocyclic aromatic and polycyclic-heterocyclic compds., and derivs. where


2l ring C-H bonds are replaced by C-X bonds; X = F, Cl, CF3 and Rf, and mixts., and Rf = C2-C5 hydrocarbon in which ≥50% C-H bonds are replaced by C-F bonds; m and n = 1-3. The compds., when used as polymeric

materials in optical communications devices result in devices with low power losses.


ACCESSION NUMBER: 2004:286761 CAPLUS  
 DOCUMENT NUMBER: 140:304235  
 TITLE: Halogenated styrene compounds, radiation curable composition, low-absorption-loss polymers and their synthesis  
 INVENTOR(S): Roux, Stephane; Arnoud, Olivier; Moroni, Marc  
 PATENT ASSIGNEE(S): Corning Incorporated, USA  
 SOURCE: Eur. Pat. Appl., 22 pp.  
 CODEN: EPXKDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1403849	A1	20040407	EP 2002-292461	20021004
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, HK, CY, AL, TR, BG, CZ, EE, SK				
US 2004077874	A1	20040422	US 2003-667530	20030922
PRIORITY APPLN. INFO.:			EP 2002-292461	A 20021004


OTHER SOURCE(S): MARPAT 140:304235  
 IT 676515-14-3P  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (halogenated styrene compds. for low-absorption-loss polymers for waveguides)  
 RN 676515-14-3 CAPLUS  
 CN Benzene, 1-ethenyl-3,5-difluoro-2,4,6-tris(pentafluorophenoxy)-, homopolymer (9CI) (CA INDEX NAME)  
 CM 1  
 CRW 676515-05-2  
 CHF C26 H3 F17 O3




IT 676515-04-1P  
 RL: IMF (Industrial manufacture); PUR (Purification or recovery); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
 (halogenated styrene compds. for low-absorption-loss polymers for waveguides)  
 RN 676515-04-1 CAPLUS  
 CN Benzaldehyde, 2,4,6-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



IT 676515-05-2P  
 RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and polymerization; halogenated styrene compds. for low-absorption-loss polymers for waveguides)  
 RN 676515-05-2 CAPLUS  
 CN Benzene, 1-ethenyl-3,5-difluoro-2,4,6-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)





AB The F-containing poly(aryl ethers), showing high heat resistance and mech. strength, contain I units (R1 = C1-150 divalent organic group; Z = divalent organic group, single bond; m = 1-4) and have OH and/or phosphoric acid groups in R1. Solvent-soluble widely-useful engineering plastics are manufactured using compds. containing 2 of phenolic OH groups and ≥1 alc. OH groups as starting materials. Also claimed are ionic conductors, useful for electrolyte membranes in fuel cells, etc., comprising F-containing poly(aryl ethers) having OH, carboxy, and/or PO3H groups and proton conductivity-imparting agents. Thus, 4,4'-bis(2,3,4,5,6-pentafluorobenzoyl) di-Ph ether was copolycond. with Epicure 171N (resin) to give F-containing polyether-polyketone, which was mixed with tungstophosphoric acid and cured to give a film showing electrocond.  $3.2 \times 10^{-5}$  and  $6.4 \times 10^{-6}$  S/cm, at 80 and 140°, resp.


ACCESSION NUMBER: 2004:139363 CAPLUS  
DOCUMENT NUMBER: 140:182769  
TITLE: Fluorine-containing poly(aryl ethers), curable compositions, cured materials, adhesives, and ionic conductors therefrom, and manufacture of solvent-soluble engineering plastics therefor

INVENTOR(S): Akutagawa, Hironobu; Omote, Kazushi; Matsumoto, Takeshi; Nishiji, Ai; Yoshida, Masaya  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 28 pp.  
CODEN: J000AF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004051978	A2	20040219	JP 2003-155624	20030530
PRIORITY APPLN. INFO.:			JP 2002-160397	A 20020531

IT 659720-09-0P  
RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(fluorine-containing poly(aryl ethers) showing good heat resistance useful



AB Method for the control of weeds which comprises applying thereto a herbicidally effective amount of at least one compound which is a 3,5-dicyanophenoxy derivative of formula (I); wherein A is as defined in the description; to novel 3,5-dicyanophenoxy derivs., to new herbicidal compns. containing them, and to processes and intermediates for their preparation

ACCESSION NUMBER: 2003:490952 CAPLUS  
DOCUMENT NUMBER: 139:64820  
TITLE: Preparation of 3,5-dicyanophenoxy derivs. as herbicides

INVENTOR(S): Cornell, Clive; Cramp, Michael Colin; Gingell, Michael; Westaway, Susan  
PATENT ASSIGNEE(S): Bayer CropScience Ltd., UK  
SOURCE: PCT Int. Appl., 58 pp.  
CODEN: PIXXD2


DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003051119	A1	20030626	WO 2002-EP13232	20021125
W: AE, AG, AL, AM, AU, AZ, BA, BB, BR, BY, BZ, CA, CN, CO, CR, CU, DM, DZ, EC, GE, GR, HR, HU, ID, IL, IN, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MA, MD, MG, MK, MN, MX, NO, NZ, OM, PH, PL, RO, RU, SC, SG, SI, TJ, TM, TN, TT, UA, US, UZ, VC, VN, YU, ZA, ZW, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AG, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10161763	A1	20030717	DE 2001-10161763	20011215
PRIORITY APPLN. INFO.:			DE 2001-10161763	A 20011215



OTHER SOURCE(S): MARPAT 139:64820  
IT 548465-86-7P 548465-90-3P 548465-93-6P  
RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation as herbicide)  
RN 548465-86-7 CAPLUS  
CN 1,3-Benzenedicarbonitrile, 5-[3-(3-cyano-5-fluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)

RN 659720-09-9 CAPLUS  
CN Poly[oxy[5-(hydroxymethyl)-1,3-phenylene]oxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene] (2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 1-B




RN 548465-90-3 CAPLUS  
CN 1,3-Benzenedicarbonitrile, 5-[3-(4-cyano-2,3,5,6-tetrafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



RN 548465-93-6 CAPLUS  
CN 1,3-Benzenedicarbonitrile, 5-[3-(3-fluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT




AB The title compds. [I; R1 = (un)substituted alkyl, aryl, heteroaryl, etc.; R2 = H, alkyl; or NR1R2 = 4-8 membered heterocycloaryl; R3 = H, alkyl, halo, etc.; R4 = H, halo, (un)substituted alkyl, aryl; R5 = (un)substituted alkyl, aryl, heteroaryl, heterocycloalkyl; R6 = H, halo, (un)substituted aryl, etc.; X = O, S], useful in the treatment of diseases through the inhibition of sodium ion flux through voltage-dependent sodium channels, were prepared. Thus, amidation of 5-(3,4-dichlorophenyl)-7-trifluoromethylpyrazolo[1,5-a]pyrimidine-2-carboxylic acid with PhCH2NH2 in the presence of BOP in THF containing Et3N afforded 89% I [R1 = CH2Ph; R2 = H; R3 = CF3; R4 = H; R5 = 3,4-Cl2C6H3; R6 = H; X = O]. More particularly, the invention provides pyrazolopyrimidines, compns. and methods that are useful in the treatment of central or peripheral nervous system disorders, particularly pain and chronic pain by blocking sodium channels associated with the onset or recurrence of the indicated conditions. The compds., compns. and methods of the present invention are of particular use for treating neuropathic or inflammatory pain by the inhibition of ion flux through a channel that includes a PN3 subunit. The compds. I were tested in mech. allodynia in vivo assay and in thermal hyperalgesia in vivo assay. Results show that after oral administration the compds. I produce efficacious anti-allodynic effects at doses less than or equal to 100 mg/kg, and that after IV administration the compds. I produce efficacious anti-hyperalgesic effects at doses less than or equal to 30 mg/kg.

ACCESSION NUMBER: 2003:356452 CAPLUS  
DOCUMENT NUMBER: 138:368908  
TITLE: Preparation of pyrazolopyrimidines as sodium channel inhibitors  
INVENTOR(S): Atkinson, Robert Nelson; Gross, Michael Francis; Van Rhee, Michel Albert  
PATENT ASSIGNEE(S): Icaegen, Inc., USA  
SOURCE: PCT Int. Appl., 94 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003037900	A2	20030508	WO 2002-US35171	20021101
WO 2003037900	A3	20041118		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				

L52 ANSWER 6 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MW, MX, MY, NZ, OM, PA, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
RW: GH, GM, KE, LS, MW, ME, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
US 2003176450 A1 20030918 US 2002-286687 20021101  
US 6833371 B2 20041221 US 2001-335874P P 20011101




PRIORITY APPL. INFO.:  
OTHER SOURCE(S): MARPAT 138:368908  
IT 326892-95-9P 326893-03-2P 328088-45-5P  
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
[preparation of pyrazolopyrimidines as sodium channel inhibitors]  
RN 326892-95-9 CAPLUS  
CN Pyrazolo[1,5-a]pyrimidine-2-carboxamide, N-[3,5-bis(4-fluorophenoxy)phenyl]-3-chloro-5-phenyl-7-(trifluoromethyl)- (9CI) (CA INDEX NAME)



RN 326893-03-2 CAPLUS  
CN Pyrazolo[1,5-a]pyrimidine-2-carboxamide, N-[3,5-bis(4-fluorophenoxy)phenyl]-5-(2-thienyl)-7-(trifluoromethyl)- (9CI) (CA INDEX NAME)



RN 328088-45-5 CAPLUS  
CN Pyrazolo[1,5-a]pyrimidine-2-carboxamide, N-[3,5-bis(4-fluorophenoxy)phenyl]-3-chloro-5-(2-thienyl)-7-(trifluoromethyl)- (9CI) (CA INDEX NAME)



II

AB The aromatic diamine compound having a formula I and a polyimide having a repeating unit represented by the following formula II, which has low-temperature adhesion, are prepared, where n is an integer of 3-7, R is independently an atom or a group selected from the group consisting of a H, a halogen atom and a hydrocarbon group, the same or different two hetero atoms selected from N and O bonded to each benzene ring are at the ortho- or meta-positions to each other on at least one benzene ring, and when n is 3, the hetero atoms are at the ortho- or meta-positions to each other on all the benzene rings and Y is a tetravalent organic group.


ACCESSION NUMBER: 2003:172961 CAPLUS  
DOCUMENT NUMBER: 138:222013  
TITLE: Novel aromatic diamine and polyimide  
INVENTOR(S): Kodama, Yoichi; Mori, Minehiro; Nagai, Naoshi; Kawaguchi, Masaru  
PATENT ASSIGNEE(S): Mitsui Chemicals, Inc., Japan  
SOURCE: Eur. Pat. Appl., 24 pp.  
CODEN: EPOXIM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1288191	A2	20030305	EP 2002-19705	20020902
EP 1288191	A3	20030702		


L52 ANSWER 7 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK  
 JP 2003231753 A2 20030819 JP 2002-243952 20020823  
 JP 2004002265 A2 20040108 JP 2002-247130 20020827  
 CN 1403438 A 20030319 CN 2002-142274 20020829  
 US 2003092870 A1 20030515 US 2002-232744 20020903  
 US 6737503 B2 20040518  
 US 2004082754 A1 20040429 US 2003-718532 20031124  
 JP 2001-267218 A 20010904

PRIORITY APPLN. INFO.:  
 JP 2001-332664 A 20011030  
 JP 2002-101107 A 20020403  
 JP 2001-369100 A 20011203  
 JP 2001-369101 A 20011203  
 US 2002-232744 A3 20020903


OTHER SOURCE(S): MARPAT 138:222013  
 IT 500577-29-59 500577-29-79  
 RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation);  
 RACT (Reactant or reagent)  
 (synthesis of novel aromatic diamines and polyimides having  
 low-temperature  
 adherability)  
 RN 500577-27-5 CAPLUS  
 CN Benzene, 1,3-bis(3-fluorophenoxy)- (9CI) (CA INDEX NAME)



RN 500577-29-7 CAPLUS  
 CN Benzene, 1,3-bis[3-(3-fluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



L52 ANSWER 8 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI




AB Preventive or recurrence-suppressive agents for liver cancer containing  
 as the  
 active ingredient thyroid hormone receptor agonists having an effect of  
 inhibiting the expression of liver estrogen sulfotransferase; and usage  
 of  
 the agents. The thyroid hormone receptor agonists are preferably compds.  
 represented by the general formula I (R1 and R2 = alkyl, halogeno, or the  
 like; R3 = hydrogen, alkyl, halogeno, or the like; X = hydroxyl or the  
 like; Y = O, S, CH2, or the like; Z = alkyl, -Q-T (wherein Q = O, CH2,  
 CH(OH), or the like; and T = optionally substituted aryl or the like), or  
 the like; Z = hydrogen, alkoxy, or the like; and A = -NHCO-Y1-CO2R8,  
 -CH2CH(R9)NR10R11, or the like) or pharmaceutically acceptable salts  
 thereof.

ACCESSION NUMBER: 2002:905927 CAPLUS  
 DOCUMENT NUMBER: 138:305  
 TITLE: Preventive or recurrence-suppressive agents for liver  
 cancer  
 INVENTOR(S): Ohnata, Hideki; Hayashi, Morimichi; Kuroda, Junji;  
 Komatsu, Yoshimitsu; Nishimura, Yoshihiro  
 PATENT ASSIGNEE(S): Kissei Pharmaceutical Co., Ltd., Japan  
 SOURCE: PCT Int. Appl., 142 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002094319	A1	20021128	WO 2002-JP4601	20020513
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			JP 2001-149775	A 20010518

OTHER SOURCE(S): MARPAT 138:305  
 IT 373641-22-6P 373641-71-5P  
 RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); SPN  
 (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study);  
 PREP (Preparation); USES (Uses)


L52 ANSWER 8 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 (preventive or recurrence-suppressive agents for liver cancer contg.  
 thyroid hormone receptor agonists)  
 RN 373641-22-6 CAPLUS  
 CN Propanoic acid, 3-[[4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]-3,5-  
 dimethylphenyl]amino]-3-oxo-, ethyl ester (9CI) (CA INDEX NAME)



RN 373641-71-5 CAPLUS  
 CN Propanoic acid, 3-[[4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]-3,5-  
 dimethylphenyl]amino]-3-oxo- (9CI) (CA INDEX NAME)




IT 373642-43-4P 477274-17-2P, 4-(4-Amino-2,6-  
 dimethylphenoxy)-2-(4-fluorophenoxy)phenol  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preventive or recurrence-suppressive agents for liver cancer  
 containing  
 thyroid hormone receptor agonists)  
 RN 373642-43-4 CAPLUS  
 CN Benzene, 4-(2,6-dimethyl-4-nitrophenoxy)-2-(4-fluorophenoxy)-1-  
 (phenylmethoxy)- (9CI) (CA INDEX NAME)




RN 477274-17-2 CAPLUS  
 CN Phenol, 4-(4-amino-2,6-dimethylphenoxy)-2-(4-fluorophenoxy)- (9CI) (CA  
 INDEX NAME)

L52 ANSWER 8 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT




AB Comps. represented by the general formula (I) or pharmacol. acceptable salts thereof [wherein W represents oxygen, sulfur, methylene, CO, SO, or SO2; R represents hydrogen, C1-6 alkyl or aryl-C1-6 alkyl; R1 and R2 represent each C1-3 alkyl, CF3, or halogeno; R3 represents hydrogen, C1-3 alkyl, halogeno, or CF3; Y represents C1-6 alkyl, CF3, 6-oxo-1,6-dihydropyridazin-3-ylmethyl, or -Q-T (wherein Q represents oxygen, methylene, hydroxymethylene, or CO; and T represents optionally substituted aryl or arylmethyl or cycloalkylmethyl optionally containing the ring); and Z represents hydrogen or C1-3 alkoxy or Y and Z are linked together to form tetramethylene] are prepared. These compts. I have excellent effects of lowering neutral fat level and non-HDL cholesterol level in the blood, inhibiting or suppressing the accumulation of neutral fat in the liver and protecting or ameliorating the liver function and, therefore, are useful as preventives or remedies for circulatory diseases such as hyperlipemia, arteriosclerosis, fatty liver, and hepatitis.

Thus, 4-[3-(4-fluorobenzoyl)-4-hydroxyphenoxy]-3,5-dimethylmalonanilic acid Et ester was reduced by NaBH4 in THF at room temperature for 13 h to give 4-[3-[(4-fluorophenyl)hydroxymethyl]-4-hydroxyphenoxy]-3,5-dimethylmalonanilic acid Et ester which was converted into 4-[3-[(4-fluorophenyl)hydroxymethyl]-4-hydroxyphenoxy]-3,5-dimethylmalonanilic acid potassium salt (II). II at 30 nmol/kg twice a day for 2 wk lowered the triglyceride level in liver of male KK-Ay mice from 16.1 (control) to 2.8 mg/l g liver.


ACCESSION NUMBER: 2001:833261 CAPLUS  
DOCUMENT NUMBER: 135:371762  
TITLE: Preparation of malonanilic acid derivatives as preventives or remedies for circulatory disease  
INVENTOR(S): Shiohara, Hiroaki; Nakamura, Tetsuya; Kikuchi, Norihiko; Ohnata, Hideki; Kozumi, Takashi; Kitazawa, Makio  
PATENT ASSIGNEE(S): Kissei Pharmaceutical Co., Ltd., Japan  
SOURCE: PCT Int. Appl., 118 pp.  
CODEN: PIXOXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001085670	A1	20011115	WO 2001-JP3499	20010424
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BE, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO,				




REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS  
FORMAT RECORD. ALL CITATIONS AVAILABLE IN THE RE

OTHER SOURCE(S): MARPAT 135:371762  
IT 373641-22-6P 373641-71-5P  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
[preparation of malonanilic acid derivs. lowering neutral fat level and non-HDL cholesterol level in blood as preventives or remedies for circulatory diseases]  
RN 373641-22-6 CAPLUS  
CN Propanoic acid, 3-[[4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]-3,5-dimethylphenyl]amino]-3-oxo-, ethyl ester (9CI) (CA INDEX NAME)



RN 373641-71-5 CAPLUS  
CN Propanoic acid, 3-[[4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]-3,5-dimethylphenyl]amino]-3-oxo- (9CI) (CA INDEX NAME)




IT 373642-43-4P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
[preparation of malonanilic acid derivs. lowering neutral fat level and non-HDL cholesterol level in blood as preventives or remedies for circulatory diseases]  
RN 373642-43-4 CAPLUS  
CN Benzene, 4-(2,6-dimethyl-4-nitrophenoxy)-2-(4-fluorophenoxy)-1-(phenylmethoxy)- (9CI) (CA INDEX NAME)

L52 ANSWER 10 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
AB The compts. QCO-p-C6H4O-p-C6H4R (Q = 2,3,4,5,6-pentafluorophenyl; R = OH, COO) are manufactured. Thus, 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone (prepared from 2,3,4,5,6-pentafluoro-4'-methoxybenzophenone) was polymerized in dimethylacetamide at 160° to give a polymer showing good solubility in dimethylacetamide and m-cresol, 10% weight loss (in air) temperature 421°.  
ACCESSION NUMBER: 2001:176773 CAPLUS  
DOCUMENT NUMBER: 134:208664  
TITLE: (2,3,4,5,6-pentafluorobenzoyl)diphenyl ethers and fluorine-containing aryl ether ketone polymers having high solubility and heat stability  
INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Casiday, Rachel E.; Fitch, John W., III; Reddy, V. Sreenivasulu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
CODEN: JXOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001064226	A2	20010313	JP 1999-183950	19990629
PRIORITY APPL. INFO.: US 1998-106270 A 19980629				
			JP 1999-180091	A 19990625

IT 213693-17-5P  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
[(pentafluorobenzoyl)phenyl ether as monomers for F-containing aryl ether ketone polymers having high solubility and heat stability]  
RN 213693-17-5 CAPLUS  
CN Poly[oxy-1,3-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene]carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene) (9CI) (CA INDEX NAME)



PAGE 1-A

PAGE 1-B



L52 ANSWER 11 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

AB The compns., useful for elec. insulating materials, contain (p-C6F4COC6H4-Q)QORIm)n (R1 = C6H4-R' (CO-p-C6F4O)2pO; R2 = divalent aryl; X, Y = halo, lower alkyl, alkoxy; q, r = 0-4; m, p = 0, 1). Thus, 2,2-bis(4-hydroxyphenyl)-1,1,1,3,3,3-hexafluoropropane was polymerized with 4,4'-bis(2,3,4,5,6-pentafluorobenzoyl)diphenyl ether to give a polymer showing dielec. constant 3.06 at 25°, 10% weight loss temperature 524° under N, and Tg 174°.

ACCESSION NUMBER: 2001:124290 CAPIUS  
DOCUMENT NUMBER: 134:179345  
TITLE: Low dielectric fluorinated aromatic polyether ketone compositions with good heat resistance  
INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 21 pp.  
CODEN: JKOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001049110	A2	20010220	JP 1999-226981	19990810
JP 3539897	B2	20040707		


PRIORITY APPLN. INFO.: JP 1999-226981 19990810

IT 213693-17-5P  
RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(low dielec. fluorinated aromatic polyether ketone compns. with good heat resistance)


RN 213693-17-5 CAPIUS

CN Poly[oxy-1,3-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylenecarbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



L52 ANSWER 12 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

AB Ketonic di-Ph ethers of p-R1COC6H4OC6H4R2-p type compds. (R1 = 2,3,4,5,6-pentafluorobenzoyl; R2 = OH, pentafluorobenzoyl group) and polyether-polyketone polymers containing tetrafluorophenylene and phenylene groups are provided which have good mech. strength, toughness, elec. property, thermal oxidative stability and solubility. Thus, heating 0.5 g 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone with 0.36 ground K carbonate, 2 mL N-methyl-2-pyrrolidone and 1 mL PhMe at 160° while stirring for 3 h gave a polymer at 85% yield and having viscosity 0.5 g/dL in AcMe2.

ACCESSION NUMBER: 2001:25791 CAPIUS  
DOCUMENT NUMBER: 134:86663  
TITLE: (2,3,4,5,6-Pentafluorobenzoyl)diphenyl ether compound, and fluorine-containing aryl ether ketone polymer  
INVENTOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.; Fitch, John W., III; Reddy, V. Sreenivasulu  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: U.S., 22 pp., Cont.-in-part of U.S. Ser. No. 106,270, abandoned.  
CODEN: USXXGM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6172181 B1		20010109	US 1999-354976	19990716


PRIORITY APPLN. INFO.: MARPAT 134:86663 US 1998-106270 19980629

IT 213693-17-5P  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(manufacture of (2,3,4,5,6-pentafluorobenzoyl)diphenyl ether compound and fluorine-containing aryl ether ketone polymer)

RN 213693-17-5 CAPIUS


CN Poly[oxy-1,3-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylenecarbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A






PAGE 1-B



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT



AB The reaction of 4-(4-CF<sub>3</sub>C<sub>6</sub>F<sub>4</sub>O)C<sub>6</sub>F<sub>4</sub>Li (RFLi) with halogens, dicyanogen, cyanogen halides, and XeF<sub>2</sub> was examined. The halogenated aroms. RFCl, RFB, and RFI were formed upon reaction with Cl<sub>2</sub>, Br<sub>2</sub>, and I<sub>2</sub>, resp. Essentially the same products were isolated in the reaction of RFLi with cyanogen halides. The nitrile RFCN was obtained from the reaction of RFLi with dicyanogen only as a minor product. The reaction of RFLi with XeF<sub>2</sub> resulted in the isolation of RFCNMeEt. All products were identified and characterized by anal. and spectroscopic methods. In addition, the mol. structures of RFCl, RFB, and RFI were determined by x-ray crystallog. Reaction enthalpies for the gas-phase reactions of C<sub>6</sub>F<sub>5</sub>Li, as a model for RFLi, with XCN (X = F, Cl, Br, I) were calculated.

ACCESSION NUMBER: 2000:745014 CAPLUS  
DOCUMENT NUMBER: 134:41943  
TITLE: The reactivity of CF<sub>3</sub>C<sub>6</sub>F<sub>4</sub>OC<sub>6</sub>F<sub>4</sub>Li towards halogens and cyanogen halides  
AUTHOR(S): Klapotke, Thomas M.; Krumm, Burkhard; Polborn, Kurt  
CORPORATE SOURCE: Department Chemie der Univ., München, Germany  
SOURCE: Zeitschrift fuer Anorganische und Allgemeine Chemie (2000), 626(10), 2047-2052  
CODEN: ZNACAB; ISSN: 0044-2313  
PUBLISHER: Wiley-VCH Verlag GmbH  
DOCUMENT TYPE: Journal  
LANGUAGE: German  
OTHER SOURCE(S): CASREACT 134:41943  
IT 312583-57-6P  
RL: BYP (Byproduct); PREP (Preparation)  
(reaction of [(fluoromethyl)fluorophenoxy]fluorophenyllithium with halogens and cyanogen halides)  
RN 312583-57-6 CAPLUS  
CN Benzene, 1,2,4-trifluoro-3,5-bis(2,3,5,6-tetrafluorophenoxy)-6-(trifluoromethyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

GI



AB The title compds. (I) [wherein R<sub>1</sub>-R<sub>3</sub> = independently H, halo, alkyl, CF<sub>3</sub>, CN, OCF<sub>3</sub>, or alkoxy; R<sub>4</sub> = H or (un)substituted alkyl; or R<sub>3</sub> and R<sub>4</sub> together form an (un)substituted carbocyclic ring, (CH<sub>2</sub>)<sub>b</sub>, or a heterocyclic ring, Q(CH<sub>2</sub>)<sub>c</sub> or (CH<sub>2</sub>)<sub>j</sub>Q(CH<sub>2</sub>)<sub>k</sub>; b = 3-7; c = 2-6; j and k = independently 2-6; Q = O, S, or NR<sub>1</sub>; R<sub>5</sub> = F, OH, alkoxy, or carboxy; or

R<sub>4</sub> and R<sub>5</sub> together form a heterocyclic ring; R<sub>6</sub> = H, halo, alkyl, or CF<sub>3</sub>; R<sub>7</sub> = H or alkyl; R<sub>8</sub> = OH, alkoxy, or (un)substituted amino; W = O, S(O)<sub>d</sub>, CH<sub>2</sub>, NH, or N(alkyl); d = 0-2], prodrugs, geometric and optical isomers, and pharmaceutically acceptable salts were prepared as thyroid receptor ligands. Thus, 2',6'-dichloro-4-methoxy-4'-nitrodiphenyl ether was treated with ClSO<sub>2</sub>H and pyrrolidine in two steps to give 1-[5-(2,6-dichloro-4-nitrophenoxy)-2-methoxybenzenesulfonyl]pyrrolidine. Demethylation using BCl<sub>3</sub>, followed by reduction using Pd/C, addition of


di-Et oxalate, and deesterification, yielded II. An in vivo oxygen consumption assay designed to evaluate the efficacy and cardiac effects of tissue-selective thyroid hormone agonists and a thyroid hormone receptor (TRα and TRβ) binding assay for thyromimetic compds. are described (no data). I are useful for the treatment of obesity, hyperlipidemia, glaucoma, cardiac arrhythmia, skin disorders, thyroid disease, hypothyroidism, and related disorders and diseases, such as diabetes mellitus, atherosclerosis, hypertension, coronary heart disease, hypercholesterolemia, depression, and osteoporosis. An anorectic agent or lipase inhibitor may be administered with I to treat these conditions.

ACCESSION NUMBER: 2000:628106 CAPLUS  
DOCUMENT NUMBER: 133:207681  
TITLE: Preparation of 4-(sulfamoylphenoxy)phenyloxamic acids and derivatives as thyroid receptor ligands  
INVENTOR(S): Chiang, Yuan-Ching Phoebe; Dow, Robert Lee  
PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
SOURCE: PCT Int. Appl., 128 pp.  
CODEN: PFXDX2  
DOCUMENT TYPE: Patent


LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000051971	A1	20000908	WO 2000-1B183	20000221
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, NG, TD, TG				
CA 2363145	AA	20000908	CA 2000-2363145	20000221
EP 1157001	A1	20011128	EP 2000-902835	20000221
EP 1157001	B1	20040804		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 2000008701	A	20011226	BR 2000-8701	20000221
TR 200102561	T2	20020221	TR 2001-200102561	20000221
JP 2002538133	T2	20021112	JP 2000-602199	20000221
EE 200100464	A	20021216	EE 2001-464	20000221
NZ 513449	A	20040227	NZ 2000-513449	20000221
AU 727282	B2	20040422	AU 2000-24575	20000221
AT 272609	E	20040815	AT 2000-902835	20000221
US 6326398	B1	20011204	US 2000-514862	20000228
ZA 2001006730	A	20020805	ZA 2001-6730	20010815
HR 2001006633	A1	20021031	HR 2001-633	20010830
NO 2001004217	A	20011011	NO 2001-4217	20010831
BG 105954	A	20020628	BG 2001-105954	20010926
US 2002049226	A1	20020425	US 2001-966467	20010927
US 6545018	B2	20030408		
US 2003114521	A1	20030619	US 2002-324948	20021220
PRIORITY APPLN. INFO.:			US 1999-122292P	P 19990301
			WO 2000-1B183	W 20000221
			US 2000-514862	AJ 20000228


OTHER SOURCE(S): MARPAT 133:207681  
IT 290352-19-1P, N-[3,5-Dichloro-4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]phenyl]oxamic acid ethyl ester 290352-20-4P, N-[3,5-Dichloro-4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]phenyl]oxamic acid 290352-21-5P, N-[4-[3-(4-Fluorophenoxy)-4-hydroxyphenoxy]-3,5-dimethylphenyl]oxamic acid  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of 4-(sulfamoylphenoxy)phenyloxamic acids and derivs. as thyroid receptor ligands by treatment of 4-methoxy-4'-nitrodiphenyl ethers with ClSO<sub>2</sub>H and amines, reduction, and amidation with oxalates)  
RN 290352-19-1 CAPLUS  
CN Acetic acid, [1,3,5-dichloro-4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]phenyl]amino]oxo-, ethyl ester (9CI) (CA INDEX NAME)



RN 290352-20-4 CAPLUS  
CN Acetic acid, [[3,5-dichloro-4-([3-(4-fluorophenoxy)-4-hydroxyphenoxy]phenyl)amino]oxo- (9CI) (CA INDEX NAME)




RN 290352-21-5 CAPLUS  
CN Acetic acid, [[4-[3-(4-fluorophenoxy)-4-hydroxyphenoxy]-3,5-dimethylphenyl]amino]oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

US 6265428 B1 20010724 US 1999-327661 19990608  
US 2001014694 A1 20010816 US 2001-808771 20010314  
PRIORITY APPLN. INFO.: US 1996-583871 B2 19960105  
US 1996-590139 A 19960123  
WO 1997-US366 W 19970103  
US 1999-327661 A1 19990608  
OTHER SOURCE(S): MARPAT 131:322420  
IT 193757-00-5P  
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of phenoxybenzoic acids and phenoxybenzimidinecarboxylic acids for modulation of activity of endothelin)  
RN 193757-00-5 CAPLUS  
CN Benzoic acid, 2-(1,3-benzodioxol-5-yloxy)-6-(4-fluorophenoxy)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 56 THERE ARE 56 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

Ar1-X-Ar3-Y-Ar1 I

AB Methods, compns., and compds. for modulating the activity of an endothelin peptide are provided. The methods use compns. that contain carboxylic acid compds. I (X and Y are selected from groups that include O, S, and NH; and Ar1, Ar2 and Ar3 are independently selected from substituted or unsubstituted groups that include 5 to 6 membered aryl groups and heteroaryl groups that contain one or two heteroatom(s)). Twenty-seven compds. were prepared and claimed. For example, 2-[3,4-(methylenedioxy)phenoxy]-6-(4-methylphenoxy)benzoic acid was prepared in

33 1 yield by the reaction of Na 4-methylphenoxide with Et 2-fluoro-6-[3,4-(methylenedioxy)phenoxy]benzoate followed by deesterification or 4,6-diphenoxy-2-(methylthio)pyrimidine-3-carboxylic acid was prepared in 71 1 yield by the reaction of 4,6-diphenoxy-2-(methylthio)pyrimidine with BuLi and dry ice. The activity of endothelin receptors are modulated by contacting with one or more of the compds. or with compns. containing one or more of the compds. prior to, simultaneously with, or subsequent to contacting the receptors with an endothelin peptide.


ACCESSION NUMBER: 1999:704991 CAPLUS  
DOCUMENT NUMBER: 131:322420  
TITLE: Substituted phenyl compounds and derivatives thereof that modulate the activity of endothelin  
INVENTOR(S): Chan, Ming Fai; Balaji, Vitukudi Narayanaiyengar; Castillo, Rosario Silvestre; Koia, Adam; Raju, Bore Gowda; Wu, Chengde  
PATENT ASSIGNEE(S): Texas Biotechnology Corporation, USA  
SOURCE: U.S., 45 pp., Cont.-in-part of U.S. Ser. No. 583,871, abandoned.  
CODEN: USXXGH  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5977117	A	19991102	US 1996-590139	19960123
WO 9725321	A2	19970717	WO 1997-US366	19970103
WO 9725321	A3	19970912		
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9715324	A1	19970801	AU 1997-15324	19970103
EP 876364	A2	19981111	EP 1997-901420	19970103
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,			

AB Perfluoro(4-phenoxy)-2,5-cyclohexadienone reacts with sodium 4-nitro- and 4-methoxycarbonylphenoxides in glyme at 65°C to give the corresponding 2,4,6-trifluoro-3,5-bis(aryloxy)-4-[2,4,6-trifluoro-3,5-bis(aryloxy)phenoxy]-2,5-cyclohexadienones. Reduction of the latter to phenols, followed by reaction with perfluorotoluene, results in formation of branched polyfluorinated polyphenyl ethers containing NO<sub>2</sub> and CO<sub>2</sub>H functional groups. Reduction of the dinitro polyphenyl ether yields the corresponding diamino derivative. A similar reaction sequence gives rise to a


linear polyphenyl ether, starting from 6-chloro-2,3,4,5,6-pentafluoro-2,4-cyclohexadien-1-one and tetrafluororesorcinol.

ACCESSION NUMBER: 1999:513659 CAPLUS  
DOCUMENT NUMBER: 131:257279  
TITLE: Synthesis of fluorinated polyphenyl ethers by reaction of polyfluorinated cyclohexadienones with substituted phenols  
AUTHOR(S): Kovtonyuk, V. N.; Kobrina, L. S.  
CORPORATE SOURCE: Novosibirsk Institute of Organic Chemistry, Siberian Division, Russian Academy of Sciences, Novosibirsk, 630090, Russia  
SOURCE: Russian Journal of Organic Chemistry (Translation of Zhurnal Organicheskoi Khimii) (1999), 35(1), 74-79  
CODEN: RUOCEQ; ISSN: 1070-4280  
PUBLISHER: MAIK Nauka/Interperiodica Publishing  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 245126-31-2P 245126-34-5P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of fluorinated polyphenyl ethers by reaction of polyfluorinated cyclohexadienones with phenols)  
RN 245126-31-2 CAPLUS  
CN Benzene, 1,3,5-trifluoro-2,4-bis(4-nitrophenoxy)-6-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)




RN 245126-34-5 CAPLUS

L52 ANSWER 16 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 CN Phenol, 3,3'-[2,4,5,6-tetrafluoro-1,3-phenylene]bis(oxy)]bis[2,4,5,6-tetrafluoro- (9CI) (CA INDEX NAME)




IT 245126-32-3P 245126-33-4P 245126-36-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of fluorinated polyphenyl ethers by reaction of polyfluorinated cyclohexadienones with phenols)  
 RN 245126-32-3 CAPLUS  
 CN Benzoic acid, 4,4'-[2,4,6-trifluoro-5-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]-1,3-phenylene]bis(oxy)]bis-, dimethyl ester (9CI) (CA INDEX NAME)




RN 245126-33-4 CAPLUS  
 CN Benzenamine, 4,4'-[2,4,6-trifluoro-5-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]-1,3-phenylene]bis(oxy)]bis- (9CI) (CA INDEX NAME)

L52 ANSWER 16 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)





RN 245126-36-7 CAPLUS  
 CN Benzene, 1,2,3,5-tetrafluoro-4,6-bis[2,4,5,6-tetrafluoro-3-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy)phenoxy]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI



AB Title aromatic polyamide with good heat-resistance, low dielec. constant, colorlessness and transparency, processability and thermal stability is composed of 1 or 2 repeating units selected from the group described by a structural formula of (I), where R1 is F, Cl-3 alkyl or fluorinated alkyl, Cl-3 alkoxy, Ph, phenoxy, benzyl, naphthyl or monovalent group having formula of (a) or (b) (R2: halogen, Cl-3 alkyl, alkoxy, halogenated alkyl or halogenated alkoxy, p = 1-5), n is an integer of 1-4, i is an integer of 0-3, A is a group described by the formula of (c) or (d) (Ar1 and Ar2: substituted aryl groups), while A is described by b, R1 is not F or fluorinated alkyl group. The polyamide is prepared by polycondensation of fluorinated alkyl-substituted bis(aminophenoxy)benzene with aromatic dicarboxylic acid derivative. Thus, 1,3-bis(3-aminophenoxy)-4-trifluoromethylbenzene 18.02 g was polymerized with terephthaloyl chloride 10.13 g to give a polyamide with yield of 97.1, logarithmic viscosity of 1.11 dl/g, glass transition temperature of 213°, and 5% weight reducing temperature of 508°.

ACCESSION NUMBER: 1999:253765 CAPLUS  
 DOCUMENT NUMBER: 130:338828  
 TITLE: Preparation of fluorine-containing aromatic polyamide resin for organic optical components  
 INVENTOR(S): Yamashita, Wataru; Yoshimura, Tomoyoshi; Shibuya, Atsushi; Sakata, Yoshihiro; Oikawa, Hideaki; Takuma, Katsuke; Ohta, Masahiro

L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PATENT ASSIGNEE(S): Mitsui Chemicals Inc., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 62 pp.  
 CODEN: JKKKAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 11106503	A2	19990420	JP 1997-226052	19970822
JP 11071316	A2	19990316	JP 1997-196009	19970722
US 6103860	A	20000815	US 1999-332534	19990614
PRIORITY APPLN. INFO.:				JP 1996-228551 A 19960829
				JP 1997-159422 A 19970617
				JP 1997-159424 A 19970617
				JP 1997-212855 A 19970807
				JP 1996-201825 A 19960731
				JP 1996-204614 A 19960802
				JP 1996-204615 A 19960802
				JP 1996-331831 A 19961212
				JP 1996-331832 A 19961212
				JP 1996-331833 A 19961212
				JP 1997-138355 A 19970528
				JP 1997-138356 A 19970528
				JP 1997-138357 A 19970528
				JP 1997-138358 A 19970528
				JP 1997-138359 A 19970528
				JP 1997-159421 A 19970617
				JP 1997-159423 A 19970617
				US 1997-917387 A3 19970826

IT 203808-02-0P 203808-05-3P 223791-44-4P  
 223791-45-5P 223791-46-6P  
 RL: DEV (Device component use); IMF (Industrial manufacture); PRP (Properties); PREP (Preparation); USES (Uses)  
 (preparation and properties of fluorine-containing aromatic polyamide resin for organic optical components)  
 RN 203808-02-0 CAPLUS  
 CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with 3,3'-[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-fluorobenzenamine]

L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
(9CI) (CA INDEX NAME)


CM 1

CRN 203807-91-4  
CMF C19 H13 F5 N2 O2



CM 2


CRN 2351-37-3  
CMF C14 H8 C12 O2



RN 203808-05-3 CAPLUS  
CN Benzoyl chloride,  
4,4'-(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)bis-  
polymer with 3,3'-[2,4,5-trifluoro-6-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine]  
(9CI) (CA INDEX NAME)


CM 1

CRN 203807-99-2  
CMF C21 H4 F18 N2 O2



CM 2


L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 223791-45-5 CAPLUS  
CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with  
1,4-benzenedicarbonyl dichloride and 3,3'-[2,4,5-trifluoro-6-  
(trifluoromethyl)-1,3-phenylene]bis(oxy)bis[2,4,5-trifluoro-6-  
(trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)


CM 1

CRN 203807-99-2  
CMF C21 H4 F18 N2 O2



CM 2

CRN 2351-37-3  
CMF C14 H8 C12 O2




CM 3

CRN 100-20-9  
CMF C8 H4 C12 O2

L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


CRN 1102-92-7  
CMF C17 H8 C12 F6 O2



RN 223791-44-4 CAPLUS  
CN Benzoyl chloride,  
4,4'-(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)bis-  
polymer with 3,3'-[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)bis[4-  
fluorobenzenamine] and 3,3'-[4-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)bis[6-fluorobenzenamine] (9CI) (CA INDEX NAME)


CM 1

CRN 203807-97-0  
CMF C19 H13 F5 N2 O2



CM 2


CRN 203807-94-7  
CMF C19 H13 F5 N2 O2



CM 3

CRN 1102-92-7  
CMF C17 H8 C12 F6 O2


L52 ANSWER 17 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 223791-46-6 CAPLUS  
CN 1,3-Benzenedicarbonyl dichloride, 4-(trifluoromethyl)-, polymer with  
1,3-benzenedicarbonyl dichloride, 3,3'-[4-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)bis[5-fluorobenzenamine] and  
3,3'-[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)bis[5-(trifluoromethyl)benzenamine] (9CI) (CA  
INDEX NAME)


CM 1

CRN 203807-91-4  
CMF C19 H13 F5 N2 O2




CM 2

CRN 161824-61-9  
CMF C21 H13 F9 N2 O2




CM 3


CRN 1483-49-4  
CMF C9 H3 C12 F3 O2




CM 4  
CRN 99-63-8  
CMF C8 H4 Cl2 O2




IT 223791-39-7P 223791-40-0P 223791-41-1P  
223791-42-2P  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(preparation and properties of fluorine-containing aromatic polyamide  
resin for  
organic optical components)  
RN 223791-39-7 CAPIUS  
CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with  
1,4-benzenedicarbonyl dichloride and 3,3'-[[4-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)]bis[6-fluorobenzenamine] (9CI) (CA INDEX NAME)  
CM 1  
CRN 203807-94-7  
CMF C19 H13 F5 N2 O2




CM 2  
CRN 2351-37-3




CM 3  
CRN 99-63-8  
CMF C8 H4 Cl2 O2




RN 223791-41-1 CAPIUS  
CN 1,3-Benzenedicarbonyl dichloride, 4-(trifluoromethyl)-, polymer with  
3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[4-fluorobenzenamine]  
and 3,3'-[[2,4,5-trifluoro-6-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)]bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine]  
(9CI) (CA INDEX NAME)  
CM 1  
CRN 203807-99-2  
CMF C21 H4 F18 N2 O2




CM 2  
CRN 203807-97-0  
CMF C19 H13 F5 N2 O2




CM 3  
CRN 100-20-9  
CMF C8 H4 Cl2 O2




RN 223791-40-0 CAPIUS  
CN 1,3-Benzenedicarbonyl dichloride, polymer with  
3,3'-[[4-(trifluoromethyl)-  
1,3-phenylene]bis(oxy)]bis[5-fluorobenzenamine] and 3,3'-[[4-(  
trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-(  
trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)  
CM 1  
CRN 203807-91-4  
CMF C19 H13 F5 N2 O2




CM 2  
CRN 161824-61-9  
CMF C21 H13 F9 N2 O2




CM 3  
CRN 1483-49-4  
CMF C9 H3 Cl2 F3 O2



RN 223791-42-2 CAPIUS  
CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with  
3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-  
fluorobenzenamine], 3,3'-[[4-(trifluoromethyl)-1,3-  
phenylene]bis(oxy)]bis[6-fluorobenzenamine] and 4,4'-[2,2,2-trifluoro-1-(  
trifluoromethyl)ethylidene]bis[benzoyl chloride] (9CI) (CA INDEX NAME)  
CM 1  
CRN 203807-94-7  
CMF C19 H13 F5 N2 O2




CM 2  
CRN 203807-91-4  
CMF C19 H13 F5 N2 O2



CM 3

CRN 2351-37-3


CMF C14 H8 C12 O2



CM 4

CRN 1102-92-7

CMF C17 H8 C12 F6 O2




IT 203807-92-5P 203807-93-6P 203807-95-6P  
 203807-96-9P 203807-98-1P 203808-00-8P  
 RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
 (preparation of fluorine-containing aromatic polyamide resin for  
 organic optical  
 components)  
 RN 203807-92-5 CAPLUS  
 CM Benzoyl chloride,  
 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis-  
 , polymer with 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-  
 fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-91-4

CMF C19 H13 F5 N2 O2




RN 203807-95-8 CAPLUS  
 CM (1,1'-Biphenyl)-4,4'-dicarbonyl dichloride, polymer with  
 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[6-fluorobenzenamine]  
 (9CI) (CA INDEX NAME)

CM 1

CRN 203807-94-7


CMF C19 H13 F5 N2 O2



CM 2

CRN 2351-37-3

CMF C14 H8 C12 O2




RN 203807-96-9 CAPLUS  
 CM 1,4-Benzenedicarbonyl dichloride, polymer with  
 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[6-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-94-7


CMF C19 H13 F5 N2 O2



CM 2

CRN 1102-92-7

CMF C17 H8 C12 F6 O2




RN 203807-93-6 CAPLUS  
 CM 1,3-Benzenedicarbonyl dichloride, polymer with  
 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-91-4


CMF C19 H13 F5 N2 O2



CM 2

CRN 99-63-8


CMF C8 H4 C12 O2



CM 2

CRN 100-20-9

CMF C8 H4 C12 O2




RN 203807-98-1 CAPLUS  
 CM [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with  
 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[4-fluorobenzenamine]  
 (9CI) (CA INDEX NAME)

CM 1

CRN 203807-97-0


CMF C19 H13 F5 N2 O2



CM 2

CRN 2351-37-3


CMF C14 H8 C12 O2



RN 203808-00-8 CAPIUS  
CN 1,3-Benzenedicarbonyl dichloride, 4-(trifluoromethyl)-, polymer with 3,3'-[1,2,4,5-trifluoro-6-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)



CM 1

CRN 203807-99-2  
CHF C21 H4 F18 N2 O2




CM 2

CRN 1483-49-4  
CHF C9 H3 Cl2 F3 O2



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT



AB The triazole derivs, I [T = (un)substituted triazolone, etc.; U = halo or (halo)alkyl; V = H, halo, alkyl, CN, NO2 or alkoxy; YZ = group of ≥5 atoms (C, N, O S Si and/or Ge), provided that ≥2 atoms are C], their N-oxides and/or salts, are prepared and enriched in the more

active isomer with respect to the relative positions of U, T and YZ. The enantiomerically-enriched compds. control fungi and arthropods.

ACCESSION NUMBER: 1999:184085 CAPIUS  
DOCUMENT NUMBER: 130:219484  
TITLE: Preparation of enantiomerically-enriched triazolone derivatives as fungicides and arthropodicides  
INVENTOR(S): Brown, Richard James; Casalnuovo, Albert Loren; Chan, Dominic Ming-Tak  
PATENT ASSIGNEE(S): E. I. Du Pont de Nemours & Co., USA  
SOURCE: PCT Int. Appl., 213 pp.  
CODEN: PIXXK2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9911129	A1	19990311	WO 1998-US17747	19980827
W: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GR, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9892065	A1	19990322	AU 1998-92065	19980827
PRIORITY APPLN. INFO.:			US 1997-57917P	P 19970904
			WO 1998-US17747	W 19980827


OTHER SOURCE(S): MARPAT 130:219484  
IT 221117-66-4P  
RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
[preparation as as fungicide and arthropodicide]  
RN 221117-66-4 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-(2-{3-(2-fluorophenoxy)phenoxy}-6-methylphenyl)-2,4-dihydro-5-methoxy-2-methyl-, stereoisomer (9CI) (CA INDEX NAME)


L52 ANSWER 19 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN  
AB Fluoropolymer-polyether-polyketones were obtained by homopolycondensation of 4-hydroxy-4'-(pentafluorobenzoyl)diphenyl ether and by copolycondensation of 4,4'-bis(pentafluorobenzoyl)diphenyl ether with benzenediols or bisphenols. The polymers have very good heat resistance, with 10% weight loss temps. ≥500°.

ACCESSION NUMBER: 1998:532324 CAPIUS  
DOCUMENT NUMBER: 129:260978  
TITLE: New polymers derived from 2,3,4,5,6-pentafluorobenzoic acid  
AUTHOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.; Fitch, John W., III; Reddy, V. Sreenivasulu; Sakaguchi, Yoshimitu  
CORPORATE SOURCE: Faculty of Environmental Science and Technology, Okayama University, Okayama, 700-8530, Japan  
SOURCE: Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1998), 39(2), 790-791  
CODEN: ACPPAY; ISSN: 0032-3934  
PUBLISHER: American Chemical Society, Division of Polymer Chemistry  
DOCUMENT TYPE: Journal  
LANGUAGE: English


IT 213693-17-5P, 1,3-Benzenediol-4,4'-bis(pentafluorobenzoyl)diphenyl ether copolymer, SRU  
RL: SPN (Synthetic preparation); PREP (Preparation)  
[preparation of thermally stable fluoropolymer-polyether-polyketones]  
RN 213693-17-5 CAPIUS  
CN Poly[oxy-1,3-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylenecarbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A





REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT



AB Title compds. I and their N-oxides and agriculturally suitable salts are disclosed [wherein E = (un)substituted 1,2-phenylene, naphthalene or heterocyclyl; A = O, S, N, NR3 or CR4; G = C or N; when G is C, then A is O, S or NR3 and the floating double bond is attached to G; and when G is N, then A is N or CR4 and the floating double bond is attached to A; W = O, S, NH, N(C1-C6 alkyl) or NO(C1-C6 alkyl); X = H, OR1, SOR1, halo, C1-C6 alkyl, C1-C6 haloalkyl, C3-C6 cycloalkyl, cyano, NH2, NHR1, N(C1-C6 alkyl)R1, NH(C1-C6 alkoxy) or N(C1-C6 alkoxy)R1; R2 = H, C1-C6 alkyl, C1-C6 haloalkyl, C2-C6 haloalkyl, C2-C6 alkenyl, C2-C6 haloalkenyl, C2-C6 alkynyl, C2-C6 haloalkynyl, C3-C6 cycloalkyl, C2-C6 alkylcarbonyl, C2-C6 alkoxycarbonyl, hydroxy, C1-C2 alkoxy, or acetyloxy; R1 = (halo)alkyl, (halo)alkenyl, etc.; R3 = H, (halo)alkyl, etc.; Y = O, CO, SO, etc.; Z = (un)substituted alkyl, alkenyl or alkynyl, R4 = H, halo, alkyl, etc.; m = 0, 1 or 2]. Claims cover methods of arthropod and fungal control, novel compds., arthropodocidal and fungicidal compns., and novel intermediates. Approx. 1000 invention compds. were prepared. For instance, 5-chloro-2,4-dihydro-4-(2-methoxyphenyl)-2-methyl-3H-1,2,4-triazol-3-one (preparation given) underwent a sequence of cleavage of the Me ether


with BB3, methoxylation of the chloride with NaOMe, and etherification of the phenolic hydroxy group with 5-chloro-3-[3,5-bis(trifluoromethyl)phenyl]-1,2,4-thiadiazole, to give title compound II. Selected I were active in screens against *Erysiphe graminis*, *Pyricularia oryzae*, *Spodoptera frugiperda*, *Tetranychus urticae*, and a variety of other standard pests.

ACCESSION NUMBER: 1998:385479 CAPIUS  
DOCUMENT NUMBER: 129:54375  
TITLE: Arthropodocidal and fungicidal cyclic amides [triazolones] and their preparation, use, and compositions  
INVENTOR(S): Brown, Richard James; Chan, Dominic Ming-Tak; Howard, Michael Henry, Jr.; Daniel, Dillon Jancey; Clark, David  
PATENT ASSIGNEE(S): Alan; Selby, Thomas Paul  
SOURCE: E. I. Du Pont de Nemours & Co., USA  
CODEN: FIMX02  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2

## PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9823155	A1	19980604	WO 1996-US18916	19961126
W: JP, KR				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
ZA 9709943	A	19990505	ZA 1997-9943	19971105
WO 9823156	A1	19980604	WO 1997-US21944	19971125
W: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9854633	A1	19980622	AU 1998-54633	19971125
EP 944314	A1	19990929	EP 1997-948597	19971125
R: CH, DE, DK, ES, FR, GB, IT, LI, NL, IE				
BR 9713415	A	20000418	BR 1997-13415	19971125
JP 2001506984	T2	20010529	JP 1998-524889	19971125
MX 9904789	A	20000131	MX 1999-4789	19990524
KR 2000057254	A	20000915	KR 1999-704639	19990526
PRIORITY APPL. INFO.:				
			WO 1996-US18916	A 19961126
			US 1996-33614P	P 19961219
			US 1997-48844P	P 19970606
			WO 1997-US21944	W 19971125

OTHER SOURCE(S): MWPAT 129:54375  
IT 186979-62-4P 186979-63-5P 186979-65-7P  
186980-15-4P 186980-17-6P 186980-20-1P  
186980-51-8P 186980-52-9P 186980-53-0P  
186981-79-3P 186981-82-8P 186981-83-9P  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation as arthropodicide and fungicide)  
RN 186979-62-4 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




## RN 186979-63-5 CAPIUS

CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)-2-nitrophenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




## RN 186979-65-7 CAPIUS

CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2-fluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



## RN 186980-15-4 CAPIUS


CN Benzonitrile, 2-[3-[2-(1,5-dihydro-3-methoxy-1-methyl-5-oxo-4H-1,2,4-triazol-4-yl)phenoxy]phenoxy]-6-fluoro- (9CI) (CA INDEX NAME)




## RN 186980-17-6 CAPIUS

CN Benzonitrile, 2-[3-[2-(1,5-dihydro-3-methoxy-1-methyl-5-oxo-4H-1,2,4-triazol-4-yl)phenoxy]phenoxy]-3-fluoro- (9CI) (CA INDEX NAME)







RN 186980-20-1 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,5-difluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




RN 186980-51-8 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(3-fluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




RN 186980-52-9 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,3-difluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




RN 186981-83-9 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,6-difluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT



RN 186980-53-0 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,4-difluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186981-79-3 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2-fluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186981-82-8 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)

AB A series of bis(ether amines), primarily 1,2-bis(4-aminophenoxy)benzenes but including 1,3- and 1,4-bis(4-aminophenoxy)benzenes, with fluoro and alkyl substituents were synthesized. These diamines were prepared by F displacement with 4-FC6H4NO<sub>2</sub> or its derivs., or NO<sub>2</sub> displacement with 1,4-C6H4(NO<sub>2</sub>)<sub>2</sub>, and various phenylenediols and their alkyl or fluoro derivs. The resulting bis(ether nitro) compds. were reduced to the corresponding bis(ether amines).

ACCESSION NUMBER: 1998:376601 CAPLUS

DOCUMENT NUMBER: 129:135942

TITLE: Methyl- and fluoro-substituted bis(4-aminophenoxy)benzenes. A convenient method of synthesis

AUTHOR(S): Eastmond, G. C.; Paprotny, J.  
CORPORATE SOURCE: Donnan Laboratories, University Liverpool, Liverpool, L69 7ZD, UK

SOURCE: Synthesis (1998), (6), 894-898

CODEN: SYNTBF; ISSN: 0039-7881

PUBLISHER: Georg Thieme Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 129:135942


IT 210492-61-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of methylated and fluorinated [(aminophenoxy)phenoxy]benzenamines)

RN 210492-61-8 CAPLUS

CN Benzene, 1,3-bis(2-fluoro-4-nitrophenoxy)- (9CI) (CA INDEX NAME)




IT 210492-62-9P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of methylated and fluorinated [(aminophenoxy)phenoxy]benzenamines)

RN 210492-62-9 CAPLUS

CN Benzenamine, 4,4'-[1,3-phenylenebis(oxy)]bis[3-fluoro- (9CI) (CA INDEX NAME)






AB The title compds. (I: E = (un)substituted 1,2-phenylene; A = O, S, N, NR5, CR6; G = C, N (provided that when G = C, then A = O, S, NR5 and the floating double bond is attached to G; and when G = N, then A = N, CR6 and the floating double bond is attached to A); W = O, S, NH, N(C1-6 alkyl), NO(C1-6 alkyl); X = OR1, S(O)NR1, halo; Y = O, S(O)n, NR7, etc.; Z = substituted Ph, pyrimidinyl, triazinyl; R1 = C1-6 alkyl, C1-6 haloalkyl, C2-6 alkenyl, etc.; R2 = H, C1-6 alkyl, C1-6 haloalkyl, etc.; R5 = H, C1-6 alkyl, C1-6 haloalkyl, etc.; R6 = H, halo, C1-6 alkyl, etc.), useful for controlling plant diseases caused by fungal plant pathogens, were prepared

Thus, 6-step synthesis of the title compound II, which showed 100% control against *Erysiphe graminis* f. sp. *tritici* and *Puccinia recondita* at 500 g/ha, is described.  
ACCESSION NUMBER: 1998:323238 CAPLUS  
DOCUMENT NUMBER: 129:4664  
TITLE: Preparation of fungicidal cyclic amides  
INVENTOR(S): Walker, Michael Paul  
PATENT ASSIGNEE(S): E. I. Du Pont de Nemours & Co., USA; Walker, Michael Paul  
SOURCE: PCT Int. Appl., 60 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9820003	Al	19980514	WO 1997-US17608	19971001
W: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9746603	Al	19980529	AU 1997-46603	19971001
EP 937051	Al	19990825	EP 1997-945385	19971001
R: DE, ES, FR, GB, IT				

AB The title compds. are new low application rate post-emergence herbicides with safety on broadleaf crops. The paper describes the elaboration of existing chemistries which led to the discovery of the aryloxybenzenes as well as the synthetic methodologies which were developed.

Structure-activity relationships are given.  
ACCESSION NUMBER: 1998:294658 CAPLUS  
DOCUMENT NUMBER: 129:13444  
TITLE: Synthesis and herbicidal activity of bisaryloxybenzenes: a new structural class of protox inhibitors derived from N-phenylbenzotriazoles  
AUTHOR(S): Crews, A. D.; Condon, M. E.; Gill, S. D.; Karp, G. M.; Manfredi, M. C.; Birk, J. H.  
CORPORATE SOURCE: Cyanamid Agric. Res. Cent., Am. Cyanamid Co., Princeton, NJ, 08543-0400, USA  
SOURCE: ACS Symposium Series (1998), 686(Synthesis and Chemistry of Agrochemicals VI), 48-54  
CODEN: ACSMCH; ISSN: 0097-6156  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 165379-64-6P  
RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation as herbicide)  
RN 165379-64-6 CAPLUS  
CN Acetic acid, [2-[5-(2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy)-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

BR 9712713	A	19991026	BR 1997-12713	19971001
CN 1242767	A	20000126	CN 1997-181160	19971001
JP 2001503424	T2	20010313	JP 1998-521383	19971001
ZA 9708958	A	19990407	ZA 1997-8958	19971007
MX 9904066	A	20000131	MX 1999-4066	19990430
KR 2000052948	A	20000825	KR 1999-703821	19990430
PRIORITY APPLN. INFO.:				US 1996-29965P P 19961101
				WO 1997-US17608 W 19971001

OTHER SOURCE(S): MARPAT 129:4664  
IT 207504-84-5P  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of fungicidal cyclic amides)  
RN 207504-84-5 CAPLUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,4-difluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT


AB The title polymer is prepared by condensing 21 fluorine-containing aromatic ether diamine with 21 dicarboxylic acid (derivative). This aromatic polyamide has a low permittivity and birefringence, excellent transparency, processability, mech. strength, and thermal stability. Thus, a film having low birefringence and heat stability was molded from the copolymer of 1,3-bis(3-aminophenoxy)-4-trifluoromethylbenzene and terephthaloyl dichloride.

ACCESSION NUMBER: 1998:175770 CAPLUS  
DOCUMENT NUMBER: 128:205262  
TITLE: An optical fluorine-containing aromatic polyether-polyamide or polyimide  
INVENTOR(S): Yamashita, Wataru; Yoshimura, Tomomi; Shibuya, Atsushi; Sakata, Yoshihiro; Oikawa, Hideaki; Takuma, Katsuke; Ohta, Masahiro  
PATENT ASSIGNEE(S): Mitsui Toatsu Chemicals, Inc., Japan  
SOURCE: Eur. Pat. Appl., 58 pp.  
CODEN: EPXKDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 3  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 826717	Al	19980304	EP 1997-306601	19970828
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
US 5990261	A	19991123	US 1997-917387	19970826
CN 1184123	A	19980610	CN 1997-120637	19970829
US 6103860	A	20000815	US 1999-332534	19990614
PRIORITY APPLN. INFO.:				JP 1996-228551 A 19960829
				US 1997-917387 A3 19970826

IT 203807-92-5P 203807-93-6P 203807-95-6P  
203807-96-9P 203807-98-1P 203808-00-6P  
203808-01-9P 203808-02-0P 203808-05-3DP,  
reaction products with benzoyl chloride 203808-08-6DP, reaction products with benzoyl chloride 203808-09-7DP, reaction products with benzoyl chloride  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation) (fluorine-containing aromatic polyether-polyamide or polyimide manufacture for film having low permittivity and birefringence, excellent transparency, processability, mech. strength, and thermal stability)  
RN 203807-92-5 CAPLUS  
CN Benzoyl chloride,  
4,4'-(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)bis-, polymer with 3,3'-[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)bis[5-fluorobenzenamine] (9CI) (CA INDEX NAME)


CM 1  
CRN 203807-91-4  
CMF C19 H13 F5 N2 O2



CM 2

CRN 1102-92-7

CMF C17 H8 C12 F6 O2




RN 203807-93-6 CAPLUS

CN 1,3-Benzenedicarbonyl dichloride, polymer with 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-91-4


CMF C19 H13 F5 N2 O2



CM 2

CRN 99-63-8


CMF C8 H4 C12 O2



CM 2

CRN 100-20-9

CMF C8 H4 C12 O2



RN 203807-98-1 CAPLUS


CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with

3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[4-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-97-0


CMF C19 H13 F5 N2 O2



CM 2

CRN 2351-37-3

CMF C14 H8 C12 O2



RN 203807-95-8 CAPLUS


CN [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with

3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[6-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-94-7


CMF C19 H13 F5 N2 O2



CM 2

CRN 2351-37-3

CMF C14 H8 C12 O2



RN 203807-96-9 CAPLUS


CN 1,4-Benzenedicarbonyl dichloride, polymer with

3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[6-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-94-7

CMF C19 H13 F5 N2 O2




RN 203808-00-8 CAPLUS

CN 1,3-Benzenedicarbonyl dichloride, 4-(trifluoromethyl)-, polymer with 3,3'-[[2,4,5-trifluoro-6-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 203807-99-2


CMF C21 H4 F18 N2 O2



CM 2

CRN 1483-49-4

CMF C9 H3 C12 F3 O2




RN 203808-01-9 CAPLUS

CN 1,3-Benzenedicarbonyl dichloride, 4-(trifluoromethyl)-, polymer with


3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[4-fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

L52 ANSWER 24 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 CRN 203807-97-0  
 CMF C19 H13 F5 N2 O2




CM 2  
 CRN 1483-49-4  
 CMF C9 H3 C12 F3 O2




RN 203808-02-0 CAPIUS  
 CM [1,1'-Biphenyl]-4,4'-dicarbonyl dichloride, polymer with  
 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[5-fluorobenzenamine]  
 (9CI) (CA INDEX NAME)

CM 1  
 CRN 203807-91-4  
 CMF C19 H13 F5 N2 O2




CM 2  
 CRN 2351-37-3

L52 ANSWER 24 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 CRN 203807-94-7  
 CMF C19 H13 F5 N2 O2




CM 2  
 CRN 1102-92-7  
 CMF C17 H8 C12 F6 O2




RN 203808-09-7 CAPIUS  
 CM 1,4-Benzenedicarbonyl dichloride, polymer with 3,3'-[[2,4,5-trifluoro-6-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)

CM 1  
 CRN 203807-99-2  
 CMF C21 H4 F18 N2 O2




CM 2  
 CRN 100-20-9  
 CMF C8 H4 C12 O2

L52 ANSWER 24 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 CMF C14 H8 C12 O2




RN 203808-05-3 CAPIUS  
 CM Benzoyl chloride,  
 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis-  
 , polymer with 3,3'-[[2,4,5-trifluoro-6-(trifluoromethyl)-1,3-  
 phenylene]bis(oxy)]bis[2,4,5-trifluoro-6-(trifluoromethyl)benzenamine]  
 (9CI) (CA INDEX NAME)

CM 1  
 CRN 203807-99-2  
 CMF C21 H4 F18 N2 O2




CM 2  
 CRN 1102-92-7  
 CMF C17 H8 C12 F6 O2



RN 203808-08-6 CAPIUS  
 CM Benzoyl chloride,  
 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis-  
 , polymer with 3,3'-[[4-(trifluoromethyl)-1,3-phenylene]bis(oxy)]bis[6-  
 fluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

L52 ANSWER 24 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L52 ANSWER 25 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Mixing a resin and a certain phthalocyanine compound having the ability  
 to  
 absorb near IR rays, provides a resin composition which is useful as a  
 heat  
 radiation-shielding material which is semi-transparent or transparent for  
 visible light but blocks heat rays. C black may also be added to enhance  
 the heat-shielding effect. Thus, polycarbonate containing 0.003%  
 VOPc(BuNH)8F8 [Pc = phthalocyanine; substituent  
 octakis(butylamino)octafluorophthalocyanine] was molded into a film  
 having transmittance and heat transmittance (JIS R-3106) 78.9 and 62.4%  
 vs. 89.2 and 84.6, resp., for polycarbonate without the IR absorber.


ACCESSION NUMBER: 1998:79369 CAPLUS  
 DOCUMENT NUMBER: 128:141505  
 TITLE: Compositions for shields for heat radiation  
 INVENTOR(S): Kaleda, Osamu; Yodoshi, Takashi; Morita, Ken;  
 Matsuura, Michio  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Japan  
 SOURCE: U.S., 15 pp., Cont.-in-part of U.S. Ser. No. 180,488,  
 abandoned.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5712332	A	19980127	US 1995-478739	19950607
			JP 1993-4326	A 19930113
PRIORITY APPLN. INFO.:			US 1994-180488	B2 19940112


OTHER SOURCE(S): MURPAT 128:141505  
 IT 163464-04-4  
 RL: MOA (Modifier or additive use); USES (Uses)  
 (polymer comps. containing certain phthalocyanines for shields for  
 heat  
 radiation)  
 RN 163464-04-4 CAPLUS  
 CH Zinc, [N,N',N'',N'''-tetrahexyl-1,3,4,8,10,11,15,17,18,22,24,25-  
 dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-  
 tetraminato(2-)-κN29,κN30,κN31,κN32]-, (SP-4-1)-  
 (9CI) (CA INDEX NAME)

L52 ANSWER 25 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A




PAGE 2-A




L52 ANSWER 25 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 3-A




PAGE 4-A



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR  
 THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L52 ANSWER 26 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI



AB The title compds. Ar2-X-Ar3-Y-Ar1 [I: X, Y = O, S, NH, etc.; Ar1, Ar2 =  
 aryl and heteroaryl containing one ring or 2-3 fused rings; Ar3 = aryl,  
 heteroaryl], useful in the treatment of hypertension, cardiovascular  
 disease, asthma, pulmonary hypertension, inflammatory diseases,  
 ophthalmol. disease, menstrual disorders, obstetric conditions, wounds,  
 gastroenteric disease, renal failure, immunosuppressant-mediated renal  
 vasoconstriction, erythropoietin-mediated vasoconstriction endotoxin  
 shock, anaphylactic shock and hemorrhagic shock, were prepared. Thus,  
 reaction of Et 2,6-difluorobenzoate and sodium  
 3,4-methylenedioxyphenoxide  
 in DMSO followed by reaction of the resulting Et 2-fluoro-6-[3,4-  
 (methylenedioxy)phenoxy]benzoate with sodium 4-methoxyphenoxide in DMSO,  
 and hydrolysis of the ester with NaOH/EtOH afforded the title compound  
 II.


Almost all of the compds. I have an IC50 of less than 10 μM and many  
 have an IC50 less than about 1 μM for either or both of the ETA and ETB  
 receptors.

ACCESSION NUMBER: 1997:564939 CAPLUS  
 DOCUMENT NUMBER: 127:161816  
 TITLE: Preparation of aryl- and/or heteroaryl-substituted  
 benzoic acids as endothelin antagonists and/or  
 agonists  
 INVENTOR(S): Chan, Ming Fai; Balaji, Vitukudi Narayanaiyengar;  
 Castillo, Rosario Silverat; Kols, Adam; Raju, Bore  
 Gowda; Wu, Chengde  
 PATENT ASSIGNEE(S): Texas Biotechnology Corp., USA  
 SOURCE: PCT Int. Appl., 136 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9725321	A2	19970717	WO 1997-US366	19970103
WO 9725321	A3	19970912		
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR,				


L52 ANSWER 26 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML,  
 MR, NZ, SN, TD, TG  
 US 5977117 A 19991102 US 1996-590139 19960123  
 AU 9715324 A1 19970801 AU 1997-15324 19970103  
 EP 876364 A2 19981111 EP 1997-901420 19970103  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI  
 PRIORITY APPLN. INFO.: US 1996-583871 A 19960105  
 US 1996-590139 A 19960123  
 WO 1997-US366 W 19970103

OTHER SOURCE(S): MARPAT 127:161816  
 IT 193757-00-SP  
 RL: BAC (Biological activity or effector, except adverse); BSU  
 (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use);  
 BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (preparation of aryl- and/or heteroaryl-substituted benzoic acids as  
 endothelin antagonists and/or agonists)  
 RN 193757-00-5 CAPLUS  
 CN Benzoic acid, 2-(1,3-benzodioxol-5-yloxy)-6-(4-fluorophenoxy)- (9CI) (CA  
 INDEX NAME)




L52 ANSWER 27 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 EP 836384 A1 19980422 EP 1996-919422 19960613  
 R: DE, FR, GB, IT  
 CN 1188394 A 19980722 CN 1996-194937 19960613  
 BR 9609001 A 19990629 BR 1996-9001 19960613  
 JP 11508257 T2 19990721 JP 1996-503876 19960613  
 ZA 9605196 A 19971219 ZA 1996-5196 19960619  
 PRIORITY APPLN. INFO.: US 1995-341P P 19950620  
 WO 1996-US10326 W 19960613

OTHER SOURCE(S): MARPAT 126:153997  
 IT 186979-62-4P 186979-63-5P 186979-65-7P  
 186980-15-4P 186980-17-6P 186980-20-1P  
 186980-51-8P 186980-52-9P 186980-53-0P  
 186981-79-3P 186981-82-0P 186981-83-9P  
 RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological  
 study); PREP (Preparation); USES (Uses)  
 (preparation as arthropodicide and fungicide)  
 RN 186979-62-4 CAPLUS  
 CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)phenoxy]phenyl]-2,4-  
 dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186979-63-5 CAPLUS  
 CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)-2-nitrophenoxy]phenyl]-2,4-  
 dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186979-65-7 CAPLUS  
 CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2-fluorophenoxy)phenoxy]phenyl]-2,4-  
 dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)

L52 ANSWER 27 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI




AB Preparation and title uses are given for I (E = (un)substituted  
 1,2-phenylene,  
 naphthalene or heterocyclyl; A = O, S, N, NR3 or CR4; G = C or N; when G  
 is C, then A is O, S or NR3 and a the floating double bond is attached to  
 G; and when G is N, then A is N or CR4 and the floating double bond is  
 attached to A; W = O, S, NH, N(C1-C6 alkyl) or NO(C1-C6 alkyl); X = H,  
 OR1, SO2R1, halo, C1-C6 alkyl, C1-C6 haloalkyl, C3-C6 cycloalkyl; cyano,  
 NH2, NR1, N(C1-C6 alkyl)R1, NH(C1-C6 alkoxy) or N(C1-C6 alkoxy)R1; R2 =  
 H, C1-C6 alkyl, C1-C6 haloalkyl, C2-C6 haloalkyl, C2-C6 alkenyl, C2-C6  
 haloalkenyl, C2-C6 alkynyl, C2-C6 haloalkynyl, C3-C6 cycloalkyl, C2-C4  
 alkylcarbonyl, C2-C6 alkoxy, C2-C6 alkoxyalkenyl, hydroxy, C1-C2 alkoxy or acetyloxy;  
 R1 = (halo)alkyl, (halo)alkenyl, etc.; R3 = H, (halo)alkyl, etc.; Y = O,  
 CO,  
 SO, etc.; Z = (un)substituted alkyl, alkenyl or alkynyl, R4 = H, halo,  
 alkyl, etc.; m = 0, 1 or 2).


ACCESSION NUMBER: 1997:168566 CAPLUS  
 DOCUMENT NUMBER: 126:153997  
 TITLE: Preparation of arthropodicide and fungicide cyclic  
 amides  
 INVENTOR(S): Brown, Richard James; Chan, Dominic Ming-Tak; Howard,  
 Michael Henry, Jr.; Daniel, Dillon Jancey; Clark,  
 David  
 PATENT ASSIGNEE(S): Alan; Selby, Thomas Paul  
 E. I. Du Pont de Nemours & Co., USA; Brown, Richard  
 James; Chan, Dominic Ming-Tak; Howard, Michael Henry,  
 Jr.; Daniel, Dillon Jancey; Clark, David Alan; Selby,  
 Thomas Paul  
 SOURCE: PCT Int. Appl., 20 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9700612	A1	19970109	WO 1996-US10326	19960613
W:	AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KE, KS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9661770	A1	19970122	AU 1996-61770	19960613


L52 ANSWER 27 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 186980-15-4 CAPLUS  
 CN Benzonitrile, 2-[3-[2-(1,5-dihydro-3-methoxy-1-methyl-5-oxo-4H-1,2,4-  
 triazol-4-yl)phenoxy]phenoxy]-6-fluoro- (9CI) (CA INDEX NAME)




RN 186980-17-6 CAPLUS  
 CN Benzonitrile, 2-[3-[2-(1,5-dihydro-3-methoxy-1-methyl-5-oxo-4H-1,2,4-  
 triazol-4-yl)phenoxy]phenoxy]-3-fluoro- (9CI) (CA INDEX NAME)




RN 186980-20-1 CAPLUS  
 CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,5-difluorophenoxy)phenoxy]phenyl]-2,4-  
 dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)





RN 186980-51-8 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(3-fluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




RN 186980-52-9 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,3-difluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)




RN 186980-53-0 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,4-difluorophenoxy)phenoxy]phenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186981-79-3 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186981-82-8 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(4-fluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)



RN 186981-83-9 CAPIUS  
CN 3H-1,2,4-Triazol-3-one, 4-[2-[3-(2,6-difluorophenoxy)phenoxy]-6-methylphenyl]-2,4-dihydro-5-methoxy-2-methyl- (9CI) (CA INDEX NAME)

AB The introduction of tetrafluoro- and pentafluorophenoxy moieties into a variety of pentafluorobenzenes C6F5R (R = CF3, CN, NO2) is accomplished by


employing the trimethylsilyl ethers (siloxanes) 4-HC6F4OSiMe3 (1) and C6F5OSiMe3 (2) as transfer agents. Depending on the nature of the electrophile, the stoichiometry of the reaction, and the reaction conditions, polysubstituted polyfluorodiphenyl ethers are obtained. Excess C6F5R results in the formation of 1,4-monosubstituted benzenes (di-Ph ethers) 4-(4'-XC6F4O)C6F4R [R = CF3, X = H (3), F (4); R = CN, X = H (5), F (6); R = NO2, X = H, F]. When R = NO2, the 1,2-substituted isomers are also detected. Addnl. byproducts that are isolable are the disubstituted benzenes 2,4-(4'-XC6F4O)2C6F3R (R = CN, X = H; R = CN, X = F; R = NO2, X = H; R = NO2, X = F). Excess 1 or 2, when reacted with C6F5R, results in the formation of the trisubstituted benzenes 2,4,6-(4'-XC6F4O)3C6F2R [R = CN, X = H (13); R = CN, X = F (14); R = NO2, X = H (15); R = NO2, X = F (16)]. Hydrolysis of nitrile-containing di-Ph ethers (5, 6, 13, and 14) under acidic conditions results in the substituted benzoic acids 4-(4'-XC6F4O)C6F4COOH [X = H (17), F (18)] and 2,4,6-(4'-XC6F4O)3C6F2COOH (X = H, F). These acids are decarboxylated to form the resp. hydropolyfluoro aroms. (4-HC6F4)2O (23), 4-(C6F5O)C6F4H, and 2,4,6-(4'-XC6F4O)3C6F2H (X = H, F). In addition to acid 17, alkaline hydrolysis of 5 gives the  $\alpha$ -hydroxy-substituted acid 4-(4'-HC6F4O)C6F3(2-OH)COOH. Alkaline hydrolysis under milder conditions enables the isolation of the amide 4-(4'-HC6F4O)C6F4CONH2 (26). The compds. 3, 4, 14-18, 23, and 26 have been characterized by single-crystal x-ray diffraction anal. The presence of a hydrogen atom in 3, as well as protection of the reactive 4'-position with a trifluoromethyl group, gives

4-(4'-CF3C6F4O)C6F4Li (3a) on reaction with n-butyllithium. In situ reactions between 3a and ketones or acid chlorides result in novel mono- or bis(perfluorodiphenyl ether)-substituted tertiary alca. 4-(4'-CF3C6F4O)C6F4C(R)(R')OH (R/R' = CF3, C6F5, Ph, C3F7/C8F17, C6F5/CH3), [4-(4'-CF3C6F4O)C6F4]2C(R)OH (R = CF3, C3F7, C7F15, i-C3H7). When R = i-C3H7, the major product is the ester


[4-(4'-CF3C6F4O)C6F4]2C(i-C3H7)OC(O)(i-C3H7). The ketone C3F7(C8F17)CO is synthesized and characterized. Reaction of 3a with hexafluoroglutaric chloride gives [4-(4'-CF3C6F4O)C6F4]2C(OH)(CF2)3C(O)C6F4O(4''-C6F4CF3), whereas with di-Me carbonate or carbonyl fluoride, [4-(4'-CF3C6F4O)C6F4]2CO as well as small amts. of [4-(4'-CF3C6F4O)C6F4]3COH and [4-(4'-CF3C6F4O)C6F4]3COOC(O)C4H9 are formed. Residual n-butyllithium cleaves the intermediate 4-(4'-CF3C6F4O)C6F4COOCH3 to form 4'-CF3C6F4C4H9 and 4-HOC6F4COOCH3.

ACCESSION NUMBER: 1997:96834 CAPIUS  
DOCUMENT NUMBER: 126:89102  
TITLE: Studies on the Reactivity of Tetrafluoro- and Pentafluorophenyl Trimethylsilyl ether with Pentafluorobenzenes. Chemistry and X-ray Structural Investigations of Polyfluorodiphenyl ethers  
AUTHOR(S): Krumm, Burkhard; Vij, Ashwani; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 8344-2343, USA  
SOURCE: Inorganic Chemistry (1997), 36(3), 366-381  
CODEN: INOCAJ; ISSN: 0020-1669  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English

L52 ANSWER 28 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 IT 16031-36-0P 185697-15-8P 185697-20-5P  
 185697-22-7P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (chemical of polyfluorodiphenyl ethers)  
 RN 16031-36-0 CAPLUS  
 CN Benzonitrile, 3,5-difluoro-2,4,6-tris(pentafluorophenoxy)- (8CI, 9CI)  
 (CA INDEX NAME)




RN 185697-15-8 CAPLUS  
 CN Benzonitrile, 3,5-difluoro-2,4,6-tris(2,3,5,6-tetrafluorophenoxy)- (9CI)  
 (CA INDEX NAME)




RN 185697-20-5 CAPLUS  
 CN Benzoic acid, 3,5-difluoro-2,4,6-tris(2,3,5,6-tetrafluorophenoxy)- (9CI)  
 (CA INDEX NAME)

L52 ANSWER 28 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 185697-22-7 CAPLUS  
 CN Benzoic acid, 3,5-difluoro-2,4,6-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)




IT 15963-72-1P 185697-11-4P 185697-13-6P  
 185697-14-7P 185697-17-0P 185697-18-1P  
 185697-21-6P 185697-23-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (chemical of polyfluorodiphenyl ethers)  
 RN 15963-72-1 CAPLUS  
 CN Benzonitrile, 2,3,5-trifluoro-4,6-bis(pentafluorophenoxy)- (8CI, 9CI)  
 (CA INDEX NAME)


L52 ANSWER 28 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 185697-11-4 CAPLUS  
 CN Benzonitrile, 2,3,5-trifluoro-4,6-bis(2,3,5,6-tetrafluorophenoxy)- (9CI)  
 (CA INDEX NAME)



RN 185697-13-6 CAPLUS  
 CN Benzene, 1,2,4-trifluoro-6-nitro-3,5-bis(2,3,5,6-tetrafluorophenoxy)- (9CI) (CA INDEX NAME)




RN 185697-14-7 CAPLUS  
 CN Benzene, 1,2,4-trifluoro-6-nitro-3,5-bis(pentafluorophenoxy)- (9CI) (CA INDEX NAME)




L52 ANSWER 28 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

RN 185697-17-0 CAPLUS  
 CN Benzene, 1,3-difluoro-5-nitro-2,4,6-tris(2,3,5,6-tetrafluorophenoxy)- (9CI) (CA INDEX NAME)




RN 185697-18-1 CAPLUS  
 CN Benzene, 1,3-difluoro-5-nitro-2,4,6-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)




RN 185697-21-6 CAPLUS  
 CN Benzene, 2,4-difluoro-1,3,5-tris(2,3,5,6-tetrafluorophenoxy)- (9CI) (CA INDEX NAME)





RN 185697-23-8 CAPIUS  
CN Benzene, 2,4-difluoro-1,3,5-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L52 ANSWER 30 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The title compds. (I; ring A = imidazolyl, tetrazolyl, Q, Q1; wherein X = O, S, NH; Y = N, CH; R1 = H, halo, lower alkyl, lower hydroxyalkyl, lower alkoxy, CF3, NO2, CO2H, lower alkoxy carbonyl, CH2 NHCONHCO2R5, CH:NOH; wherein R5 = H, lower alkyl; R2, R3 = H, halo; R4 = H, HO; n = 0,1),

which lower blood sugar based on the enhancement of insulin sensitivity, have low toxicity, and are useful as antidiabetics for treating or preventing noninsulin-dependent diabetes and various diabetes complications (no data), are prepared Thus, 3-(4-trifluoromethylphenoxy)phenol 6, K2CO3 3.3, and 4-fluorobenzaldehyde 3.0 g were stirred in DMSO at 100° for 10 h to give 6 g 4-[3-(4-trifluoromethylphenoxy)phenoxy]benzaldehyde (II; R = CHO), which (6 g) was condensed with 1.8 g hydroxylamine hydrochloride in the presence of 2.0 g NH4OAc in aqueous MeOH at room temperature for 2 h and under reflux for 30 min to give the oxime II (R = CH:NOH) (4.0 g). The latter oxime (3.0 g) was dissolved in 30 mL EtOH and after adding 1.2 g pyridine-borane complex, treated dropwise with 12 mL 4 N aqueous HCl, and left to stand at room temperature for 4 h to give 2.5 g II (R = CH2NHOH), which (1.5 g) was dissolved in THF, treated with 0.7 g ethoxycarbonyl isocyanate, left to stand for 30 min, made alkaline with 1 N aqueous NaOH, left to stand at room temperature for 2 h, and made acidic with 6 N aqueous HCl to give 1.0 g the 1,2,4-oxadiazolidine-3,5-dione derivative II (R = Q2).

ACCESSION NUMBER: 1996:326164 CAPIUS  
DOCUMENT NUMBER: 125:10826  
TITLE: Preparation of p-[(phenoxy or benzyloxy)phenoxy]benzylazole derivatives for

lowering

INVENTOR(S): Niigata, Kunihiro; Takahashi, Takumi; Maruyama, Tatsuya; Suzuki, Takayuki; Onda, Kenichi; Konya, Tooru; Noshiro, Osamu  
PATENT ASSIGNEE(S): Yamanouchi Pharma Co Ltd, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 26 pp.  
CODEN: JKOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08059638	A2	19960305	JP 1994-202503	19940826

PRIORITY APPLN. INFO.: JP 1994-202503 19940826

OTHER SOURCE(S): MRRPAT 125:10826

IT 177031-89-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of p-[(phenoxy or benzyloxy)phenoxy]benzylazole derivs.)

for


Page 343

L52 ANSWER 29 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN


AB A general method for the synthesis of 1,3-bis(aryloxy)-5-nitrobenzenes by the condensation of phenols with 1,3,5-trinitrobenzene or aryl 3,5-dinitrophenyl ethers in the presence of solid K2CO3 in amide-type dipolar aprotic solvents has been developed.

ACCESSION NUMBER: 1996:477446 CAPIUS  
DOCUMENT NUMBER: 125:221267  
TITLE: A general method for the synthesis of 1,3-bis(aryloxy)-5-nitrobenzenes  
AUTHOR(S): Shevelev, Svyatoslva; Dutov, Mikhail D.; Vatsadze, Irina A.; Korolev, Maksim A.; Rusanov, Aleksandr L.; N. D. Zelinsky Inst. Organic Chem., Russian Academy of Sci., Moscow, 117913, Russia  
SOURCE: Mendeleev Communications (1996), (4), 155-157  
CODEN: MENCEX; ISSN: 0959-9436  
PUBLISHER: Russian Academy of Sciences  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 168839-60-9P 181507-42-6P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 168839-60-9 CAPIUS  
CN Benzene, 1-(4-fluorophenoxy)-3-nitro-5-phenoxy- (9CI) (CA INDEX NAME)



RN 181507-42-6 CAPIUS  
CN Acetamide, N-[4-[3-(4-fluorophenoxy)-5-nitrophenoxy]phenyl]- (9CI) (CA INDEX NAME)




L52 ANSWER 30 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

lowering blood sugar as antidiabetics)

RN 177031-89-9 CAPIUS

CN Benzaldehyde, 4-[3-[4-[(3,5-dioxo-1,2,4-oxadiazolidin-2-yl)methyl]-2-fluorophenoxy]phenoxy]-3-fluoro-, 1-oxime (9CI) (CA INDEX NAME)




IT 163301-37-9  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of p-[(phenoxy or benzyloxy)phenoxy]benzylazole deriva.

for

lowering blood sugar as antidiabetics)

RN 163301-37-9 CAPIUS

CN Benzaldehyde, 4,4'-[1,3-phenylenebis(oxy)]bis[3-fluoro- (9CI) (CA INDEX NAME)




IT 177032-00-7P 177032-01-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of p-[(phenoxy or benzyloxy)phenoxy]benzylazole derivs.

for


lowering blood sugar as antidiabetics)

RN 177032-00-7 CAPIUS

CN Benzaldehyde, 4,4'-[1,3-phenylenebis(oxy)]bis[3-fluoro-, dioxime (9CI) (CA INDEX NAME)



RN 177032-01-8 CAPIUS  
 CN Benzaldehyde,  
 3-fluoro-4-[3-[2-fluoro-4-[(hydroxyamino)methyl]phenoxy]phenoxy]-, oxime (9CI) (CA INDEX NAME)




AB Alkyl aryl sulfones, e.g., 2-(4-methylsulfonylphenoxy)-6-(3-trifluoromethylphenoxy)pyridine, are disclosed which are useful as high temperature and magnetic recording media lubricants. Also disclosed are lubricating mixts. containing the alkyl aryl sulfones, magnetic recording media containing the alkyl aryl sulfones, and a process for lubricating

the magnetic recording media with the alkyl aryl sulfones.  
 ACCESSION NUMBER: 1996:71502 CAPIUS  
 DOCUMENT NUMBER: 124:150632  
 TITLE: Alkyl aryl sulfones and their use as lubricants in high temperature and magnetic recording media applications  
 INVENTOR(S): Pawloski, Chester E.; Nader, Bassam S.  
 PATENT ASSIGNEE(S): The Dow Chemical Company, USA  
 SOURCE: U.S., 12 pp.  
 CODEN: USXKAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5480568	A	19960102	US 1994-279036	19940722

PRIORITY APPLN. INFO.:  
 US 1994-279036 19940722

OTHER SOURCE(S): MARPAT 124:150632  
 IT 173447-54-6P  
 RL: IMF (Industrial manufacture); NUU (Other use, unclassified); PHU (Preparation, unclassified); PREP (Preparation); USES (Uses)  
 (preparation of alkyl aryl sulfones and their use as lubricants in high temperature and magnetic recording media applications)  
 RN 173447-54-6 CAPIUS  
 CN Benzene, 1-(3-fluorophenoxy)-3-[4-(methylsulfonyl)phenoxy]- (9CI) (CA INDEX NAME)



GI




AB Title compds. [I; A = (un)substituted Ph, 2-pyridyl, 1-alkylimidazolyl, etc.; R = NO2, halo, cyano, (halo)alkylsulfonyl; R1 = H, halo, NO2; R2 = H, halo, (halo)alkyl; R6 = (CR3R4)nR5; R3,R4 = H, (halo)alkyl; R3R4 = (CH2)2-5; R5 = cyano, CO2H, alkoxy, carbonyl, etc.; W, W1 = O, S, (alkyl)imino; m = 0 or 1; n = 0-5] were prepared. Thus, Ph ether II (R7 = NO2) was condensed with 4-(HO)C6H4OCHMeCO2Me to give II [R7 = OC6H4(OCHMeCO2Me)-4]. II [R7 = OC6H4(CO2Me)-3] gave 45-100% control of 4 weeds (e.g., 100% control of Amaranthus retroflexus) at 0.125kg/ha postemergent with slight to trace effect on 4 crops, e.g., winter wheat.

ACCESSION NUMBER: 1995:701871 CAPIUS  
 DOCUMENT NUMBER: 123:111672  
 TITLE: Preparation of [(aryloxy)phenoxy]phenoxy]alkanoates and analogs as herbicides  
 INVENTOR(S): Condon, Michael E.; Crews, Alvin D., Jr.  
 PATENT ASSIGNEE(S): American Cyanamid Co., USA  
 SOURCE: Eur. Pat. Appl., 68 pp.  
 CODEN: EPXKDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 647612	A1	19950412	EP 1994-115265	19940928
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
US 5424280	A	19950613	US 1993-133698	19931007
CA 2133690	A	19950408	CA 1994-2133690	19941005
JP 07165659	A2	19950627	JP 1994-266156	19941005
NO 9403765	A	19950410	NO 1994-3765	19941006
AU 9474470	A1	19950427	AU 1994-74470	19941006
BR 9404010	A	19950613	BR 1994-4010	19941006
HU 69022	A2	19950828	HU 1994-2891	19941006
ZA 9407834	A	19960206	ZA 1994-7834	19941006
CN 1104207	A	19950628	CN 1994-116340	19941007

PRIORITY APPLN. INFO.:  
 US 1993-133698 A 19931007


OTHER SOURCE(S): MARPAT 123:111672  
 IT 165379-41-9P 165379-42-0P 165379-43-1P  
 165379-44-2P 165379-46-4P 165379-59-9P  
 165379-60-2P 165379-62-4P 165379-63-5P  
 165379-64-6P 165379-65-7P 165379-78-2P  
 165379-83-9P 165379-86-2P 165379-82-0P  
 165379-93-1P 165379-94-2P 165380-05-2P  
 RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (preparation of [(aryloxy)phenoxy]phenoxy]alkanoates and analogs as herbicides)  
 RN 165379-41-9 CAPIUS  
 CN Propanoic acid, 2-[4-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-42-0 CAPIUS  
 CN Benzoic acid, 2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-43-1 CAPIUS  
 CN Benzoic acid, 4-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-44-2 CAPIUS  
 CN Benzoic acid, 3-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-46-4 CAPLUS  
CN Benzenepropanoic acid, 2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-59-9 CAPLUS  
CN Acetic acid, [4-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-60-2 CAPLUS  
CN Propanoic acid, 2-[3-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-62-4 CAPLUS  
CN Propanoic acid, 2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-83-9 CAPLUS  
CN Acetic acid, [2-[5-(4-fluorophenoxy)-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-86-2 CAPLUS  
CN Acetic acid, [2-[5-[2-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-92-0 CAPLUS  
CN Acetic acid, [2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-93-1 CAPLUS  
CN Benzoic acid, 4-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-63-5 CAPLUS  
CN Acetic acid, [3-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-64-6 CAPLUS  
CN Acetic acid, [2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-65-7 CAPLUS  
CN Butanoic acid, 4-[2-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-78-2 CAPLUS  
CN Acetic acid, [2-[5-(2-chloro-4-fluorophenoxy)-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)



RN 165379-94-2 CAPLUS  
CN Benzoic acid, 3-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165380-05-2 CAPLUS  
CN Acetic acid, [2-[5-[2-chloro-5-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]phenoxy]-, methyl ester (9CI) (CA INDEX NAME)




IT 165379-47-5P 165379-53-3P 165379-54-4P  
165379-55-5P 165379-56-6P 165379-57-7P  
165379-58-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or Reagent)  
(Preparation of [(aryloxy)phenoxy]phenoxy)alkanoates and analogs as herbicides)


RN 165379-47-5 CAPLUS  
CN Benzene, 4-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-(4-methoxyphenoxy)-1-nitro- (9CI) (CA INDEX NAME)




RN 165379-53-3 CAPLUS  
CN Phenol, 4-[5-[2-chloro-6-fluoro-4-(trifluoromethyl)phenoxy]-2-nitrophenoxy]-, methyl ester (9CI) (CA INDEX NAME)




RN 165379-54-4 CAPLUS  
CN Phenol, 4-[5-(4-chloro-2-fluorophenoxy)-2-nitrophenoxy]- (9CI) (CA INDEX NAME)



RN 165379-55-5 CAPLUS  
CN Phenol, 3-[5-(4-chloro-2-fluorophenoxy)-2-nitrophenoxy]- (9CI) (CA INDEX NAME)



RN 165379-56-6 CAPLUS  
CN Phenol, 2-[5-(4-chloro-2-fluorophenoxy)-2-nitrophenoxy]- (9CI) (CA INDEX NAME)



RN 165379-57-7 CAPLUS  
CN Phenol, 2-[3-(4-chloro-2-fluorophenoxy)-2-fluoro-6-nitrophenoxy]- (9CI) (CA INDEX NAME)

AB Facile synthesis of  $Cp^*Ru^+$  ( $Cp^*$  = pentamethylcyclopentadienyl)  $\pi$ -complexes of highly electron deficient aroms., including 1,3,5-trichloro-, 1,2,4,5-tetrachloro-, pentachloro-, and hexachlorobenzene, is accomplished by the ligand exchange reaction with  $(Cp^*Ru(CH_3CN)_3)SO_3CF_3^-$  in polar solvents under mild reaction conditions. The extraordinary activating ability of the  $Cp^*Ru^+$  moiety is demonstrated by rapid and quant. nucleophilic substitution reactions of the 1,3,5-tri- and 1,2,4,5-tetrachlorobenzene  $\pi$ -complexes with potassium phenoxide, thiophenoxide, 4-aminophenoxide, 4-chlorophenoxide, and 4-fluorophenoxide.

This methodol. allows syntheses of highly substituted and functionalized aromatic compds.

ACCESSION NUMBER: 1995:701727 CAPLUS

DOCUMENT NUMBER: 123:169866

TITLE: Synthesis and Nucleophilic Substitution of Highly

Chlorinated Arene ( $\eta^5$ -Pentamethylcyclopentadienyl)ruthenium  $\pi$ -Complexes

Dembek, Alexa A.; Fagan, Paul J.

Experimental Station, DuPont Central Research and

Development, Wilmington, DE, 19880-0328, USA

Organometallics (1995), 14(8), 3741-5

CODEN: ORGN7; ISSN: 0276-7333

American Chemical Society

PUBLISHER: Journal

DOCUMENT TYPE: English

LANGUAGE: CASREACT 123:169866

OTHER SOURCE(S):

IT 163709-89-5P 167227-89-6P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

RN 163709-89-5 CAPLUS


CN Ruthenium(1+), [(1,2,3,4,5- $\eta$ )-1,2,3,4,5-pentamethyl-2,4-cyclopentadien-1-yl][(1,2,3,4,5,6- $\eta$ )-1,3,5-tris(4-fluorophenoxy)benzene]-, salt with trifluoromethanesulfonic acid (1:1) (9CI) (CA INDEX NAME)

CM 1


CRN 163709-88-4

CMF C34 H30 F3 O3 Ru

CCI CCS




RN 165379-58-8 CAPLUS  
CN Phenol, 2-[2-chloro-3-(4-chloro-2-fluorophenoxy)-6-nitrophenoxy]- (9CI) (CA INDEX NAME)




PAGE 1-A



PAGE 2-A



PAGE 3-A



CM 2

CRN 37181-39-8  
CMF C F3 O3 S

RN 167227-89-6 CAPIUS  
CN Ruthenium(II+), [(1,2,3,4,5-η)-1,2,3,4,5-pentamethyl-2,4-cyclopentadien-1-yl][(1,2,3,4,5,6-η)-1,2,4,5-tetrakis(4-fluorophenoxy)benzene]-, salt with trifluoromethanesulfonic acid (1:1) (9CI) (CA INDEX NAME)


CM 1

CRN 167227-88-5  
CMF C40 H33 F4 O4 Ru  
CCI CCS


PAGE 1-A



PAGE 2-A



PAGE 3-A



CM 2

CRN 37181-39-8  
CMF C F3 O3 S

L52 ANSWER 34 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN  
AB A method of preparing 5-nitroresorcinol diaryl ethers bearing both

different and equal substituents was elaborated. The method is based on the nucleophilic substitution of a nitro group in 1,3,5-(O2N)3C6H3 or in 3,5-dinitrophenyl aryl ethers by phenols in the presence of a base. Similarly, tris-Ph ether of phloroglucinol was obtained from 1-nitro-3,5-diphenoxybenzene.

ACCESSION NUMBER: 1995:656708 CAPIUS  
DOCUMENT NUMBER: 123:256255

TITLE: Phenol substitution of nitro groups in 1,3,5-trinitrobenzene - method of preparation of 5-nitroresorcinol diaryl ethers and 3,5-dinitrophenyl aryl ethers

AUTHOR(S): Shevelev, S. A.; Dutov, M. D.; Vatsadze, I. A.; Serushkina, O. V.; Korelev, M. A.; Rusanov, A. L.  
CORPORATE SOURCE: N. D. Zelinsky Inst. Org. Chem., Moscow, 117913, Russia

SOURCE: Izvestiya Akademii Nauk, Seriya Khimicheskaya (1995), (2), 393-4  
CODEN: IASKEA

PUBLISHER: Institut Organicheskoi Khimii im. N. D. Zelinskogo  
Rossiiskoi Akademii Nauk


DOCUMENT TYPE: Journal  
LANGUAGE: Russian  
OTHER SOURCE(S): CASREACT 123:256255  
IT 168839-60-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(phenol substitution of nitro groups in trinitrobenzene in preparation of

nitroresorcinol diaryl ethers and dinitrophenyl aryl ethers)

RN 168839-60-9 CAPIUS


CN Benzene, 1-(4-fluorophenoxy)-3-nitro-5-phenoxy- (9CI) (CA INDEX NAME)






OTHER SOURCE(S): MARPAT 122:316019  
 IT 163464-84-4  
 RL: MDA (Modifier or additive use); POF (Polymer in formulation); FRP  
 (Properties); TEM (Technical or engineered material use); USES (Uses)  
 (near-IR absorber; heat-radiation-shielding compns. containing  
 polymers  
 and)  
 RN 163464-84-4 CAPIUS  
 CN Zinc, [N,N',N'',N''']-tetrahexyl-1,3,4,8,10,11,15,17,18,22,24,25-  
 dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-  
 tetraminato(2-)-<N29,<N30,<N31,<N32>-; (SP-4-1)-  
 (9CI) (CA INDEX NAME)


PAGE 1-A





PAGE 2-A



PAGE 3-A



PAGE 4-A



AB Novel ruthenium pi-arene complexes, I-III' (R1, R2 = independently H, halo, R3; R3 = inert monovalent radical having Hammett constant > -0.35 and < -0.12; R4 = cyclopentadienyl, pentamethylcyclopentadienyl; X = perfluoroalkylsulfonate, hexafluorophosphate), in which the arene ring has 3 or more halogen atoms bound to it are disclosed. These compds. are reacted with phenoxides and thiophenoxides to form ruthenium pi-arene complexes of polyfunctional aromatic ethers and thioethers which are useful as monomers and crosslinking agents. Thus, reaction of 1,3,5-trichlorobenzene with [Cp\*Ru(MeCN)] [SO3CF3] [Cp\* = pentamethylcyclopentadienyl] in THF gave [Cp\*Ru(1,3,5-trichlorobenzene)] [SO3CF3], which on treatment with potassium p-aminophenoxide in MeCN gave [Cp\*Ru(1,3,5-tris(4-aminophenoxy)benzene)] [SO3CF3].

ACCESSION NUMBER: 1995:403412 CAPIUS  
 DOCUMENT NUMBER: 123:9696  
 TITLE: Preparation and reactions of polyhaloaromatic ruthenium complexes  
 INVENTOR(S): Dembek, Alexa A.  
 PATENT ASSIGNEE(S): E. I. Du Pont de Nemours & Co., USA  
 SOURCE: U.S., 6 pp.  
 CODEN: USXQUM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE

OTHER SOURCE(S): MARPAT 123:9696  
 IT 163709-89-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 163709-89-5 CAPIUS  
 CN Ruthenium(1+), [(1,2,3,4,5-η)-1,2,3,4,5-pentamethyl-2,4-cyclopentadien-1-yl][(1,2,3,4,5,6-η)-1,3,5-tris(4-fluorophenoxy)benzene]-, salt with trifluoromethanesulfonic acid (1:1) (9CI) (CA INDEX NAME)


CH 1

CRN 163709-88-4  
 CMF C34 H30 F3 O3 Ru  
 CCI CCS


PAGE 1-A



PAGE 2-A



PAGE 3-A



CH 2

CRN 37181-39-8  
 CMF C F3 O3 S



L52 ANSWER 38 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB Hexafluorobenzene reacts readily with a variety of trimethylsilyl ethers  
 ROSiMe3 (R = CF3CH2, FCH2CH2, H(CF2)nCH2 (n = 2, 4), CF3(CF2)6CH2,  
 CF3(CF2)5CH2CH2, Me3SiOCH2CH2, C6F5OCH2CH2, C6H5, 4-FC6H4) to give from  
 mono- to hexapolyfluoroalkoxy- and polyfluoroaryloxy-benzenes. The  
 structure of C6(OCH2CF3)6 has been confirmed by single-crystal X-ray  
 anal.

The perfluorinated ether C6F5OCF2CF3 may be synthesized from C6F5OCH2CF3  
 by chlorination and subsequent fluorination with SbF3/SbCl5. The  
 chlorination of 5,6,7,8-tetrafluoro-1,4-benzodioxane is also discussed.


ACCESSION NUMBER: 1995:57102 CAPIUS  
 DOCUMENT NUMBER: 122:31037  
 TITLE: Reaction of hexafluorobenzene with trimethylsilyl  
 ethers  
 AUTHOR(S): Zhang, Y. F.; Kirchmeier, Robert L.; Shreeve, Jean'ne  
 M.  
 CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow,  
 ID, 83844-2343, USA

SOURCE: Journal of Fluorine Chemistry (1994), 68(3), 287-92  
 CODEN: JFLCAR; ISSN: 0022-1139

DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 122:31037


IT 159752-90-6 159752-93-9 159752-94-0  
 RL: PREP (Properties)  
 (reaction of hexafluorobenzene with trimethylsilyl ethers)

RN 159752-90-6 CAPIUS  
 CN Benzene, hexakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)




RN 159752-93-9 CAPIUS  
 CN Benzene, fluoropentakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)







RN 159752-94-0 CAPLUS  
CN Benzene, 1,4-difluoro-2,3,5,6-tetrakis(4-fluorophenoxy)- (9CI) (CA INDEX NAME)



L52 ANSWER 39 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Six F-containing poly(arylene ethers) were prepared by polymerizing decafluorobiphenyl with 4,4'-(hexafluoroisopropylidene)diphenol (bisphenol AP), 9,9-bis(4-hydroxyphenyl)fluorene, 1,1-bis(4-hydroxyphenyl)-1-phenylethane (bisphenol AP), phenolphthalein, fluorescein, and Me 3,5-dihydroxybenzoate. The polymers exhibited low dielec. consts. and moisture absorption and excellent thermal and mech. properties and may be useful in electronic applications.

ACCESSION NUMBER: 1994:484111 CAPLUS  
DOCUMENT NUMBER: 121:84111  
TITLE: Synthesis and characterization of new poly(arylene ethers) with low dielectric constant  
AUTHOR(S): Mercer, Frank W.; Coffin, Chris; Duff, David W.  
CORPORATE SOURCE: Raychem Corp., Menlo Park, CA, 94025-1164, USA  
SOURCE: ACS Symposium Series (1994), 537(Polymer for Microelectronics), 546-53  
CODEN: ACSMC8; ISSN: 0097-6156  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 156589-15-0P, Decafluorobiphenyl-methyl 3,5-dihydroxybenzoate copolymer, SRU  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (preparation and properties of)  
RN 156589-15-0 CAPLUS  
CN Poly[oxy[5-(methoxycarbonyl)-1,3-phenylene]oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)




AB Title compds. I [A = C1-12 alkylene, C2-12 alkenylene, C3-8 cycloalkylene;  
R1 = H, C1-6 alkylthio, (substituted) PhS, (substituted) phenyl-C1-6 alkylthio, halo, 2-, 3-, 4-pyridylthio, 2-, 3-thienylthio, 2-thiazolylthio, etc.; R2 = R5CONH where R5 = C1-6 alkyl, substituted ureido, etc., R5NR6CO, wherein R6 = H, C1-6 alkyl, etc.; R3 = phenyl-C1-6-alkyl, hetero-C1-6-alkyl, etc.; R4 = C1-6 alkyl, C1-12 alkoxy, (substituted) Ph, (substituted) PhO, (substituted) phenylalkyloxy, (substituted) naphthylloxy, (substituted) heteroarylalkyloxy] or a salt, ester, or amide, are prepared To I (A = Me2CCH2, R1 = Me3CS, R2 = HO2C, R3 = 4-ClC6H4CH2, R4 = 5-Me2CH) in C6H6 were added Et3N and N3PO(OPh)2, the mixture was refluxed for 1 h, and HONRMe.HCl in Et3N was added to give after workup I (A = Me2CCH2, R1 = Me3CS, R2 = MeN(OH)CONH, R3 = 4-ClC6H4CH2, R4 = 5-Me2CH) (II). II inhibited LTB4 biosynthesis in vitro in human whole blood and human polymorphonuclear leukocytes with IC50 of 1.3 and 0.15 μM, resp.


ACCESSION NUMBER: 1992:407801 CAPLUS  
DOCUMENT NUMBER: 117:7801  
TITLE: Preparation of indole derivatives which inhibit leukotriene biosynthesis  
INVENTOR(S): Brooks, Dee W.; Carter, George W.; Dellaria, Joseph F.; Maki, Robert G.; Rodrigues, Karen E.  
PATENT ASSIGNEE(S): Abbott Laboratories, USA  
SOURCE: PCT Int. Appl., 101 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9203132	A1	19920305	WO 1991-US5621	19910807
W: CA, JP, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE				
US 5095031	A	19920310	US 1990-570248	19900820
CA 2090006	AA	19920221	CA 1991-2090006	19910807
EP 544819	A1	19930609	EP 1991-917405	19910807
EP 544819	B1	19951206		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 06500557	T2	19940120	JP 1991-516031	19910807
AT 131051	E	19951215	AT 1991-917405	19910807
ES 2083595	T3	19960416	ES 1991-917405	19910807
US 5459150	A	19951017	US 1993-969294	19930216

L52 ANSWER 40 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
PRIORITY APPLN. INFO.: US 1990-570248 A2 19900820  
WO 1991-US5621 W 19910807

OTHER SOURCE(S): MARPAT 117:7801  
IT 141863-03-8P  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of, as leukotriene biosynthesis inhibitor)  
RN 141863-03-8 CAPLUS  
CN Acetic acid, [[3-[1-[(4-chlorophenyl)methyl]-3-[(1,1-dimethylethyl)thio]-5-[3-(4-fluorophenoxy)phenoxy]-1H-indol-2-yl]-2,2-dimethylpropylidene]amino]oxy]- (9CI) (CA INDEX NAME)






AB Reaction of perfluorinated cyclohexadienone I with C6F5ONa in MeCN or with PhOH or 3,5-(C6F5O)2C6F3OH in MeCN containing K2CO3 at 20° gave 71-86% products of substitution of I in the 3- and 5-positions of the diene ring, which isomerize upon heating. Reaction of I with phenols and K2CO3 in MeCN at 70° gave products in which 4 F atoms at positions 3 and 5 in the diene ring and 3 and 5 in the aromatic ring were substituted.

Reduction with Na2S2O4 of the tetrakis(aryloxy)dienones thus obtained gave 63-94% fluorinated 3,5-bis(aryloxy)phenols, which reacted with C6F6 to give 1,3,5-trisubstituted fluorinated polyphenyl ethers.


ACCESSION NUMBER: 1992:235168 CAPIUS  
DOCUMENT NUMBER: 116:235168  
TITLE: Nucleophilic substitution of a fluorine atom in perfluoro(phenoxycyclohexadienones). Synthesis of 1,3,5-trisubstituted fluorine-containing polyphenyl ethers  
AUTHOR(S): Kovtonyuk, V. N.; Kobrina, L. S.  
CORPORATE SOURCE: Novosib. Inst. Org. Khim., Novosibirsk, USSR  
SOURCE: Zhurnal Organicheskoi Khimii (1991), 27(11), 2289-97  
CODEN: ZORXAE; ISSN: 0514-7492  
DOCUMENT TYPE: Journal  
LANGUAGE: Russian  
OTHER SOURCE(S): CASREACT 116:235168

IT 141215-99-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and etherification of, with hexafluorobenzene)  
RN 141215-99-8 CAPIUS  
CN Phenol, 2,4,6-trifluoro-3,5-bis[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)




IT 141215-97-6P 141228-91-3P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and reduction of)  
RN 141215-97-6 CAPIUS  
CN 2,5-Cyclohexadien-1-one, 2,4,6-trifluoro-3,5-bis[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]-4-[2,4,6-trifluoro-3,5-bis[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]phenoxy]- (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 2-A




RN 141228-91-3 CAPIUS  
CN 2,5-Cyclohexadien-1-one, 2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)-4-[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



IT 141215-95-4P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and substitution reaction of, with perfluoro(phenoxycyclohexadienone))  
RN 141215-95-4 CAPIUS  
CN Phenol, 2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)- (9CI) (CA INDEX NAME)




IT 141216-00-4P 141216-01-5P 141228-90-2P  
141228-92-4P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 141216-00-4 CAPIUS  
CN Benzene, 1,3,5-trifluoro-2,4,6-tris(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



RN 141216-01-5 CAPIUS  
CN Benzene, 1,3,5-trifluoro-2-(pentafluorophenoxy)-4,6-bis[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 2-A

F

RN 141228-90-2 CAPIUS  
CN 2,5-Cyclohexadien-1-one, 2,3,4,5,6-pentafluoro-4-[2,4,6-trifluoro-3,5-bis(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



RN 141228-92-4 CAPIUS  
CN Benzene, pentafluoro(2,4,6-trifluoro-3,5-diphenoxypheoxy)- (9CI) (CA INDEX NAME)



AB Paddy fungicides, herbicides or plant-growth regulators are formulated with solid carrier and an oil. The oil (sp. gr.  $\leq 1$ ; b.p.  $>150^\circ$ ) is water-immiscible and miscible with the active ingredient. Granules comprised Me (E)-2-[2-{3-(pyrimidin-2-yloxy)phenoxy}phenyl]-3-methoxycarboxylate 1.0, surfactant B246 0.05, soybean oil 7.5, and Na sesquicarbonate to 100% weight/weight. In pot expts., the granules (3 Kg active ingredient/ha) prevented artificial Pyricularia oryzae infection of rice.


ACCESSION NUMBER: 1991:487516 CAPIUS  
DOCUMENT NUMBER: 115:87516  
TITLE: Agrochemical compositions for paddy  
INVENTOR(S): Warrington, Roger Paul; Ramsay, Guy; Bird, Neal  
PATENT ASSIGNEE(S): Imperial Chemical Industries PLC, UK  
SOURCE: Eur. Pat. Appl., 14 pp.  
CODEN: EPXXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 415569	A2	19910306	EP 1990-308478	19900801
EP 415569	A3	19921028		
EP 415569	B1	19980701		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
AT 167778	E	19980715	AT 1990-308478	19900801
ES 2117626	T3	19980816	ES 1990-308478	19900801
AU 9060220	A1	19910307	AU 1990-60220	19900806
AU 642151	B2	19931014		
IL 95305	A1	19941128	IL 1990-95305	19900806
HU 57536	A2	19911230	HU 1990-4960	19900810
BR 9003989	A	19910903	BR 1990-3989	19900813
CA 2023547	AA	19910302	CA 1990-2023547	19900817
JP 03193705	A2	19910823	JP 1990-228511	19900831
JP 2880778	B2	19990412		
CN 1049771	A	19910313	CN 1990-107234	19900901
CN 1044070	B	19990714		
US 5466693	A	19951114	US 1993-125702	19930923
PRIORITY APPLN. INFO.:				GB 1989-19833 A 19890901
				GB 1990-7995 A 19900409
				GB 1990-16783 19900731
				US 1990-570588 B1 19900821

OTHER SOURCE(S): MARPAT 115:87516

IT 122281-57-6  
RL: PROC (Process)  
(fungicidal formulation of, solid, for rice paddy)  
RN 122281-57-6 CAPIUS  
CN Benzenecetic acid, 2-[3-(2-fluorophenoxy)phenoxy]- $\alpha$ -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.






AB 3,4-Dinitrochlorobenzene reacted with 2,4-dichlorophenol and K<sub>2</sub>CO<sub>3</sub> in Me<sub>2</sub>NCHO to give 2,4-bis(2,4-dichlorophenoxy)nitrobenzene (I, R = 2,4-dichloro). 3-Nitrophenol reacted with 3,4-dichlorobenzene to yield 2-chloro-4-(trifluoromethyl)phenyl 3-nitrophenyl ether, which was nitrated to give 2-chloro-4-(trifluoromethyl)phenyl 3,4-dinitrophenyl ether (III). I (R = 2,4-dichloro) reacted with KOH to give 2,4-dichlorophenyl 3-hydroxy-4-nitrophenyl ether (III, R = 2,4-dichloro). Similarly

prepared were I (R = H, 2,4-dichloro-6-fluoro, 2- or 3-chloro, 2-chloro-4-nitro, 2-methyl-4-chloro, 2-methyl-4,6-dichloro, 2,4,6-trichloro, or 2-chloro-4-trifluoromethyl) and III (R = H, 2- or 3-chloro, 3-trifluoromethyl, 4-methylthio, 2-chloro-3-trifluoromethyl, 2-chloro-4-nitro, 4-chloro-2-Me, 2,4-dichloro-6-fluoro, 2,4-dichloro-6-Me, or 2,4,6-trichloro).

ACCESSION NUMBER: 1991:448952 CAPIUS  
DOCUMENT NUMBER: 115:48952  
TITLE: Synthesis of intermediates related to diphenyl ether herbicides  
AUTHOR(S): Yoshimoto, Takeo; Igarashi, Keiichi; Fujita, Takashi; Harayama, Takeo  
CORPORATE SOURCE: Life Sci. Bus. Dev. Dep., Mitsui-Toatsu Chem. Inc., Tokyo, 100, Japan  
SOURCE: Nippon Noyaku Gakkaishi (1990), 15(3), 341-52  
CODEN: NNGADV; ISSN: 0385-1559  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 115:48952  
IT 60793-43-3P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 60793-43-3 CAPIUS  
CN Benzene, 2,4-bis(2,4-dichloro-6-fluorophenoxy)-1-nitro- (9CI) (CA INDEX NAME)



L52 ANSWER 44 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA issue.

AB The title compds. [I; A, B, E = H, halo, OH, alkyl, alkoxy, haloalkyl, etc.; X = O, S; X = O, S, SO, SO<sub>2</sub>, CO, (un)substituted NH, etc.; Z = (un)substituted aryl, heterocyclyl] are prepared as fungicides. A mixture of

Me (E)-2-[2-(3-hydroxyphenoxy)phenyl]-3-methoxypropenoate (preparation given).

PhCH<sub>2</sub>Br, K<sub>2</sub>CO<sub>3</sub> and dry DMF was stirred 3 h to give Me (E)-2-[2-(3-benzyloxyphenoxy)phenyl]-3-methoxypropenoate (II). At 0.05% I totally controlled Puccinia recondita on wheat, Venturia inaequalis on apple, Pyricularia oryzae on rice, and other fungi.

ACCESSION NUMBER: 1989:514869 CAPIUS  
DOCUMENT NUMBER: 111:114869  
TITLE: Preparation of phenylpropenoates as fungicides  
INVENTOR(S): Anthony, Vivienne Margaret; Heaney, Stephen Paul; Beaumont, Kevin; Clough, John Martin; Crowley, Patrick Jelf; Godfrey, Christopher Richard Ayles; De Fraine, Paul John; Buckley, Alan John; Hutchings, Michael Gordon; Ferguson, Ian  
PATENT ASSIGNEE(S): Imperial Chemical Industries PLC, UK  
SOURCE: Eur. Pat. Appl., 119 pp.  
CODEN: EPOXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 307103	A2	19890315	EP 1988-307746	19880822
EP 307103	A3	19910417		
EP 307103	B1	19940921		
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
GB 2210041	A1	19890601	GB 1988-19878	19880822
GB 2210041	B2	19920701		
ES 2059527	T3	19941116	ES 1988-307746	19880822
IL 87573	A1	19930114	IL 1988-87573	19880826
AU 8821665	A1	19890316	AU 1988-21665	19880830
AU 611934	B2	19910627		
ZA 8806490	A	19890426	ZA 1988-6490	19880831
HU 47527	A2	19890328	HU 1988-4607	19880907
HU 202820	B	19910429		
DD 287401	A5	19910228	DD 1988-319606	19880907
PL 158290	B1	19920831	PL 1988-274572	19880908
RU 2014320	C1	19940615	RU 1988-4356509	19880908
DK 8805040	A	19890310	DK 1988-5040	19880909
BR 8804672	A	19890418	BR 1988-4672	19880909
CN 1032782	A	19890510	CN 1988-107270	19880909
CN 1021009	B	19930602		
JP 01121243	A2	19890512	JP 1988-224814	19880909
JP 2858756	B2	1990217		
CA 1309401	A1	19921027	CA 1988-577060	19880909
US 5334748	A	19940802	US 1988-242760	19880909
KR 127749	B1	19880402	KR 1988-11630	19880909
RU 2026282	C1	19950109	RU 1989-4742419	19891121
RU 2014321	C1	19940615	RU 1990-4743256	19900307
SU 1836328	A3	19930823	SU 1990-4743319	19900312
GB 2241236	A1	19910828	GB 1991-5678	19910318
GB 2241236	B2	19920708		

L52 ANSWER 44 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

GB 2241237 A1 19910828 GB 1991-5679 19910318  
GB 2241237 B2 19920708  
GB 2241238 A1 19910828 GB 1991-5680 19910318  
GB 2241238 B2 19920708  
AU 9173806 A1 19910711 AU 1991-73806 19910325  
AU 632207 B2 19921217  
RU 2017721 C1 19940815 RU 1991-5010110 19911120  
LV 10242 B 19950420 LV 1992-490 19921224  
LT 3673 B 19960125 LT 1993-1427 19931028  
US 5468857 A 19951121 US 1994-191163 19940203  
US 5468857 A 19951121 US 1994-191163 19940203  
PRIORITY APPLN. INFO.: GB 1987-21221 A 19870909


GB 1987-21706 A 19870915  
GB 1988-1485 A 19880122  
GB 1988-6317 A 19880317  
GB 1988-14734 A 19880621  
GB 1988-19878 A 19880822  
US 1988-242760 A3 19880909

OTHER SOURCE(S): CASREACT 111:114869; MARPAT 111:114869

IT 122281-57-6P 122281-58-7P 122297-49-8P  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of, as fungicide)


RN 122281-57-6 CAPIUS  
CN Benzenecetic acid, 2-[3-(2-fluorophenoxy)phenoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 122281-58-7 CAPIUS  
CN Benzenecetic acid, 2-[3-(4-fluorophenoxy)phenoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.





RN 122297-49-8 CAPLUS  
 CN Benzeneacetic acid, 2-[(3-fluorophenoxy)phenoxy]- $\alpha$ -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)

RN 113397-96-9 CAPLUS  
 CN Phenol, 2,4,5-trifluoro-6-methyl-3-[(2,4,5-trifluoro-6-methyl-3-(2,3,4,5-tetrafluoro-6-methylphenoxy)phenoxy)]- (9CI) (CA INDEX NAME)




AB C6F5OH reacted with DMSO-DCC-H3PO4 below room temperature to give C6F5OCH2SMe and the [2,3]-rearrangement products I (R = F, C6F5O). Under similar conditions, 2,3,5,6-F4C6HOH and perfluoro-2-naphthol gave similar ethers and rearrangement products (e.g., dihydronaphthalene II; Z = S). Reaction of polyfluoroarenes with DMSO-(CF3CO)2O at low temps. followed by deprotonation with Et3N resulted in more efficient rearrangement reactions. 4-Bromo-3,5,6-trifluoro-2-pyridinol gave the corresponding ether and both C and N rearrangement products; 2,4,5,6-tetrafluoro-3-pyridinol gave products resulting from the overall replacement of the 2-F by CHO and by CH(SMe)2; both 2,5,6-trifluoro-4-pyrimidinol and 5-fluoro-4,6-dimethoxy-2-pyrimidinol gave simple rearrangement products. NaBH4 reduction and Raney Ni desulfurization of some of the rearranged

compds. gave phenolic products. Reaction of sulfone II (Z = SO2) with DBU effected overall replacement of the 1-F in the starting perfluoro-2-naphthol to give hydroxy aldehyde derivs. III [R1 = CHO, CH(SO2Me)2].

ACCESSION NUMBER: 1988:131238 CAPLUS  
 DOCUMENT NUMBER: 108:131238  
 TITLE: Reactions of polyfluoroarenes and -heteroarenes with activated dimethyl sulfoxide. Facile

[2,3]-sigmatropic rearrangement reactions giving dearomatized products  
 AUTHOR(S): Brooke, Gerald M.; Ferguson, J. A. K. Jamie  
 CORPORATE SOURCE: Dep. Chem., Science Lab., Durham, DH1 3LE, UK  
 SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1972-1999) (1987), (9), 2091-7  
 CODEN: JCPRB4; ISSN: 0300-922X

DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 108:131238  
 IT 113397-96-9P




AB Herbicidal (no data) nitrophenoxylanilines I [R-R2 = H, alkyl, alkenyl, alkynyl, alkoxy, cyano, F3C, MeCO, halo; R3 = H, halo, Me; R4 = H, halo; R5 = H, alkyl, alkenyl, amino, (un)substituted alkyl, PhCH2] were prepared

Thus, 2,3,4-trichloro-6-nitrophenyl ether was treated with PhOH in the presence of Bu4N+ HSO4- to give 97.4% 3,2,4-trichloro-6-nitrophenyl ether. This was treated with NH3 in (CH3OCH2CH2OH)2 at 165° 6h to give 95.8% I (R = R1 = R2 = R4 = R5 = H, R3 = Cl).


ACCESSION NUMBER: 1985:203708 CAPLUS  
 DOCUMENT NUMBER: 102:203708  
 TITLE: Diphenyl ethers  
 INVENTOR(S): Sehring, Richard; Buck, Wolfgang  
 PATENT ASSIGNEE(S): Celamerck G.m.b.H. und Co. K.-G., Fed. Rep. Ger.  
 SOURCE: Ger. Offen., 27 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM.: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3209878	A1	19830929	DE 1982-3209878	19820318
EP 89517	A1	19830928	EP 1983-101949	19830228
EP 89517	B1	19860611		
R: AT, BE, CH, DE, FR, GB, IT, LI, NL				
AT 20340	E	19860615	AT 1983-101949	19830228
JP 58170736	A2	19831007	JP 1983-44033	19830315
JP 04080898	B4	19921221		
CA 1264763	A1	19900123	CA 1983-423736	19830316
PRIORITY APPLN. INFO.:			DE 1982-3209878	A 19820318
			EP 1983-101949	A 19830228




OTHER SOURCE(S): CASREACT 102:203708  
 IT 88483-48-1P 88483-51-EP 88483-46-3P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and aminolysis of)  
 RN 88483-48-1 CAPLUS  
 CN Benzene, 2-chloro-1,3-bis(4-fluorophenoxy)-4-nitro- (9CI) (CA INDEX NAME)



RN 88483-51-6 CAPLUS  
CN Benzene, 1,3-bis(4-fluorophenoxy)-2-methyl-4-nitro- (9CI) (CA INDEX NAME)



RN 88483-66-3 CAPLUS  
CN Benzene, 1,3-bis(2-fluorophenoxy)-2-methyl-4-nitro- (9CI) (CA INDEX NAME)



AB Tetrahydrofurans I (R, R2 = H, halogen; R1 = halogen, CF3) were prepared Thus, 3-tetrahydrofuranol was treated with 2,4-Cl(CF3)C6H3OC6H3(NO2)2-3,4 to give 65.6% I (R = Cl, R1 = NO2, R2 = H) which at 1 g/are post-emergence

gave total control of, e.g., Echinochloa crusgalli in rice.

ACCESSION NUMBER: 1982:6554 CAPLUS

DOCUMENT NUMBER: 96:6554

TITLE: Tetrahydrofuran derivatives and their use as herbicides

INVENTOR(S): Yoshimoto, Takeo; Toyama, Teruhiko; Igarashi, Keiichi;

Ura, Masaaki; Enomoto, Yuji; Funakoshi, Yasunobu;

Hojo, Yoshikata

Mitsui Toatsu Chemicals, Inc., Japan

Fr. Demande, 20 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2471979	A1	19810626	FR 1980-26654	19801216
FR 2471979	B1	19830812		
JP 56086179	A2	19810713	JP 1979-162711	19791217
JP 59051950	B4	19841217		
US 4308051	A	19811229	US 1980-216609	19801215
BR 8008236	A	19810707	BR 1980-8236	19801216
DE 3047629	A1	19810917	DE 1980-3047629	19801217
DE 3047629	C2	19890105		

PRIORITY APPLN. INFO.:

OTHER SOURCE(S): CASREACT 96:6554


IT 60793-43-3

RL: RCT (Reactant); RACT (Reactant or reagent)

(reaction of, with hydroxytetrahydrofuran)

RN 60793-43-3 CAPLUS

CN Benzene, 2,4-bis(2,4-dichloro-6-fluorophenoxy)-1-nitro- (9CI) (CA INDEX NAME)



AB 2,4-Diphenoxynitrobenzenes I (R = halo, alkyl, CF3, NO2; n = 0-3) were prepared by treating RnC6H5-nOH (II) with 3,4-dinitrochlorobenzene (III)

in the presence of alkali. III is more readily available than 2,4-dichloronitrobenzene. Thus, a mixture of II (Rn = 2,4-Cl2) 35.9, III 20.3, and K2CO3 30.4 parts in DMF was heated at 70° for 2 hr and at 140° for 3 hr to give 90.3% I (Rn = 2,4-Cl2). Me2SO, sulfolane, or diglyme was also a good solvent, whereas ethylene glycol instead of DMF lowered the yield. Among 9 addnl. I prepared were I (Rn = H, 2,4-Me(Cl), 2,4-Cl(CF3), and 2,4,6-Cl3.

ACCESSION NUMBER: 1976:539665 CAPLUS

DOCUMENT NUMBER: 85:139665

TITLE: 2,4-Diphenoxynitrobenzenes

INVENTOR(S): Yoshimoto, Takeo; Harayama, Takeo; Ura, Masaaki

Mitsui Toatsu Chemicals, Inc., Japan

Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 51088931	A2	19760804	JP 1975-13958	19750204
JP 55030702	B4	19800813		

PRIORITY APPLN. INFO.:


IT 60793-43-3P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

RN 60793-43-3 CAPLUS

CN Benzene, 2,4-bis(2,4-dichloro-6-fluorophenoxy)-1-nitro- (9CI) (CA INDEX NAME)




L52 ANSWER 49 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The PbO<sub>2</sub> oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).

ACCESSION NUMBER: 1974:520135 CAPIUS  
DOCUMENT NUMBER: 81:120135  
TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol  
AUTHOR(S): Deniville, Leon; Huynh Anh Hoa  
CORPORATE SOURCE: Lab. Chim. Text. Tintoriale, Conservatoire Natl. Arts  
SOURCE: Metiers, Paris, Fr.  
Bulletin de la Societe Chimique de France (1974), [3-4, Pt. 2], 487-90  
CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal  
LANGUAGE: French  
IT 53279-71-3P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 53279-71-3 CAPIUS  
CN 1,4-Benzenediol, 2,3,5,6-tetrakis(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



L52 ANSWER 50 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.


AB The reaction of (C6Cl<sub>5</sub>)<sub>2</sub>O with KF in an autoclave did not give the expected (C6F<sub>5</sub>)<sub>2</sub>O; only C6F<sub>6</sub>, C6F<sub>5</sub>Cl, C6F<sub>4</sub>Cl<sub>2</sub>, C6F<sub>3</sub>Cl<sub>3</sub>, and C6F<sub>2</sub>Cl<sub>4</sub> were obtained. The same products were obtained from C6Cl<sub>6</sub>. The reaction of C6F<sub>5</sub>NO<sub>2</sub> with 2,3,5,6,4-F<sub>4</sub>XC<sub>6</sub>OK (X is Me, H, Br, CF<sub>3</sub>, NO<sub>2</sub>) gave 46-801 2,3,5,6,4-F<sub>4</sub>XC<sub>6</sub>OC<sub>6</sub>(NO<sub>2</sub>)-F<sub>4</sub>-4,2,3,5,6 which were reduced to the corresponding amino-derivs. The amino derivs. were aminated to diamino ethers. The oxidation of 4,4'-diaminooctafluorodiphenyl ether (I) with CF<sub>3</sub>-CO<sub>2</sub> h gave 4,4'-dinitrooctafluorodiphenyl ether (II), which was reacted with NH<sub>3</sub>(g) to the 3,3'-diamino analog (III) of II. The reduction of

III gave 3,3',4,4'-tetraaminohexafluorodiphenyl ether.  
4-Bromooctafluorodiphenyl ether or 4,4'-dibromooctafluorodiphenyl ether were also prepared from 4-aminooctafluorodiphenyl or I, resp., via the

San dmeier reaction.  
ACCESSION NUMBER: 1970:110946 CAPIUS  
DOCUMENT NUMBER: 72:110946  
TITLE: Aromatic fluoro derivatives. XXXVIII. Derivatives of polyfluorodiphenyl ether

AUTHOR(S): Kobrina, L. S.; Furin, G. G.; Yakobson, G. G.  
CORPORATE SOURCE: Novosibirsk. Inst. Org. Khim., Novosibirsk, USSR  
SOURCE: Zhurnal Organicheskoi Khimii (1970), 6(2), 340-5  
CODEN: ZORXAE; ISSN: 0514-7492  
DOCUMENT TYPE: Journal  
LANGUAGE: Russian  
IT 28096-26-6P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 28096-26-6 CAPIUS  
CN Benzene, 1,2,3,5-tetrafluoro-4,6-bis(2,3,5,6-tetrafluoro-4-nitrophenoxy)- (8CI) (CA INDEX NAME)




L52 ANSWER 51 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

AB NaOC<sub>6</sub>F<sub>5</sub> reacted with C<sub>6</sub>F<sub>5</sub>CN in Me<sub>2</sub>CO to give 4-cyanooctafluorodiphenyl ether (I), while in AcNMe<sub>2</sub> solvent, isomeric and polysubstituted products accompanied the formation of I. C<sub>6</sub>F<sub>5</sub>CN is more reactive than C<sub>6</sub>F<sub>5</sub>CF<sub>3</sub> toward NaOC<sub>6</sub>F<sub>5</sub>, which reacts with p-CF<sub>3</sub>C<sub>6</sub>F<sub>4</sub>CN to give 2,6-bis(pentafluorophenoxy)-4-trifluoromethylfluorobenzonitrile. The observations suggest that a CN substituent is a potent activator for

ortho and para nucleophilic substitution reactions on fluorinated aromatic compds.


ACCESSION NUMBER: 1968:104689 CAPIUS  
DOCUMENT NUMBER: 68:104689  
TITLE: Reactions of fluoroaromatic nitriles with sodium pentafluorophenolate  
AUTHOR(S): De Pasquale, Ralph J.; Tamborski, Christ  
CORPORATE SOURCE: Air Force Mater. Lab., Wright-Patterson Air Force Base, OH, USA  
SOURCE: Journal of Organic Chemistry (1968), 33(4), 1658-61  
CODEN: JOCEAH; ISSN: 0022-3263  
DOCUMENT TYPE: Journal  
LANGUAGE: English

IT 15963-72-1P 16031-36-0P 16065-60-4P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 15963-72-1 CAPIUS  
CN Benzonitrile, 2,3,5-trifluoro-4,6-bis(pentafluorophenoxy)- (8CI, 9CI) (CA INDEX NAME)




RN 16031-36-0 CAPIUS  
CN Benzonitrile, 3,5-difluoro-2,4,6-tris(pentafluorophenoxy)- (8CI, 9CI) (CA INDEX NAME)

L52 ANSWER 51 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




RN 16065-60-4 CAPIUS  
CN p-Tolunitrile, alpha,alpha,3,5-pentafluoro-2,6-bis(pentafluorophenoxy)- (8CI) (CA INDEX NAME)




L52 ANSWER 52 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A number of perfluorinated and partially fluorinated phenyl and polyphenyl ethers were synthesized, characterized for thermal stability, fire resistance, and viscosity, and compared with their H analogs to assess the potential use of this class of compds. as functional fluids. Without exception, polyfluorination and perfluorination lower thermal stability; the decrease in stability depends on the position and number of fluorine substituents. The autoignition temperature and fire resistance are not improved over the H analog, and viscosity is degraded. These data coupled with the comparatively high melting points do not suggest a bright future for this class of compds. as useful functional fluids. 6 references.

ACCESSION NUMBER: 1968:77890 CAPLUS  
 DOCUMENT NUMBER: 68:77890  
 TITLE: Synthesis, thermal stability, flammability, and viscosity of some partially fluorinated and perfluorinated aromatic and polyaromatic ethers  
 AUTHOR(S): Richardson, George Albert; Blake, Edward S.  
 CORPORATE SOURCE: Monsanto Res. Corp., Dayton, OH, USA  
 SOURCE: Industrial & Engineering Chemistry Product Research and Development (1968), 7(1), 17-21  
 CODEN: IEPRA6; ISSN: 0196-4321  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 5026-82-4P 17742-25-5P 17742-29-9P  
 17742-30-2P 17742-35-7P 17742-36-8P  
 17742-37-9P 17742-40-4P 17742-42-6P  
 19228-15-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
 RN 5026-82-4 CAPLUS  
 CN Benzene, 1-(m-fluorophenoxy)-3-(m-phenoxyphenoxy)- (7CI, 8CI) (CA INDEX NAME)




RN 17742-25-5 CAPLUS  
 CN Benzene, 1-(pentafluorophenoxy)-3-phenoxy- (8CI) (CA INDEX NAME)




RN 17742-29-9 CAPLUS  
 CN Benzene, 1-(pentafluorophenoxy)-3-(m-phenoxyphenoxy)- (8CI) (CA INDEX NAME)


L52 ANSWER 52 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 17742-40-4 CAPLUS  
 CN Biphenyl, 2,2',3,3',5,5',6,6'-octafluoro-4,4'-bis(m-phenoxyphenoxy)- (8CI)  
 (CA INDEX NAME)




RN 17742-42-6 CAPLUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(m-phenoxyphenoxy)phenoxy- (8CI)  
 (CA INDEX NAME)




RN 19228-15-0 CAPLUS  
 CN Benzene, 1,2,4-trifluoro-3,5,6-tris(m-phenoxyphenoxy)- (7CI, 8CI) (CA INDEX NAME)


L52 ANSWER 52 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 17742-30-2 CAPLUS  
 CN Benzene, 1-(m-phenoxyphenoxy)-3-(2,3,5,6-tetrafluorophenoxy)- (8CI) (CA INDEX NAME)



RN 17742-35-7 CAPLUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(m-phenoxyphenoxy)- (8CI) (CA INDEX NAME)




RN 17742-36-8 CAPLUS  
 CN Benzene, 1,4-difluoro-2,5-bis(m-phenoxyphenoxy)- (8CI) (CA INDEX NAME)



RN 17742-37-9 CAPLUS  
 CN Benzene, 1-[m-(pentafluorophenoxy)phenoxy]-3-(m-phenoxyphenoxy)- (8CI)  
 (CA INDEX NAME)

L52 ANSWER 52 OF 54 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)





L52 ANSWER 53 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB 2,3,5,6-Tetrafluoro-O,O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]hydroquinones (I) and 2,4,5,6-tetrafluoro-O,O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]resorcinols (II) are prepared by the

treatment of the compds. of the general formulas C6F5X, where X is CN, CF3, CONH2, and CO2H, with 2,3,5,6-tetrafluorohydroquinone di-Na salt and 2,4,5,6-tetrafluororesorcinol di-Li salt (III), resp. 19F-N.M.R. and ir data for the prepared I and II are given. III is treated with decafluorocyclohexene to give a mixture of 2,4,5,6-tetrafluoro-O,O'-bis(perfluoro-1-cyclohexen-1-yl)resorcinol and 2,4,5,6-tetrafluoro-O-(perfluoro-1-cyclohexen-1-yl)-O'-(perfluoro-2-cyclohexen-1-yl)resorcinol.

ACCESSION NUMBER: 1968:68571 CAPIUS

DOCUMENT NUMBER: 68:68571

TITLE: Reactions of perfluorophenolates with substituted pentafluorobenzenes and perfluorocyclohexene  
AUTHOR(S): De Pasquale, Ralph J.; Tamborski, Christ  
CORPORATE SOURCE: Wright-Patterson Air Force Base, Dayton, OH, USA  
SOURCE: Journal of Organic Chemistry (1968), 33(2), 830-3  
CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal


LANGUAGE: English

IT 14796-04-4P 14901-49-6P 15038-90-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)


RN 14796-04-4 CAPIUS

CN Benzonitrile, 4,4'-[(2,4,5,6-tetrafluoro-m-phenylene)dioxy]bis(2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



RN 14901-49-6 CAPIUS

CN Benzoic acid, 4,4'-[(2,4,5,6-tetrafluoro-m-phenylene)dioxy]bis(2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



RN 15038-90-1 CAPIUS

L52 ANSWER 54 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

AB Polyphenyl ethers containing 4-7 Ph groups and in which one of the terminal groups is substituted by 1, 2, or 5 F atoms and by 2 m-phenoxyphenyl groups are heat-stable compds. with low m.p.s. The mono-F compds. were prepared by condensation of an alkali phenoxide with bromofluorobenzene in the presence of Cu. The penta-F compds. were prepared by condensation of an

alkali phenoxide with hexafluorobenzene. 1,3,4-Trifluorotris(m-phenoxyphenoxy)benzene was prepared by condensation of 3 equivalent of K m-phenoxyphenoxide with hexafluorobenzene. Thus, 58.4 g. m-(m-phenoxyphenoxy)phenol was mixed with 12.4 g. KOH and 35 ml. PhMe; water was removed with the PhMe by azeotropic distillation. The dry phenoxide was dissolved in 100 ml. diglyme and added to a solution of 35 g. m-bromofluorobenzene in 100 ml. diglyme containing 0.5 g. Cu powder, over a period of 2.25 hrs. at 155°. The mixture was then stirred for 25.5 hrs. at 165° cooled, and filtered. The diglyme was distilled and the residue dissolved in ether. The ether solution was washed with acid, dilute alkali, and water, and distillation gave 44.5 g. (59.5% theory) of pure m(m-fluorophenoxy)phenyl m-phenoxyphenyl ether, decomposition 462° b0.15 212°, n25D 1.6094.

ACCESSION NUMBER: 1965:487872 CAPIUS

DOCUMENT NUMBER: 63:87872

ORIGINAL REFERENCE NO.: 63:16116t-g

TITLE: Heat-stable hydraulic fluids

PATENT ASSIGNEE(S): Monsanto Co.

SOURCE: 18 pp.

DOCUMENT TYPE: Patent

LANGUAGE: Unavailable

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NL 6414074	---	19650608	NL	---
PRIORITY APPLN. INFO.:			US	19631204


IT 5026-81-3, Benzene, 1,2,4-trifluoro-3,5,6-tris(m-phenoxyphenoxy)-  
5026-82-4, Benzene, 1-(m-fluorophenoxy)-3-(m-phenoxyphenoxy)-  
(as hydraulic fluid)

RN 5026-81-3 CAPIUS

CN Benzene, 1,3,4-trifluoro-2,5,6-tris(m-phenoxyphenoxy)- (8CI) (CA INDEX NAME)

L52 ANSWER 53 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN


CN Benzene, 1,2,3,5-tetrafluoro-4,6-bis[(a,a,a,2,3,5,6-heptafluoro-p-tolyl)oxy]- (8CI) (CA INDEX NAME)



(Continued)


L52 ANSWER 54 OF 54 CAPIUS COPYRIGHT 2005 ACS on STN

(Continued)



RN 5026-82-4 CAPIUS

CN Benzene, 1-(m-fluorophenoxy)-3-(m-phenoxyphenoxy)- (7CI, 8CI) (CA INDEX NAME)



```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          271.71      3828.46

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          -39.42      -265.72
```

FILE 'REGISTRY' ENTERED AT 20:28:26 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
 DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
 information enter HELP PROP at an arrow prompt in the file or refer  
 to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=> FIL STNGUIDE
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          1.29      3829.75

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          0.00      -265.72
```

FILE 'STNGUIDE' ENTERED AT 20:30:20 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Jan 7, 2005 (20050107/UP).

```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          0.06      3829.81

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          0.00      -265.72
```

FILE 'REGISTRY' ENTERED AT 20:30:30 ON 12 JAN 2005




1:CLASS 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
 11:Atom 12:Atom 13:CLASS 14:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom  
 21:Atom 22:Atom 23:CLASS 25:CLASS 26:CLASS 28:CLASS 29:CLASS 30:CLASS  
 31:CLASS 32:CLASS 33:CLASS 35:CLASS

L53 STRUCTURE UPLOADED

=> d query

L53 STR



G1 N,X

Structure attributes must be viewed using STN Express query preparation.

=> s 153

SAMPLE SEARCH INITIATED 20:31:18 FILE 'REGISTRY'  
 SAMPLE SCREEN SEARCH COMPLETED - 110 TO ITERATE

100.0% PROCESSED 110 ITERATIONS  
 SEARCH TIME: 00.00.01

7 ANSWERS

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
 BATCH \*\*COMPLETE\*\*  
 PROJECTED ITERATIONS: 1571 TO 2829  
 PROJECTED ANSWERS: 7 TO 298

L54 7 SEA SSS SAM L53

=> s .153 full

FULL SEARCH INITIATED 20:31:24 FILE 'REGISTRY'  
 FULL SCREEN SEARCH COMPLETED - 2122 TO ITERATE

100.0% PROCESSED 2122 ITERATIONS  
 SEARCH TIME: 00.00.01

103 ANSWERS

L55 103 SEA SSS FUL L53

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
161.76	3991.57

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION

CA SUBSCRIBER PRICE

0.00 -265.72

FILE 'CAPLUS' ENTERED AT 20:31:30 ON 12 JAN 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 155

L56            89 L55


=> d 56 1-89 abs ibib hitstr

L56 ANSWER 56 OF 89 CAPLUS COPYRIGHT 2005 ACS ON STN


AB In the presence of fluoride ion, per- or polyfluoroarom. siloxanes C6F5OSi(CH<sub>3</sub>)<sub>3</sub> (1), (CH<sub>3</sub>)<sub>3</sub>SiOC6F4OSi(CH<sub>3</sub>)<sub>3</sub> (2), 1-fluoro-2,3-bis((trimethylsilyl)oxy)benzene (3), and (CH<sub>3</sub>)<sub>3</sub>SiOC(CF<sub>3</sub>)<sub>2</sub>C6F4(CF<sub>3</sub>)<sub>2</sub>OSi(CH<sub>3</sub>)<sub>3</sub> (4) are useful transfer reagents for the synthesis of mono- and diethers. Siloxane 1 forms RfOCF<sub>2</sub>Br (5), RfOCF<sub>3</sub> (6), RfOCH<sub>2</sub>ORf (7), RfOC(CH<sub>3</sub>)<sub>2</sub>ORf (8), RfOC<sub>2</sub>SO (9), RfOC<sub>2</sub>CO (10), C<sub>3</sub>N<sub>3</sub>(ORf)<sub>3</sub> (11), RfOC(O)(CF<sub>2</sub>)<sub>3</sub>C(O)ORf (12), CF<sub>3</sub>SO<sub>2</sub>ORf (13), [cyclic] CF<sub>2</sub>C(ORf):C(ORf)CF<sub>2</sub> (14), and [cyclic] CF<sub>2</sub>C(ORf):C(F)CF<sub>2</sub> (15) (Rf = C6F<sub>5</sub>) with CF<sub>2</sub>Br<sub>2</sub>, CF<sub>2</sub>Br<sub>3</sub>, CH<sub>2</sub>Br<sub>2</sub>, Br<sub>2</sub>CHCHBr<sub>2</sub>, SOF<sub>2</sub>, COF<sub>2</sub>, (CNF)<sub>3</sub>, ClC(O)(CF<sub>2</sub>)<sub>3</sub>C(O)Cl, CF<sub>3</sub>SO<sub>2</sub>F, 1,2-dichlorotetrafluorocyclobutene, and perfluorocyclobutene, resp. Compound 5 is readily converted to C6F<sub>5</sub>OCF<sub>2</sub>Si(CH<sub>3</sub>)<sub>3</sub> (16) with hexaethylphosphorous triamide and (CH<sub>3</sub>)<sub>3</sub>SiCl in benzonitrile. With C6F<sub>5</sub>CH, CF<sub>3</sub>CF<sub>3</sub>, C<sub>6</sub>F<sub>5</sub>SH, CH<sub>3</sub>I, perfluorocyclobutene, CF<sub>3</sub>C(O)Cl and (CNF)<sub>3</sub>, 2 forms diethers p-CNC6F4OC6F4OC6F4C<sub>2</sub>N-p (17), p-CF<sub>3</sub>C6F4OC6F4OC6F4CF<sub>3</sub>-p (18), NC6F4OC6F4OC6F4N (19), CH<sub>3</sub>OC6F4OCCH<sub>3</sub> (20), and C2F<sub>5</sub>C(O)OC6F4OC(O)C2F<sub>5</sub> (21), resp. Reaction of 3 with 1,2-diiodotetrafluorobenzene in diglyme gives 1,4,9-trifluoro-2,3-dilodophenazine (22). Disiloxane 4 with C6H<sub>5</sub>CH<sub>2</sub>Br, CH<sub>3</sub>I, C6F<sub>5</sub>CH<sub>2</sub>Br, and COF<sub>2</sub> results in C6H<sub>5</sub>CH<sub>2</sub>OC(CF<sub>3</sub>)<sub>2</sub>C6F4C(CF<sub>3</sub>)<sub>2</sub>OCCH<sub>2</sub>C6H<sub>5</sub> (23), CH<sub>3</sub>OC(CF<sub>3</sub>)<sub>2</sub>C6F4C(CF<sub>3</sub>)<sub>2</sub>OCCH<sub>3</sub> (24), C6F<sub>5</sub>CH<sub>2</sub>OC(CF<sub>3</sub>)<sub>2</sub>C6F4C(CF<sub>3</sub>)<sub>2</sub>OCCH<sub>2</sub>C6F<sub>5</sub> (25), and [cyclic] C(O)OC(CF<sub>3</sub>)<sub>2</sub>C6F4C(CF<sub>3</sub>)<sub>2</sub>OC(CF<sub>3</sub>)<sub>2</sub>OC(CF<sub>3</sub>)<sub>2</sub>C6F4C(CF<sub>3</sub>)<sub>2</sub>OC(CF<sub>3</sub>)<sub>2</sub> (26), resp. These materials are thermally and hydrolytically stable and are formed in high yields.

ACCESSION NUMBER: 1995:269201 CAPLUS  
DOCUMENT NUMBER: 122:133262  
TITLE: Per- and Polyfluoroaryl Mono- and Disiloxanes as Transfer Reagents in the Synthesis of Highly Fluorinated Mono- and Diethers  
AUTHOR(S): Patel, Mimesh R.; Chen, Janguo; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 83844-2343, USA  
SOURCE: Inorganic Chemistry (1995), 34(1), 13-17  
CODEN: INOCAL; ISSN: 0020-1669  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 122:133262  
IT 15053-71-1P 15077-30-2P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 15053-71-1 CAPLUS  
CN Benzonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)]

L56 ANSWER 56 OF 89 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)




RN 15077-30-2 CAPLUS  
CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)



L56 ANSWER 1 OF 89 CAPLUS COPYRIGHT 2005 ACS ON STN

GI



AB This invention pertains to a method for producing bis(fluorine-containing phthalic anhydride) derivs. I [wherein m = 1-3; n = 1-3; Z = a single bond or a spacer], which comprises reacting a fluorine-containing dinitrile with a fluorine-containing hydroquinone, followed by hydrolysis and dehydration reaction. For example, 3,4,5,6-tetrafluorophthalonitrile was reacted with tetrafluorohydroquinone in Me iso-Bu ketone in the presence of KF to give the bisphthalonitrile (97%). The bisphthalonitrile was hydrolyzed in propionic acid in the presence of H<sub>2</sub>SO<sub>4</sub> to afford the bis(phthalic acid) (75%). The bis(phthalic acid) was treated with acetic anhydride to provide the bis(phthalic anhydride) (98%).

ACCESSION NUMBER: 2004:1127358 CAPLUS  
TITLE: Process for preparation of bis(fluorine-containing phthalic anhydride)  
INVENTOR(S): Masuda, Go; Okumura, Yasunori; Nishimae, Shinji; Naraya, Toshihiro  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: PCT Int. Appl., 39 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004/11026	A1	20041223	WO 2004-JP8829	20040617
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RV: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPL. INFO.:		JP 2003-172046		A 20030617
		JP 2003-201123		A 20030724

IT 143363-91-1P 143363-92-2P 143376-50-5P  
RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic)


Page 364

L56 ANSWER 1 OF 89 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
preparation); PREP (Preparation); RACT (Reactant or reagent) (intermediate; prepn. of bis(fluorine-contg. phthalic anhydride))


RN 143363-91-1 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)]




RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT




AB The polyimides have repeating units I (R1 = tetravalent aromatic group containing F atoms directly bonded to C atoms on aromatic rings; R2 = divalent organic

group containing alicyclic structure). The polyimides are useful for red-emitting electroluminescent devices. The polyimides show high fluorescent intensity and good heat and H2O resistance.


ACCESSION NUMBER: 2004:931701 CAPLUS  
DOCUMENT NUMBER: 141:403244  
TITLE: Fluorescent polyimides and organic light-emitting devices using them  
INVENTOR(S): Ando, Shinji; Urano, Yuichi  
PATENT ASSIGNEE(S): Rikogaku Shinkokai, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 33 pp.  
CODEN: JI004F  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004307657	A2	20041104	JP 2004-91565	20040326
PRIORITY APPLN. INFO.:			JP 2003-84629	A 20030326

IT 790229-67-3P 790229-68-4P 790229-69-5P  
790229-70-8P 790229-71-9P 790229-72-0P  
790229-73-1P 790229-74-2P  
RL: DEV (Device component use); IMF (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses) (fluorescent fluorinated polyimides with good heat resistance for light-emitting devices)  
RN 790229-67-3 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-methylenebis(cyclohexanamine) (9CI) (CA INDEX NAME)  
CH 1  
CRN 143363-91-1  
CMF C22 F10 O8




CH 2  
CRN 1761-71-3  
CMF C13 H26 N2




RN 790229-68-4 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-cyclohexanedimethylene-1,4-cyclohexanediyl] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 790229-69-5 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 1,4-cyclohexanediamine (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CMF C22 F10 O8



CH 2  
CRN 3114-70-3  
CMF C6 H14 N2




RN 790229-70-8 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-cyclohexanedimethylene-1,4-cyclohexanediyl] (9CI) (CA INDEX NAME)



RN 790229-71-9 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis(cyclohexanediyl) (9CI) (CA INDEX NAME)


CM 1

CRN 473906-71-7  
CMF C15 H24 F6 N2




CM 2

CRN 143363-91-1  
CMF C22 F10 O8




RN 790229-72-0 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-




RN 790229-74-2 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)[2,2'-bis(trifluoromethyl)[1,1'-bicyclohexyl]-4,4'-diyl]] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B

PAGE 1-A




PAGE 1-B



RN 790229-73-1 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)[1,1'-bicyclohexyl]-4,4'-diamine (9CI) (CA INDEX NAME)

CM 1

CRN 551950-80-2  
CMF C14 H22 F6 N2



CM 2

CRN 143363-91-1  
CMF C22 F10 O8

L56 ANSWER 3 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
AB The polyimide is that obtained by polymerization of (a) a dianhydride of a diacid having F-containing aromatic rings and (b) a diamine having sulfonic acid groups.  
The composition contains the polyimide and an agent for addition of proton conductivity.  
The ionic conductor contains the polyimide or the composition A fluorinated polyamic acid obtained by polymerization of the above dianhydride and the diamine is also claimed. The polyimide showing enhanced resistance to heat, boiling water, and radicals, is suitable for an electrolytic film in a fuel cell.

ACCESSION NUMBER: 2004:898860 CAPLUS  
DOCUMENT NUMBER: 141:373556  
TITLE: Sulfonic acid group-substituted fluorinated polyimide,  
composition containing the polyimide, and ionic conductor  
INVENTOR(S): Nishichi, Ai; Matsumoto, Takeshi; Akutagawa, Hironobu;  
Omote, Kazuishi; Yoshida, Masaya  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 27 pp.  
DOCUMENT TYPE: CODEN: JKOXAF  
LANGUAGE: Patent  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION: Japanese

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004300208	A2	20041028	JP 2003-92639	20030328
PRIORITY APPLN. INFO.:			JP 2003-92639	20030328


IT 777865-72-2P  
RL: IMP (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(sulfonic acid group-substituted fluorinated polyimide for ionic conductor)

RN 777865-72-2 CAPLUS  
CN [1,1'-Biphenyl]-2,2'-disulfonic acid, 4,4'-diamino-5,5'-dimethyl-, disodium salt, polymer with 4,4'-oxybis(benzenamine) and 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8







CH 2


CRN 67027-35-4  
CHF C14 H16 N2 O6 S2 . 2 Na

● 2 Na

CH 3


CRN 101-80-4  
CHF C12 H12 N2 OCRN 143363-91-1  
CHF C22 F10 O8

CH 2

CRN 1198-63-6  
CHF C6 H4 F4 N2

RN 143433-45-8 CAPIUS

CH  
Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 484016-84-4 CAPIUS  
CH  
1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 5-chloro-2,4,6-

AB Title method comprises forming on a substrate a coating film of a fluorine-containing polyimide precursor (e.g., 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer, polyamic acid) in an atmospheric having a relative humidity thereof adjusted to a level of not more than 50 RH and then subjecting the fluorine-containing polyimide precursor to heat 30 treatment thereby forming a fluorine-containing polyimide film. The formation of the coating film is carried out by a spin coating method which comprises dropping the fluorine-containing polyimide precursor onto the substrate and spin coating the fluorine-containing polyimide precursor. The polyimide film exhibits excellent in heat resistance, resistance to chems., water repellency, dielec. properties, elec. properties, and optical properties.

ACCESSION NUMBER: 2004:732270 CAPIUS  
DOCUMENT NUMBER: 141:244610  
TITLE: Method and apparatus for production of fluorine-containing polyimide film  
INVENTOR(S): Tajiri, Kozo; Goto, Shinichi; Asako, Yoshinobu;  
PATENT ASSIGNEE(S): Kawada, Kumiko  
SOURCE: Nippon Shokubai Co., Ltd., Japan  
Eur. Pat. Appl., 24 pp.  
CODEN: EPOKDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1454945	A1	20040908	EP 2004-251311	20040305
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK,				
HR JP 2004269591	A2	20040930	JP 2003-59331	20030306
JP 2004267848	A2	20040930	JP 2003-59332	20030306
US 2004234686	A1	20041125	US 2004-795122	20040305
PRIORITY APPLN. INFO.:			JP 2003-59331	A 20030306
			JP 2003-59332	A 20030306

IT 143376-21-0P 143433-45-8P 484016-84-4P  
484016-86-6P 750397-62-7P 750397-63-8P  
RL: IMP (Industrial manufacture); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(method and apparatus for production of fluorine-containing polyimide film)  
RN 143376-21-0 CAPIUS  
CH 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CH 1

trifluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1


CRN 474805-39-5  
CHF C6 H4 Cl F3 N2

CH 2


CRN 143363-91-1  
CHF C22 F10 O8

RN 484016-86-6 CAPIUS



CH  
Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(5-chloro-2,4,6-trifluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



L56 ANSWER 4 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 RN 750597-62-7 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-(9H-fluoren-9-ylidene)bis(benzenamine) (9CI) (CA INDEX NAME)  
 CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8



CH 2  
 CRN 15499-84-0  
 CHF C25 H20 N2




RN 750597-63-8 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)-1,4-phenylene-9H-fluoren-9-ylidene-1,4-phenylene] (9CI) (CA INDEX NAME)

L56 ANSWER 5 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A new poly(arylene ether) high temperature polymer was synthesized. The polymer was converted into ionically conductive polymer by sulfonation and some of the fundamental phys. properties of sulfonated and nonsulfonated materials were evaluated.

ACCESSION NUMBER: 2004:670343 CAPLUS  
 DOCUMENT NUMBER: 141:424520  
 TITLE: New versatile and thermally stable poly(arylene ether)  
 AUTHOR(S): Bouguettaya, Mohamed; Basheer, Rafil A.  
 CORPORATE SOURCE: Delphi research labs, Delphi Corp., Shelby Township, MI, 48315, USA  
 SOURCE: Polymeric Materials: Science and Engineering (2004), 91, 492  
 CODEN: PMSEGD; ISSN: 0743-0515  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal: (computer optical disk)  
 LANGUAGE: English

IT 794592-38-4P  
 RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and characterization of new versatile and thermally stable poly(arylene ether))

RN 794592-38-4 CAPLUS  
 CN Poly[oxy(1,1':4',1''-terphenyl)-2',5'-diyoxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)




IT 794592-38-4DP, sulfonated  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (preparation and characterization of new versatile and thermally stable poly(arylene ether))


RN 794592-38-4 CAPLUS  
 CN Poly[oxy(1,1':4',1''-terphenyl)-2',5'-diyoxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

L56 ANSWER 4 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


PAGE 1-A



PAGE 1-B



L56 ANSWER 5 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L56 ANSWER 6 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB The compns. containing repeating units of AD(R) [A = F-substituted (O- or S-containing) hydrocarbylene; D = F-substituted (O- or S-containing) trivalent hydrocarbon group; R = Si(OR1)(OR2)R3; R1, R2 = hydrocarbyl (each b.p. of R1OH and R2OH under normal pressure <250°); R3 = (F-substituted) double bond-terminated hydrocarbyl] show n and dielec. constant of their cured products 1.350-1.600 and 2.00-4.00, resp. The compns. give cured products with adjustable n and dielec. constant, and improved mech. properties and solvent resistance.

ACCESSION NUMBER: 2004:411677 CAPLUS  
DOCUMENT NUMBER: 140:431126  
TITLE: Manufacture of silicon-containing curable polymer compositions for planar optical waveguides and wiring boards

INVENTOR(S): Florence, Corey Nawarage  
PATENT ASSIGNEE(S): Fujitsu Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
CODEN: J000AF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004143280	A2	20040520	JP 2002-309280	20021024
PRIORITY APPLN. INFO.: JP 2002-309280 20021024				

IT 691906-05-SDP, reaction products with allyltrichlorosilane and ethanol 691906-07-TDP, reaction products with allyltrichlorosilane and ethanol  
RL: DEV (Device component use); IMF (Industrial manufacture); PREP (Preparation); USES (Uses)  
(crosslinked: manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)

RN 691906-05-5 CAPLUS  
CN 1,3-Benzenedicarbonyl dichloride, 5-bromo-, polymer with 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)

CM 1

CRN 691906-04-4  
CMF C20 H4 C12 F8 O4



L56 ANSWER 6 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

CM 2


CRN 461-96-1  
CMF C6 H3 Br F2




IT 691906-02-2P 691906-03-3P 691906-04-4P  
691906-05-5P 691906-06-6P 691906-07-7P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation);

RACT (Reactant or reagent)  
(manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)

RN 691906-02-2 CAPLUS  
CN Benzonitrile, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



RN 691906-03-3 CAPLUS  
CN Benzoic acid, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)




RN 691906-04-4 CAPLUS  
CN Benzoyl chloride, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)

L56 ANSWER 6 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


CM 2

CRN 83558-87-6  
CMF C15 H12 F6 N2 O2



CM 3


CRN 57863-69-1  
CMF C8 H3 Br C12 O2




RN 691906-07-7 CAPLUS  
CN Phenol, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)  
with 1-bromo-3,5-difluorobenzene (9CI) (CA INDEX NAME)

CM 1

CRN 691906-06-6  
CMF C18 H6 F8 O4




L56 ANSWER 6 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 691906-05-5 CAPLUS  
CN 1,3-Benzenedicarbonyl dichloride, 5-bromo-, polymer with 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)


CM 1

CRN 691906-04-4  
CMF C20 H4 C12 F8 O4




CM 2

CRN 83558-87-6  
CMF C15 H12 F6 N2 O2




CM 3

CRN 57863-69-1  
CMF C8 H3 Br C12 O2




RN 691906-06-6 CAPLUS  
CN Phenol, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



RN 691906-07-7 CAPLUS  
CN Phenol, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro-, polymer with 1-bromo-3,5-difluorobenzene (9CI) (CA INDEX NAME)

CM 1

CRN 691906-06-6  
CMF C18 H6 F8 O4



CM 2

CRN 461-96-1  
CMF C6 H3 Br F2

L56 ANSWER 7 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB Polymers incorporating the triptycene subunit were prepared for the mol.-level design of low dielec. constant (low- $\kappa$ ) materials that can be used to manufacture faster integrated circuits. Triptycenes having restricted rotation by multiple point attachment to the polymer backbone are shown to introduce free volume into the films, thereby lowering their dielec. consts. We describe the use of the triptycene moiety as a rigid and shape-persistent component to introduce free volume into a polymer film.

This approach can produce bulk polymers with overall lower dielec. consts. without necessitating the introduction of pore generators, thus eliminating problems inherent to porous polymers. Triptycenes and higher ipitycenes were shown to increase free volume in polymer films, thereby decreasing the bulk dielec. constant of the material. The polymers do not display any glass transitions, are highly thermally stable, and exhibit low water absorption. All of these characteristics are desired by the semiconductor industry for the next generation of microprocessors and memory for computers to provide insulation in the increasingly shrinking feature sizes of faster microprocessors. We believe these materials are viable candidates for evaluation for and incorporation into future generations of microchips.


ACCESSION NUMBER: 2004:231116 CAPLUS  
DOCUMENT NUMBER: 141:141007  
TITLE: Molecular design of free volume: a route to low- $\kappa$  dielectric materials  
AUTHOR(S): Long, Timothy M.; Swager, Timothy M.  
CORPORATE SOURCE: Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA  
SOURCE: Polymeric Materials Science and Engineering (2004), 90, 731-732  
CODEN: PMSEDD; ISSN: 0743-0515  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal; (computer optical disk)  
LANGUAGE: English  
IT 632621-45-5 632621-47-7  
RL: PRP (Properties)

(free volume, dielec., thermal and surface properties of triptycene-based polymers for low dielec. constant materials)  
RN 632621-45-5 CAPLUS  
CN Poly[oxy[6,14-bis[1,1-dimethylethyl]-9,10-dihydro-9,10[1',2']-benzenoanthracene-1,4-diyl]oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)




L56 ANSWER 7 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

(Continued)




PAGE 2-A




RN 632621-47-7 CAPLUS

CN Poly[oxy(5,7,12,14-tetrahydro-5,14[1',2']-7,12[1'',2'']-dibenzenopentacene-6,13-diyl)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



REFERENCE COUNT:  
THIS  
FORMAT


23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR  
RECORD. ALL CITATIONS AVAILABLE IN THE RE

CRN 1190-63-6  
CMF C6 H4 F4 N2



RN 143433-45-8 CAPLUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



AB The waveguides, showing relative refractive index difference 20.24 between cores and cladding layers, comprise cores containing fluorinated polymers or deuterated polymers and cladding layers containing

fluorinated polymers or deuterated polymers with lower F or deuterium content than that of the cores or F- or deuterium-free polymers.

ACCESSION NUMBER: 2004:219234 CAPLUS

DOCUMENT NUMBER: 140:261194

TITLE: Low-cost polymer optical waveguides with low transmission loss

INVENTOR(S): Sakuma, Takeshi; Fujita, Daigo; Ogawa, Hironobu;

PATENT ASSIGNEE(S): Omachi, Koji; Hosoya, Hideyuki

SOURCE: Fujikura Ltd., Japan

Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JPOCAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004085937	A2	20040318	JP 2002-247401	20020827
PRIORITY APPLN. INFO.:			JP 2002-247401	20020827

IT 143376-21-0P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine copolymer

143433-45-8P

RL: IMF (Industrial manufacture); TEM (Technical or engineered material use); PRP (Preparation); USES (Uses)  
(core layer; low-cost polymer optical waveguides with low transmission loss)


RN 143376-21-0 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1

CMF C22 F10 O8



CM 2

AB The temperature dependence of in-plane and out-of-plane refractive indexes

(i.e., thermo-optic coeffs.  $dn/dT$ ) was measured for 7 kinds of aromatic polyimide (PI) films formed on Si substrates. The absolute values of  $dn/dT$

(polarization parallel to the film plane) are significantly larger than  $dn/dT$  (perpendicular to the film plane). The  $dn/dT$  for average refractive

indexes ( $n_{av}$ ) are -94 to -58 ppm/K, independent of film thickness. The anisotropies ( $dn/dT-dn/dT$ ) are -9 to -39 ppm/K. Although the values of

$n_{av}$  are independent of film thickness, the anisotropies slightly increase as the film thickness decreases for flexible PI films. As expected from the temperature derivative of the Lorentz-Lorenz (LL) equation, the amorphous PI

films exhibiting high  $n_{av}$  show large  $dn/dT$ , however, significant anisotropies are observed even for the PI films exhibiting very small birefringence.

ACCESSION NUMBER: 2003:961524 CAPLUS

DOCUMENT NUMBER: 140:153566

TITLE: Anisotropy in thermo-optic coefficients of polyimide films formed on Si substrates

AUTHOR(S): Terui, Yoshiharu; Ando, Shinji

CORPORATE SOURCE: Department of Organic and Polymeric Materials, Tokyo

Institute of Technology, 2-12-1 Ookayama, Meguro-ku,

Tokyo, 152-8552, Japan

SOURCE: Applied Physics Letters (2003), 83(23), 4755-4757

CODEN: APPLAB; ISSN: 0003-6951

PUBLISHER: American Institute of Physics

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 143433-45-8


RL: PRP (Properties)

(optical anisotropy in thermo-optic coeffs. of films formed on silicon substrates)

RN 143433-45-8 CAPLUS

CN

Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




REFERENCE COUNT:  
THIS


25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR

L56 ANSWER 10 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
AB Polymers incorporating the triptycene subunit were prepared for the  
mol.-level design of low dielec. constant (low- $\epsilon$ ) materials that can  
be used to manufacture faster integrated circuits. Triptycenes having  
restricted rotation by multiple point attachment to the polymer backbone  
are shown to introduce free volume into the films, thereby lowering their  
dielec. consts. The triptycene containing polymers exhibit a number of  
desirable  
properties including low-water absorption and high thermal stability.  
Systematic studies wherein comparisons are made between two sep. classes  
of triptycene polymers and their non-triptycene containing analogs  
demonstrate  
that proper insertion of triptycenes into a polymer backbone can give  
rise  
to a reduction in the material's dielec. constant while also improving  
its mech.  
properties. These characteristics are desired by the semiconductor  
industry for the next generation of microprocessors and memory to provide  
insulation of the increasingly shrinking features.  
ACCESSION NUMBER: 2003:827002 CAPIUS  
DOCUMENT NUMBER: 140:28127  
TITLE: Molecular Design of Free Volume as a Route to  
Low- $\epsilon$  Dielectric Materials  
AUTHOR(S): Long, Timothy M.; Swager, Timothy M.  
CORPORATE SOURCE: Department of Chemistry, Massachusetts Institute of  
Technology, Cambridge, MA, 02139, USA  
SOURCE: Journal of the American Chemical Society (2003),  
125(46), 14113-14119  
CODEN: JACSAT; ISSN: 0002-7863  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 632621-45-5P 632621-47-7P  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(synthesis, free volume, dielec., thermal and surface properties of  
triptycene-based low dielec. constant materials)  
RN 632621-45-5 CAPIUS  
CN Poly[oxy(6,14-bis(1,1-dimethylethyl)-9,10-dihydro-9,10[1',2']-  
benzoanthracene-1,4-diyl)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-  
biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




RN 632621-47-7 CAPIUS  
CN Poly[oxy(5,7,12,14-tetrahydro-5,14[1',2']-7,12[1'',2'']-dibenzenopentacene-  
6,13-diyl)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)]  
(9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 2-A



REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR  
THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT




AB The title phthalic acid derivs. (Markush structure given) are prepared by reaction of tetrafluorophthalic acid derivs. with MAM (A = divalent organic moiety; M = H, et.). Thus, I was prepared I is a raw material for the manufacture of fluorinated polyimides. The title phthalic acid derivs. are

useful as intermediates for optical materials, liquid crystals, etc.  
ACCESSION NUMBER: 2003:823334 CAPLUS  
DOCUMENT NUMBER: 139:330642  
TITLE: Phthalic acid derivatives as intermediates for halogenated polyimides, liquid crystals, etc., and process for manufacturing them  
INVENTOR(S): Okumura, Yasunori; Kuwahara, Masayoshi; Masuda, Takeshi  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JX00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003300934	A2	20031021	JP 2002-108095	20020410

PRIORITY APPLM. INFO.: JP 2002-108095 20020410

OTHER SOURCE(S): MARPAT 139:330642  
IT 615263-86-0P  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(phthalic acid derivs. as intermediates for halogenated polyimides, optical materials, liquid crystals, and process for manufacturing them)  
RN 615263-86-0 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(3,5,6-trifluoro-, tetraethyl ester (9CI) (CA INDEX NAME)



AB The invention refers to an optical waveguide and wavelength modulator comprising a fluoro-polyimide I (R1,2 = halo- or fluoroalkyl-substituted Ph groups linked by -O-, -CO-, SO2-, -S-, -Rf-, -ORf-, -C(Rf)2-, -RfO-, -(ORf)n-, -(RfO)n-, -(ORf)nO-, OC(O)-RfC(O)-; Rf = halo or fluoroalkyl; n = 1 - 10) which does not contain any C-H bonds.



ACCESSION NUMBER: 2003:582992 CAPLUS  
DOCUMENT NUMBER: 139:140706  
TITLE: Polyimide optical waveguide and polyimide filter-type wavelength combination/division device  
INVENTOR(S): Kobayashi, Junya; Kawakami, Naomi; Yamada, Noriyoshi; Kagei, Emiko; Yamamoto, Fumio; Kudo, Ayako; Matsura, Toru  
PATENT ASSIGNEE(S): NTT Advanced Technology Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
CODEN: JX00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003215364	A2	20030730	JP 2002-14719	20020123


PRIORITY APPLM. INFO.: JP 2002-14719 20020123

IT 143376-21-0 143376-23-2 143433-45-0  
143433-47-0 484016-84-4 484016-86-6  
RL: DEV (Device component use); USES (Uses)  
(polyimide optical waveguide and polyimide filter-type wavelength combination/division device)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
CMF C22 F10 O8

CM 2

CRN 1198-63-6  
CMF C6 H4 F4 N2


RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
CMF C22 F10 O8

CM 2

CRN 20115-19-9





RN 143433-45-8 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)


PAGE 1-A



RN 484016-86-6 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(5-chloro-2,4,6-trifluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




PAGE 1-B



RN 484016-84-4 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 5-chloro-2,4,6-trifluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1  
 CRN 474805-39-5  
 CHF C6 H4 Cl F3 N2



CH 2  
 CRN 143363-91-1  
 CHF C22 F10 O8

L56 ANSWER 13 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A polyimide excelling in heat resistance, chemical resistance, water repellency, dielec. characteristics, elec. characteristics, and optical characteristics and a polyamic acid useful as the raw material thereof are provided. The polyimides are prepared from polyamic acids containing chlorine atoms and fluorine atoms. A polyamic acid was prepared from 1,3-diamino-2,4,5,6-tetrafluorobenzene, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluorophthalic anhydride], and 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluorophthalic anhydride].

ACCESSION NUMBER: 2003:40181 CAPLUS  
 DOCUMENT NUMBER: 138:90676  
 TITLE: Fluorine-containing polyamic acid and polyimide, and optical material  
 INVENTOR(S): Tajiri, Kozo; Kuwabara, Masayoshi; Okumura, Yasunori; Matsui, Tohru; Yamada, Noriyoshi  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan; NTT Advanced Technology Corp.  
 SOURCE: Eur. Pat. Appl., 44 pp.  
 CODEN: EPXNDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1275679	A1	20030115	EP 2002-254908	20020712
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2003026799	A2	20030129	JP 2001-213743	20010713
JP 2003026800	A2	20030129	JP 2001-213744	20010713
US 2003050407	A1	20030313	US 2002-194791	20020711
CN 1397583	A	20030219	CN 2002-126201	20020715
PRIORITY APPL. INFO.:			JP 2001-213743	A 20010713
			JP 2001-213744	A 20010713

IT 484016-83-3P 484016-84-4P 484016-86-6P  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (fluorine-containing polyamic acid and polyimide, and optical material)


RN 484016-83-3 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine and 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

CH 1  
 CRN 474805-33-9  
 CHF C22 Cl4 F6 O8






CM 2

CRN 143363-91-1  
CMF C22 F10 O8

CM 3


CRN 1198-63-6  
CMF C6 H4 F4 N2

RN 484016-84-4 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 5-chloro-2,4,6-trifluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 474805-39-5

GI



AB The compds. are prepared by hydrolysis of aromatic tetranitriles I (X, X' = halo; n, n' = 1-3; Z = single bond, O, S, O-p-C6H4-mYmO, etc; Y = halo; m = 1-4) in the presence of acids in organic solvents. 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene was treated with H2SO4 in propionic acid under reflux for 6 h to give 95.7% 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene.


ACCESSION NUMBER: 2002:886125 CAPLUS  
DOCUMENT NUMBER: 137:384650  
TITLE: Preparation of halogen-containing aromatic carboxylic acids  
INVENTOR(S): Kuwahara, Masayoshi; Yokoo, Junko; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JJOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002332253	A2	20021122	JP 2001-142027	20010511
JP 3490407	B2	20040126		


PRIORITY APPL. INFO.: JP 2001-142027 20010511

OTHER SOURCE(S): MARPAT 137:384650  
IT 143363-92-2P 474805-31-7P  
RL: IMP (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)


CMF C6 H4 Cl F3 N2



CM 2


CRN 143363-91-1  
CMF C22 F10 O8

RN 484016-86-6 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(5-chloro-2,4,6-trifluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE


FORMAT



RN 474805-31-7 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




IT 143376-50-5, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 474805-29-3P, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrachlorobenzene  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)

L56 ANSWER 14 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 RN 474805-29-3 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




L56 ANSWER 15 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The present invention relates to halogen-containing aromatic compds. and Methods thereof. The present invention relates a halogen-containing aromatic acid dianhydride, halogen-containing aromatic tetranitrile compound, halogen-containing m-phenylenediamine compound and fluorine compound, and a method thereof. Tetrafluorophthalonitrile was reacted with tetrachlorohydroquinone in the presence of potassium fluoride and acetonitrile to give 1,4-bis(3,4-dicyanotrifluorophenoxy)tetrachlorobenzene.  
 ACCESSION NUMBER: 2002:866679 CAPIUS  
 DOCUMENT NUMBER: 137:354701  
 TITLE: Halogen-containing aromatic compound  
 INVENTOR(S): Kuvabara, Masayoshi; Okumura, Yasunori  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan; Nippon Catalytic Chem. Ind.  
 SOURCE: Eur. Pat. Appl., 38 pp.  
 CODEN: EPOIDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1256564	A1	20021113	EP 2002-253088	20020501
EP 1256564	B1	20040728		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2002332268	A2	20021122	JP 2001-142028	20010511
JP 2002332281	A2	20021122	JP 2001-142029	20010511
JP 2002332264	A2	20021122	JP 2001-142031	20010511
JP 3563040	B2	20040908		
JP 2002332254	A2	20021122	JP 2001-142032	20010511
US 2003018204	A1	20030123	US 2002-133158	20020426
EP 1462436	A1	20040929	EP 2004-10371	20020501
R: DE, FR, GB				
CN 1385427	A	20021218	CN 2002-119176	20020513
PRIORITY APPLN. INFO.:				
			JP 2001-142028	A 20010511
			JP 2001-142029	A 20010511
			JP 2001-142031	A 20010511
			JP 2001-142032	A 20010511
			EP 2002-253088	A3 20020501

OTHER SOURCE(S): MARPAT 137:354701  
 IT 474805-33-9P  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (halogen-containing aromatic compound)  
 RN 474805-33-9 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)


L56 ANSWER 15 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)



IT 474805-29-3P 474805-31-7P  
 RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation);  
 RACT (Reactant or reagent)  
 (halogen-containing aromatic compound)  
 RN 474805-29-3 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



RN 474805-31-7 CAPIUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L56 ANSWER 16 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The devices comprise: (1) a substrate/cladding comprising a 1st fluorinated polyimide ( $n = n_1$ ); (2) a core comprising a 2nd fluorinated polyimide ( $n = n_2 > n_1$ ); and (3) an upper cladding layer comprising a 1st fluorinated polyimide ( $n = n_1$ ), where the laminate (1)-(3) are heat-compressed.  
 ACCESSION NUMBER: 2002:750899 CAPIUS  
 DOCUMENT NUMBER: 137:286156  
 TITLE: Optical waveguide equipped substrate devices and manufacture  
 INVENTOR(S): Matsuura, Toru; Yamamoto, Fumio; Yokoyama, Hirokazu; Sugimoto, Noriki; Moriyama, Hideki  
 PATENT ASSIGNEE(S): NTT Advanced Technology Corp., Japan; Du Pont-Toray Co., Ltd.  
 SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
 CODEN: JYQXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002286957	A2	20021003	JP 2001-91856	20010328
PRIORITY APPLN. INFO.:			JP 2001-91856	20010328


IT 143433-45-8  
 RL: DEV (Device component use); USES (Uses)  
 (optical waveguide equipped substrate devices and manufacture)  
 RN 143433-45-8 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




L56 ANSWER 17 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The refractive indexes of eleven kinds of polyimide (PI) films formed on silicon substrates have been measured at three wavelengths (0.633, 1.320, and 1.523  $\mu\text{m}$ ) in the transverse elec. (TE) and transverse magnetic (TM) polarization modes using the prism-coupling method. The wavelength dependence of average refractive indexes was fitted using the simplified Cauchy's formula, and the estimated refractive indexes at infinite wavelength (ninf) and the coeffs. of wavelength dispersion (D) were determined. The PIs having higher ninf exhibit larger D, and the values of D are linearly proportional to ninf for aromatic PIs, while those of wholly alicyclic PIs are very small and deviate from the linearity. In a similar manner, the in-plane/out-of-plane birefringence ( $\Delta n$ ) values of the aromatic PIs having higher refractive indexes are large and show significant wavelength dependence, while the values of  $\Delta n$  of alicyclic PIs are negligible. Approx. equations are presented for aromatic PIs to estimate ninf and refractive indexes at 1.320 and 1.523  $\mu\text{m}$  from a refractive index at 0.633  $\mu\text{m}$ .  
 ACCESSION NUMBER: 2002:638977 CAPIUS  
 DOCUMENT NUMBER: 137:353613  
 TITLE: Wavelength dependence of refractive indices of polyimides in visible and near-IR regions  
 AUTHOR(S): Ando, Shinji; Watanabe, Yasufumi; Matsuura, Tohru  
 CORPORATE SOURCE: Department of Organic and Polymeric Materials, Tokyo Institute of Technology, Tokyo, 152-8552, Japan  
 SOURCE: Japanese Journal of Applied Physics, Part 1: Regular Papers, Short Notes & Review Papers (2002), 41(8), 5254-5258  
 CODEN: JAPNDE  
 PUBLISHER: Japan Society of Applied Physics  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143376-21-0 143433-45-8 148362-06-5  
 148446-36-0  
 RL: PRP (Properties)  
 (effect of refractive indexes of polyimides-(polyether-fluoropolymers) on wavelength in visible and near-IR regions)  
 RN 143376-21-0 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CMF C22 F10 O8


L56 ANSWER 17 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)



CH 2  
 CRN 1198-63-6  
 CMF C6 H4 F4 N2




RN 143433-45-8 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 148362-06-5 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-


L56 ANSWER 17 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




PAGE 1-A

PAGE 1-B


RN 148446-36-0 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diamine (9CI) (CA INDEX NAME)  
 CH 1  
 CRN 143363-91-1  
 CMF C22 F10 O8



CH 2  
CRN 341-58-2  
CMF C14 H10 F6 N2



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS  
FORMAT RECORD. ALL CITATIONS AVAILABLE IN THE RE



AB The magnetic components, useful for elec. transformers, DC converters, etc., have insulation layers of polyimides I (R1 = tetravalent aromatic group; R2 = divalent aromatic group; R1 and/or R2 have no C-H bond).


ACCESSION NUMBER: 2002:139128 CAPIUS  
DOCUMENT NUMBER: 136:209529  
TITLE: Heat-resistant thin-film magnetic components with low stray capacitance having polyimide insulation layers with low dielectric constant  
INVENTOR(S): Tajiri, Kozo; Konishi, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 17 pp.  
CODEN: JI00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002057032	A2	20020222	JP 2000-243310	20000810
PRIORITY APPLN. INFO.:			JP 2000-243310	20000810


IT 143376-21-0 143433-45-8  
RL: TEM (Technical or engineered material use); USES (Uses) (heat-resistant thin-film magnetic components with low stray capacitance having polyimide insulation layers with low dielec. constant)

RN 143376-21-0 CAPIUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CMF C22 F10 O8



CH 2  
CRN 1198-63-6  
CMF C6 H4 F4 N2



RN 143433-45-8 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



L56 ANSWER 19 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
AB The repeating units of polyimides for coating materials contain tetravalent aromatic organic groups and aromatic groups having no CH bonds. Thus,


a film of 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer had dielec. constant 2.6 at 1 MHz, compared with 3.5 for a com. polyimide.

ACCESSION NUMBER: 2002:139034 CAPIUS  
DOCUMENT NUMBER: 136:185465  
TITLE: Insulating coating materials for coil wire  
INVENTOR(S): Tajiri, Kozo; Kuwahara, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 16 pp.  
CODEN: JI00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002056720	A2	20020222	JP 2000-243309	20000810
PRIORITY APPLN. INFO.:			JP 2000-243309	20000810

IT 143433-45-8, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer, SRU  
148273-08-9, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer  
RL: PRP (Properties); TEM (Technical or engineered material use); USES (Uses) (fluoro polyimide insulating coating materials for coil wire)


RN 143433-45-8 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 148273-08-9 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

L56 ANSWER 19 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CM 1



CRN 143363-92-2  
CMF C22 H4 F10 O10



CM 2  
CRN 1198-63-6  
CMF C6 H4 F4 N2



L56 ANSWER 20 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
GI



II

AB The agent is a polyimide of I (R1 = tetravalent aromatic organic group; R2 = divalent aromatic organic group; R1 and/or R2 = aromatic organic group without C-H bonding). Thus, a coating was made from II in dimethylacetamide.

ACCESSION NUMBER: 2002:129271 CAPLUS  
DOCUMENT NUMBER: 136:169089

TITLE: Surface coating agent for printed circuit board with relatively low dielectricity, good heat resistance, and good moisture resistance  
INVENTOR(S): Tajiri, Kozo; Kuwahara, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 17 pp.  
CODEN: JJOQAF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002053802	A2	20020219	JP 2000-243308	20000810

PRIORITY APPL. INFO.: JP 2000-243308 20000810

IT 143433-45-8  
RL: PRP (Properties); TEM (Technical or engineered material use); USES (Uses)  
(surface coating agent for printed circuit board with relatively low dielectricity, good heat resistance, and good moisture resistance)  
RN 143433-45-8 CAPLUS

L56 ANSWER 20 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CM

Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



L56 ANSWER 21 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB The title circuit boards comprise substrate materials, circuit layers, adhesive layers, and surface-protection films. The substrate, adhesive, and/or surface-protection insulator materials are made from a polyimide (I: R1 = tetravalent aromatic organic group; R2 = aromatic organic group not containing C-H bonding). The polyimides gives the circuit boards improved size and dielec. stability and suitable for high-frequency flexible printed circuit boards.

ACCESSION NUMBER: 2001:796858 CAPLUS  
DOCUMENT NUMBER: 135:351388

TITLE: High-frequency multilayer circuit boards  
INVENTOR(S): Tajiri, Kozo; Konishi, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
CODEN: JJOQAF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001308542	A2	20011102	JP 2000-119916	20000420

PRIORITY APPL. INFO.: JP 2000-119916 20000420

IT 371976-28-2P 371976-29-3P  
RL: DEV (Device component use); PNU (Preparation, unclassified); PRP (Properties); PREP (Preparation); USES (Uses)  
(insulator film; high-frequency multilayer circuit boards)

RN 371976-28-2 CAPLUS


CM Poly[(4,4',6,6',7,7'-hexafluoro-1,1',3,3'-tetrahydro-1,1',3,3'-tetraoxo[2,2'-bi-2H-isoindole]-5,5'-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 371976-29-3 CAPLUS  
CM 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with hydrazine (9CI)  
(CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 O8



CM 2

CRN 302-01-2  
CHF H4 N2H<sub>2</sub>N-NH<sub>2</sub>

AB The title board comprises a dielec. film from a polyimide having a repeating unit represented by (I), where R1 = tetravalent arm. organic group, R2 = divalent arm. organic group, and R1 and/or R2 being free of a C-H bond.


The board has a superior high-frequency transmission property.

ACCESSION NUMBER: 2001:796833 CAPLUS  
DOCUMENT NUMBER: 135:351372  
TITLE: Wiring board for high-frequency application  
INVENTOR(S): Tajiri, Kozor Konishi, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 15 pp.  
CODEN: JKKKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001308474	A2	20011102	JP 2000-119917	20000420


PRIORITY APPLN. INFO.: JP 2000-119917 20000420

IT 371976-28-2 371976-29-3  
RL: DEV (Device component use); USES (Uses)  
(polyimide dielec. film for wiring board for high-frequency application)  
RN 371976-28-2 CAPLUS  
CN Poly[(4,4',6,6',7,7'-hexafluoro-1,1',3,3'-tetrahydro-1,1',3,3'-tetraoxo[2,2'-bi-isoindole]-5,5'-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 371976-29-3 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with hydrazine (9CI)  
(CA

CM 1

CRN 143363-91-1  
CHF C22 F10 O8


CM 2

CRN 302-01-2  
CHF H4 N2H<sub>2</sub>N-NH<sub>2</sub>

AB Two kinds of new aromatic poly(arylene ether)s containing sulfonic acid groups were synthesized. Polymer 1 composed of tetraphenylphenylene ether and perfluorobiphenylene units was sulfonated with chlorosulfonic acid. Sulfonation took place only at the para position of the pendant Ph rings. The average degree of sulfonation per repeating unit (m) was controlled from 1 to 4. Sulfonated polymer 2 with m = 3 was soluble in methanol and DMSO and swelled in water. Incorporating bis(3,5-dimethylphenyl)sulfone moieties into the sulfonated polymer imparts less methanol affinity. Polymers 4 with 30-65 mol % tetrakis(sulfonylphenyl)phenylene ether units has high decomposition temps. above 300°C, hydrophilicity, and good hydrolytic stability.

ACCESSION NUMBER: 2001:670068 CAPLUS  
DOCUMENT NUMBER: 135:372091  
TITLE: Synthesis and properties of poly(arylene ether)s bearing sulfonic acid groups on pendant phenyl rings  
AUTHOR(S): Miyatake, Kenji; Hay, Allan S.  
CORPORATE SOURCE: Department of Chemistry, McGill University, Montreal, QC, H3A 2K6, Can.  
SOURCE: Journal of Polymer Science, Part A: Polymer Chemistry (2001), 39(19), 3211-3217  
CODEN: JPACED; ISSN: 0887-624X  
PUBLISHER: John Wiley & Sons, Inc.  
DOCUMENT TYPE: Journal  
LANGUAGE: English

IT 338453-92-ZDP, para-phenyl-sulfonated  
RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(1-4 sulfonate groups per repeating unit; synthesis and properties of poly(arylene ether)s bearing sulfonic acid groups on pendant Ph rings)  
RN 338453-92-2 CAPLUS  
CN Poly[oxy(4',5'-diphenyl[1,1':2',1''-terphenyl]-3',6'-diyl)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI)  
(CA INDEX NAME)



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 24 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The compds. QCO-p-C6H4O-p-C6H4R (Q = 2,3,4,5,6-pentafluorophenyl; R = OH, COO) are manufactured. Thus, 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone (prepared from 2,3,4,5,6-pentafluoro-4'-methoxybenzophenone) was polymerized in dimethylacetamide at 160° to give a polymer showing good solubility in dimethylacetamide and m-cresol, 10% weight loss (in air) temperature 421°.


ACCESSION NUMBER: 2001:176773 CAPIUS  
 DOCUMENT NUMBER: 134:208664  
 TITLE: (2,3,4,5,6-pentafluorobenzoyl)diphenyl ethers and fluorine-containing aryl ether ketone polymers having high solubility and heat stability  
 INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Casiday, Rachel E.; Fisch, John W., III; Reddy, V. Sreenivasulu  
 PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
 CODEN: JQOQAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001064226	A2	20010313	JP 1999-183950	19990629
PRIORITY APPL. INFO.:			US 1998-106270	A 19980629
			JP 1999-180091	A 19990625

IT 213693-15-3P  
 RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
 (pentafluorobenzoyl)phenyl ether as monomers for F-containing aryl ether ketone polymers having high solubility and heat stability)

RN 213693-15-3 CAPIUS  
 CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene) (9CI) (CA INDEX NAME)

PAGE 1-A




L56 ANSWER 25 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The polymerization of 2,3,5,6-tetrafluorophenylhydroquinone (or 2,2',3,3',5,5'-hexafluoro-4,4'-dihydroxybiphenyl) with  $\alpha,\omega$ -tetrahydroperfluoroalkenediol and decafluorobiphenyl was carried out to synthesize a series of copolymers III (Mw = 49 100-80 900). The copolymers III are composed of arylene ether (10-30 mol %) and fluorinated alkane (90-70 mol %) moieties. The reaction of III with chlorosulfonic acid gave sulfonated polymers IV, which are soluble in polar organic solvents and form flexible and transparent films by casting from solution. The polymers IV have glass transition temps. of 109-155° and decomposition temps. of ca. 300°. The hydrated polymers show protonic conductivity (3.4  $\times 10^{-3}$  S cm $^{-1}$ ), which does not decrease at temps. up to 170 °C.

ACCESSION NUMBER: 2001:153020 CAPIUS  
 DOCUMENT NUMBER: 134:340775  
 TITLE: Synthesis and Properties of Novel Sulfonated Arylene Ether/fluorinated Alkane Copolymers  
 AUTHOR(S): Miyatake, Kenji; Oyaizu, Kenichi; Tsuchida, Eishun; Hay, Allan S.  
 CORPORATE SOURCE: Department of Chemistry, McGill University, Montreal, QC, H3A 2K6, Can.  
 SOURCE: Macromolecules (2001), 34(7), 2065-2071  
 CODEN: MAMOBX; ISSN: 0024-9297  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

IT 338453-92-2P  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (synthesis and properties of sulfonated arylene ether/fluorinated alkane copolymers)


RN 338453-92-2 CAPIUS  
 CN Poly[oxy(4',5'-diphenyl[1,1':2',1''-terphenyl]-3',6'-diyl)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)



REFERENCE COUNT: 20 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L56 ANSWER 24 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-B



L56 ANSWER 26 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The compns., useful for elec. insulating materials, contain (p-C6F4COC6H4-qXqORlm)n [R1 = C6H4-rX'r(CO-p-C6F4OR2)pO; R2 = divalent aryl; X, X' = halo, lower alkyl, alkoxy; q, r = 0-4; m, p = 0, 1]. Thus, 2,2-bis(4-hydroxyphenyl)-1,1,1,3,3,3-hexafluoropropane was polymerized with 4,4'-bis(2,3,4,5,6-pentafluorobenzoyl)diphenyl ether to give a polymer showing dielec. constant 3.06 at 25°, 10% weight loss temperature 524° under N<sub>2</sub> and Tg 174°.

ACCESSION NUMBER: 2001:124290 CAPIUS  
 DOCUMENT NUMBER: 134:179345  
 TITLE: Low dielectric fluorinated aromatic polyether ketone compositions with good heat resistance  
 INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Okumura, Yasunori  
 PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 21 pp.  
 CODEN: JQOQAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001049110	A2	20010220	JP 1999-226981	19990810
JP 3539897	B2	20040707	JP 1999-226981	19990810

PRIORITY APPL. INFO.:


IT 213693-15-3P  
 RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
 (low dielec. fluorinated aromatic polyether ketone compns. with good heat resistance)

RN 213693-15-3 CAPIUS  
 CN Poly[oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



L56 ANSWER 27 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN


AB Ketonic di-Ph ethers of p-R1OC6H4OC6H4R2-p type compds. (R1 = 2,3,4,5,6-pentafluorobenzoyl; R2 = OH, pentafluorobenzoyl group) and polyether-polyketone polymers containing tetrafluorophenylene and phenylene groups are provided which have good mech. strength, toughness, elec. property, thermal oxidative stability and solubility. Thus, heating 0.5 g 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone with 0.36 ground K carbonate, 2 mL N-methyl-2-pyrrolidone and 1 mL PhMe at 160° while stirring for 3 h gave a polymer at 85% yield and having viscosity 0.5 g/dL in AcHMe2.

ACCESSION NUMBER: 2001:25791 CAPLUS  
DOCUMENT NUMBER: 134:86663  
TITLE: (2,3,4,5,6-Pentafluorobenzoyl)diphenyl ether compound,  
and fluorine-containing aryl ether ketone polymer  
INVENTOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.;  
Fitch, John W., III; Reddy, V. Sreenivasulu  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: U.S., 22 pp., Cont.-in-part of U.S. Ser. No. 106,270, abandoned.  
CODEN: USXXAM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6172181 B1		20010109	US 1999-354976	19990716
OTHER SOURCE(S):		MARPAT 134:86663	US 1998-106270	19980629

IT 213693-15-3P  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation) (manufacture of (2,3,4,5,6-pentafluorobenzoyl)diphenyl ether compound and fluorine-containing aryl ether ketone polymer)  
RN 213693-15-3 CAPLUS  
CN Poly[oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene]carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L56 ANSWER 28 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN


AB An electrolyte is provided having a backbone that includes a plurality of aromatic constituents coupled together by at least one atom having a  $\pi$ -cloud, and in which a halogen atom and an ion exchange group are covalently bound directly to the backbone. Furthermore, the electrolyte is high temperature resistant and may comprise perhalogenated polymers, including perhalogenated polyphenylenes, perhalogenated polyamides, perhalogenated aromatic polyesters, perhalogenated polyimides, etc. Still further, the electrolyte may have acidic groups as ion exchange groups, including sulfonic acid groups, or phosphoric acid groups. A typical polymer electrolyte was manufactured by solid-liquid phase transfer polymerization of 2,3,5,6-tetrakis(benzylthio)-1,4-hydroquinone 18 h at 80° with bis(pentafluorophenyl) sulfone in the presence of K2CO3 and 18-crown-6 in AcHMe2, reaction of resulting polymer 15 h with Cl in CH2Cl2 containing HOAc and water, and conversion of the resulting sulfonyl chloride group-containing polymer to a sulfonic acid group-containing polymer by reaction with aqueous NaOH in DME.

ACCESSION NUMBER: 2001:12796 CAPLUS  
DOCUMENT NUMBER: 134:72076  
TITLE: High-temperature polymer electrolytes  
INVENTOR(S): Narang, Subhash; Ventura, Susanna  
PATENT ASSIGNEE(S): Sri International, USA  
SOURCE: PCT Int. Appl., 30 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001001510	A1	20010104	WO 2000-US40282	20000621
W: JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 6248480	B1	20010619	US 1999-340778	19990628
PRIORITY APPLN. INFO.:			US 1999-340778	A 19990628
			US 1998-91051P	P 19980629
			US 1998-109154P	P 19981120

IT 316149-04-9P  
RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses) (high-temperature polymer electrolytes)  
RN 316149-04-9 CAPLUS  
CN Poly[oxy(2,3,5,6-tetrakis((phenylmethyl)thio)-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)sulfonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)





REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

AB High temperature polybenzazole and polyether polymer electrolytes are provided.

High temperature polybenzazole polymer electrolytes may comprise a benzobisoxazole, a benzobisthiazole, a benzobisimidazole, a difluorodisulfonated Ph ring or a sulfonated bisphenylether. High temperature

polyether polymers comprise a persulfonated Ph ring, and a substituted Ph ring or a substituted bisphenylsulfonate ring system.

ACCESSION NUMBER: 2001:12792 CAPLUS  
DOCUMENT NUMBER: 134:86656


TITLE: High temperature polybenzazole and polyether electrolytes  
INVENTOR(S): Varang, Subhash; Ventura, Susanna; Koolpe, Gary  
PATENT ASSIGNEE(S): Sri International, USA  
SOURCE: PCT Int. Appl., 17 pp.

DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001001506	A1	20010104	WO 2000-US40278	20000621
W: JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 6176984	B1	20010123	US 1999-340777	19990628
EP 1243043	A1	20020925	EP 2000-957985	20000621
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
JP 2003503599	T2	20030128	JP 2001-506630	20000621
PRIORITY APPLN. INFO.:			US 1999-340777	A 19990628
			US 1998-91051P	P 19980629
			US 1998-109154P	P 19981120
			WO 2000-US40278	W 20000621

IT 316149-04-9P  
RL: IMF (Industrial manufacture); POF (Polymer in formulation); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses) (high temperature polybenzazole and polyether electrolytes)

RN 316149-04-9 CAPLUS  
CN Poly[oxy(2,3,5,6-tetrakis[(phenylmethyl)thio]-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)sulfonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

AB Following a review (50 refs.) on optical fibers and the preparation and characterization of perfluorinated diamines and dianhydrides new results are presented with respect to synthesis and properties of the corresponding perfluorinated polyimides. Novel perfluorinated polyimides were synthesized from newly prepared 1,4-bis(3,4-dicarboxyltrifluorophenoxy)tetrafluorobenzene dianhydride (10FEDA) and several perfluorinated aromatic diamines. To generate high mol. weight perfluorinated poly(amic acid)s, the reactivities of 5 diamines and their monoacyl deriva. were estimated before from end group contents (19F NMR)

of the corresponding poly(amic acid), 15N and 1H NMR shifts, and calculated ionization potentials. The perfluorinated polyimides synthesized from 10FEDA and 3 (out of the 5) selected diamines gave flexible films with glass transition temps. values >270° and with high transparencies at the wavelengths of optical communications (1.0-1.7 μm) with sufficient mech. strength. Their dielec. consts. and refractive indexes are as low as those of conventional fluorinated polyimides, and their in-plane/out-of-plane birefringence is low. These characteristics indicate that perfluorinated polyimides are promising materials for optical communication applications.

ACCESSION NUMBER: 2000:292601 CAPLUS  
DOCUMENT NUMBER: 133:193573  
TITLE: Synthesis and properties of perfluorinated polyimides  
AUTHOR(S): Ando, Shinji; Matsuura, Tohru; Sasaki, Shigekuni  
CORPORATE SOURCE: Science and Core Technology Group, Nippon Telegraph and Telephone Corp., Musashino, 180, Japan  
SOURCE: Fluoropolymers (1999), Volume 2, 277-303. Editor(s): Hougham, Gareth. Kluwer Academic/Plenum Publishers: New York, N. Y.

DOCUMENT TYPE: Conference; General Review  
LANGUAGE: English

IT 143376-21-0P 143376-22-1P 143376-23-2P  
143376-24-3P 143433-45-8P 143433-46-9P  
143433-47-0P 143433-48-1P 289650-74-4P  
289650-76-6P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (synthesis and properties of perfluorinated polyimides)

RN 143376-21-0 CAPLUS


CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-, 1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 08




PAGE 1-B




RN 143433-48-1 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)thio(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 289650-74-4 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8




CM 2

CRN 1038-66-0  
 CMF C12 H4 F8 N2



RN 289650-76-6 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 53 THERE ARE 53 CITED REFERENCES AVAILABLE FOR THIS

L56 ANSWER 31 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN

AB The lattice is manufactured by irradiating a fluorinated polyimide optical waveguide with x-ray via a x-ray mask having a desirable diffraction lattice pattern to form a region with periodically changed refractive index and a diffraction lattice region with homogeneously changed refractive index. The lattice is manufactured by x-ray radiation in a simple process.

ACCESSION NUMBER: 1999:518975 CAPIUS  
DOCUMENT NUMBER: 131:191675  
TITLE: Formation method of fluorinated polyimide-based optical waveguide diffraction lattice  
INVENTOR(S): Tanemura, Yoshiaki; Ishii, Tetsuyoshi; Kobayashi, Junya; Maruno, Toru  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 11223740	A2	19990817	JP 1998-24152	19980205

PRIORITY APPLN. INFO.: JP 1998-24152 19980205



IT 143376-21-OP  
RL: DEV (Device component use); IMF (Industrial manufacture); PREP (Preparation); USES (Uses)  
(manufacture of fluorinated polyimide-based optical waveguide diffraction lattice)  
RN 143376-21-0 CAPIUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 08



L56 ANSWER 32 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN



AB The title compds. [I: A1-A6 = H, F, Me, OMe, CF3, OCF3, etc.; ≥1 of A1-A6 contains F or F-containing group; T = O, S; Z = (un)substituted (hetero)arylene; m = 0, 1], useful, e.g., for the manufacture of polybenzoxazoles and other heat-resistant polymers, were prepared by (a) condensation of nitro compds. with alkali hydroxides or alkali hydrogen sulfides, or with dihydroxy- or dimercapto compds. HTZTH (T, Z as above) in the presence of a base, and (b) reduction of the intermediate bis-o-nitro(thio)phenols to their amino analogs. For example, adding 60

g K2CO3 in portions to a solution of 33.6 g 4-HOC6H4C(CF3)2C6H4OH-4' and 42.6 g C6F5OH in 400 mL DMSO, stirring the mixture for 24 h at ambient temperature, heating for 6 h at 80°, adding 10 g KHCO3 and heating the whole for addnl. 18 h gave 91% 2,2-bis[4-(4-nitro-3-hydroxy-2,5,6-trifluorophenoxy)phenyl]hexafluoropropane. Reduction of the latter in

1:1 EtOAc/THF at 1 bar H in the presence of Pd/C gave 93 % of the title compound II.


ACCESSION NUMBER: 1999:236983 CAPIUS  
DOCUMENT NUMBER: 130:267209  
TITLE: Preparation of bis-o-amino(thio)phenols  
INVENTOR(S): Sezi, Recai; Keitmann, Michael  
PATENT ASSIGNEE(S): Siemens Aktiengesellschaft, Germany  
SOURCE: Eur. Pat. Appl., 16 pp.  
CODEN: EPOXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 906903	A2	19990407	EP 1998-117341	19980912
EP 906903	A3	19990414		

L56 ANSWER 31 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

CM 2


CRN 1198-63-6  
CMF C6 H4 F4 N2




L56 ANSWER 32 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)


EP 906903 B1 20011212  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO  
JP 11158129 A2 19990615 JP 1998-270392 19980924  
US 6150558 A 20001121 US 1998-161144 19980924  
PRIORITY APPLN. INFO.: DE 1997-19742195 A 19970924

OTHER SOURCE(S): MARPAT 130:267209  
IT 222294-75-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn and reduction; preparation of bis-o-amino(thio)phenols)  
RN 222294-75-9 CAPIUS  
CN Phenol, 3,3'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(2,4,5-trifluoro-6-nitro- (9CI) (CA INDEX NAME)



IT 222294-71-5P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of bis-o-amino(thio)phenols)  
RN 222294-71-5 CAPIUS  
CN Phenol, 3,3'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(6-amino-2,4,5-trifluoro- (9CI) (CA INDEX NAME)






AB The title precursors, having the partial structure I (A1-6 = H, F, Me, CF<sub>3</sub>, OMe, OCF<sub>3</sub>, Et, C<sub>2</sub>F<sub>5</sub>, OEt, OC<sub>2</sub>F<sub>5</sub>; T = O or S; Z = aromatic or heterocyclic group; m = 0 or 1), are soluble in organic solvents or in aqueous alkalis free from metal ions, suitable for use in photosensitive compns., and can be spin-coated. Heating benzyl 5-hydroxy-2-nitrophenyl ether 0.1, C<sub>6</sub>F<sub>5</sub>C<sub>6</sub>F<sub>5</sub> 0.05, and K<sub>2</sub>CO<sub>3</sub> 0.22 mol in DMSO at 100° for 4 h gave 4,4'-bis[3-(benzyloxy)-4-nitro]octafluorobiphenyl, hydrogenation of which over Pd/C in THF-EtOAc gave 4,4'-bis(4-amino-3-hydroxyphenoxy)octafluorobiphenyl (II). Polymerization of 0.1 mol II with 0.1 mol 4,4'-oxydibenzoyl chloride in butyrolactone containing pyridine at 10° to room temperature gave a polymer (soluble in a variety of organic solvents an an aqueous alkaline developer), cyclization of which at 350° for 1 h gave a polybenzoxazole with moisture uptake (24 h, 76% relative humidity) 0.84% and weight loss in TGA 1% at 480°.

ACCESSION NUMBER: 1999:219825 CAPLUS  
DOCUMENT NUMBER: 130:282476  
TITLE: Precursors for polybenzoxazoles and polybenzothiazoles  
INVENTOR(S): Sezi, Recai; Schmid, Gunter; Keilmann, Michael  
PATENT ASSIGNEE(S): Siemens Aktiengesellschaft, Germany  
SOURCE: Eur. Pat. Appl., 14 pp.  
CODEN: EPOQDW  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 905169	A2	19990331	EP 1998-117333	19980912
EP 905169	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
JP 11171994	A2	19990629	JP 1998-270388	19980924
US 6153350	A	20001128	US 1998-161148	19980925
PRIORITY APPLN. INFO.:		DE 1997-19742132	A	19970924

IT 222612-40-0P 222612-41-1P  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(precursors for polybenzoxazoles and polybenzothiazoles)  
RN 222612-40-0 CAPLUS  
CN Hexanedioyl dichloride, polymer with 3,3'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[6-amino-2,4,5-trifluorophenol] (9CI) (CA INDEX NAME)

CH 1  
CRN 222294-71-5  
CMF C18 H6 F10 N2 O4




CH 2  
CRN 111-50-2  
CMF C6 H8 Cl2 O2



RN 222612-41-1 CAPLUS  
CN Poly(oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,6-trifluoro-5-hydroxy-1,4-phenylene)imino(1,6-dioxo-1,6-hexanediyl)imino(2,3,5-trifluoro-6-hydroxy-1,4-phenylene)) (9CI) (CA INDEX NAME)


PAGE 1-A



PAGE 1-B

]


IT 222294-71-5P  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(preparation of)  
RN 222294-71-5 CAPLUS  
CN Phenol, 3,3'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[6-amino-2,4,5-trifluoro- (9CI) (CA INDEX NAME)



IT 222612-40-0DP, cyclized 222612-41-1DP, cyclized  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(preparation of)  
RN 222612-40-0 CAPLUS  
CN Hexanedioyl dichloride, polymer with 3,3'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[6-amino-2,4,5-trifluorophenol] (9CI) (CA INDEX NAME)


CH 1  
CRN 222294-71-5  
CMF C18 H6 F10 N2 O4

CH 2  
CRN 111-50-2  
CMF C6 H8 Cl2 O2




RN 222612-41-1 CAPLUS  
CN Poly(oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,6-trifluoro-5-hydroxy-1,4-phenylene)imino(1,6-dioxo-1,6-hexanediyl)imino(2,3,5-trifluoro-6-hydroxy-1,4-phenylene)) (9CI) (CA INDEX NAME)

PAGE 1-A



] n



AB The films, useful for printed circuit boards and optical waveguides, comprise polyimides with repeating unit I (R1 = tetravalent organic groups from Q1-3 (Rf1, Rf2 = H, F, or CF3; X = C(CF3)2, CO, O, SO2, single bond); R2 = F-containing aryls). The films are prepared by polymerization of corresponding tetracarboxylic dianhydride with R2(NH2)2 (R2 = the same as above) in organic solvents, imidation, and removal of solvents. Thus, pyromellitic dianhydride was polymerized with equimolar 2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl at room temperature in N,N-dimethylacetamide to give a polyamic acid solution, which was applied on an Al substrate and cured at 70-350° and immersed in 10% HCl to give a water-repellent polyimide film showing thermal decomposition temperature 610° and water contact angle 74.6°.

ACCESSION NUMBER: 1999:65363 CAPLUS  
DOCUMENT NUMBER: 130:183528  
TITLE: Fluorine-containing polyimide films with excellent water repellency and heat resistance and their preparation  
INVENTOR(S): Matsuura, Toru; Sasaki, Shigekuni; Kobayashi, Shogo  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKKXAF  
DOCUMENT TYPE: Patent

LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 11021350	A2	19990126	JP 1997-187252	19970630

PRIORITY APPLN. INFO.: JP 1997-187252 19970630


IT 143376-21-OP 143376-23-2P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) ether copolymer 143376-24-3P 143433-45-OP 143433-47-OP, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) ether copolymer, sru 143433-48-1P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) sulfide copolymer, sru 148362-03-2P 148362-06-5P 148446-36-OP, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl copolymer 148446-40-6P 205451-28-1P 220592-84-7P  
RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(fluorine-containing polyimide films with excellent water repellency

and heat resistance for elec. or optical uses)

RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8



CM 2


CRN 1198-63-6  
CMF C6 H4 F4 N2



RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8



CM 2


CRN 20115-19-9  
CMF C12 H4 F8 N2 O




RN 143376-24-3 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-thiobis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

L56 ANSWER 34 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 CN 1


CRN 143363-91-1  
 CMF C22 F10 O8



CM 2  
 CRN 61907-46-8  
 CMF C12 H4 F8 N2 S




RN 143433-45-8 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPIUS

L56 ANSWER 34 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B




RN 148362-03-2 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]-1,4-phenylene] (9CI) (CA INDEX NAME)


L56 ANSWER 34 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 143433-48-1 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)thio(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

L56 ANSWER 34 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A




PAGE 1-B



RN 148362-06-5 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diyl]] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148446-36-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)-1,1'-biphenyl]-4,4'-diamine (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
 CMF C22 F10 O8



L56 ANSWER 34 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 phenyleneoxy)]bis(benzenamine) (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8




CM 2

CRN 69563-88-6  
 CMF C27 H20 F6 N2 O2




RN 220592-84-7 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)-1,4-phenyleneoxy-1,4-phenylene(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)-1,4-phenyleneoxy-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A



CM 2


CRN 341-58-2  
 CMF C14 H10 F6 N2



RN 148446-40-6 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis(benzenamine) (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2


CRN 1095-78-9  
 CMF C15 H12 F6 N2



RN 205451-28-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis(4,1-

L56 ANSWER 34 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-B





L56 ANSWER 35 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
AB The array comprises the cores and the claddings employing fluoropolyimides

and thermo-optical phase shifters.  
ACCESSION NUMBER: 1998:811477 CAPLUS  
DOCUMENT NUMBER: 130:102659  
TITLE: Wavelength modulating waveguide matrix array  
INVENTOR(S): Kobayashi, Junya; Maruno, Toru; Hirata, Yasuoki;  
Inoue, Yasuyuki; Sasaki, Shigekuni; Matsuura, Toru  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 11 pp.  
CODEN: JN00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10332959	A2	19981218	JP 1997-146806	19970604

PRIORITY APPLN. INFO.: JP 1997-146806 19970604

IT 143376-21-0, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine copolymer  
143433-45-8, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine copolymer, polyimide, sru  
219142-03-7, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine-2,2-bis(3,4-dicarboxyphenyl)hexafluoropropane dianhydride-4,4'-oxydianiline copolymer  
RL: DEV (Device component use); USES (Uses)  
(wavelength modulating waveguide matrix array)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
CHF C22 F10 O8



CH 2


CRN 1198-63-6

L56 ANSWER 35 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CHF C6 H4 F4 N2



RN 143433-45-8 CAPLUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 219142-03-7 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis(benzenamine), 2,4,5,6-tetrafluoro-1,3-benzenediamine and 5,5'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

CH 1

CRN 143363-91-1  
CHF C22 F10 O8

L56 ANSWER 35 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




CH 2

CRN 1198-63-6  
CHF C6 H4 F4 N2



CH 3


CRN 1107-00-2  
CHF C19 H6 F6 O6



CH 4


CRN 101-80-4  
CHF C12 H12 N2 O

L56 ANSWER 35 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




L56 ANSWER 36 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Fluoropolymer-polyether-polyketones were obtained by homopolycondensation of 4-hydroxy-4'-(pentafluorobenzoyl)diphenyl ether and by copolycondensation of 4,4'-bis(pentafluorobenzoyl)diphenyl ether with benzenediols or bisphenols. The polymers have very good heat resistance, with 10% weight loss temps. >500°.  
 ACCESSION NUMBER: 1998:532324 CAPLUS  
 DOCUMENT NUMBER: 129:260978  
 TITLE: New polymers derived from 2,3,4,5,6-pentafluorobenzoic acid  
 AUTHOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.; Fitch, John W., III; Reddy, V. Sreenivasulu; Sakaguchi, Yoshimitsu  
 CORPORATE SOURCE: Faculty of Environmental Science and Technology, Okayama University, Okayama, 700-8530, Japan  
 SOURCE: Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1998), 39(2), 790-791  
 PUBLISHER: American Chemical Society, Division of Polymer Chemistry  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 213693-15-3P, 4,4'-Bis(pentafluorobenzoyl)diphenyl ether-hydroquinone copolymer, SRU  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of thermally stable fluoropolymer-polyether-polyketones)  
 RN 213693-15-3 CAPLUS  
 CN Poly[oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene]carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene) (9CI) (CA INDEX NAME)

PAGE 1-A




L56 ANSWER 36 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-B




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L56 ANSWER 37 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A series of bis(ether amines), primarily 1,2-bis(4-aminophenoxy)benzenes but including 1,3- and 1,4-bis(4-aminophenoxy)benzenes, with fluoro and alkyl substituents were synthesized. These diamines were prepared by F displacement with 4-FC6H4NO2 or its deriva., or NO2 displacement with 1,4-C6H4(NO2)2, and various phenylenediols and their alkyl or fluoro deriva. The resulting bis(ether nitro) compds. were reduced to the corresponding bis(ether amines).  
 ACCESSION NUMBER: 1998:376601 CAPLUS  
 DOCUMENT NUMBER: 129:135942  
 TITLE: Methyl- and fluoro-substituted bis(4-aminophenoxy)benzenes. A convenient method of synthesis  
 AUTHOR(S): Eastmond, G. C.; Paprotny, J.  
 CORPORATE SOURCE: Donnan Laboratories, University Liverpool, Liverpool, L69 7ZD, UK  
 SOURCE: Synthesis (1998), (6), 894-898  
 PUBLISHER: Georg Thieme Verlag  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 129:135942  
 IT 210492-55-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (preparation of methylated and fluorinated [(aminophenoxy)phenoxy]benzenamines)  
 RN 210492-55-0 CAPLUS  
 CN Benzene, 1,4-bis(2-fluoro-4-nitrophenoxy)- (9CI) (CA INDEX NAME)



IT 210492-57-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of methylated and fluorinated [(aminophenoxy)phenoxy]benzenamines)  
 RN 210492-57-2 CAPLUS  
 CN Benzenamine, 4,4'-[1,4-phenylenebis(oxy)]bis[3-fluoro- (9CI) (CA INDEX NAME)

L56 ANSWER 37 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




L56 ANSWER 38 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A graded-refractive-index optical plastic material is composed of (a) an amorphous fluoropolymer which substantially has no C-H bond, e.g., a F-containing polyether, aromatic polyester, aromatic polycarbonate, or polyimide, and (b) at least one material which differs from the fluoropolymer in refractive index by at least 0.001, and the material (b) is distributed in the fluoropolymer so as to have a concentration gradient and a continuously varying refractive index in a specific direction. The optical plastic has both high transparency and heat resistance as well as humidity, chemical, and flame resistance, and provides an optical fiber or lens with a low loss for optical communication over short distances. Thus, a perfluoropolyimide (I), prepared from 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene anhydride and bis(2,3,5,6-tetrafluoro-4-aminophenyl)ether, and 15 phr N-pentafluorophenyldichloromaleimide were dissolved in hexafluoroisopropanol (II) (mixture viscosity 3 x 104), and I alone was dissolved in II (viscosity 2 x 104). The 2 mixts. were extruded through a double nozzle to give a fiber-like product having the first mixture on the inside and the latter mixture on the outside, and the II removed to give a title fiber showing light transmission loss at 650-1600 nm and no substantial change after heating 100 h at 150°.

ACCESSION NUMBER: 1998:372628 CAPLUS  
 DOCUMENT NUMBER: 129:42058  
 TITLE: Graded-refractive-index optical perfluoropolymer plastic material, production thereof, and optical products with low transmission loss therefrom  
 INVENTOR(S): Koike, Yasuhiro; Naritomi, Masaki; Murofushi, Hidenobu; Sugiyama, Norihide  
 PATENT ASSIGNEE(S): Koike, Yasuhiro, Japan  
 SOURCE: U.S., 14 pp., Cont.-in-part of U. S. Ser. No. 553,547.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5760139	A	19980602	US 1996-659867	19960607
EP 1249715	A2	20021016	EP 2002-15713	19950412
EP 1249715	A3	20030102		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE				
JP 2002311254	A2	20021023	JP 2002-8228	19950412
JP 08337723	A2	19961224	JP 1995-143710	19950609
JP 3419960	B2	20030623		
US 5783636	A	19980721	US 1995-553547	19951215
US 6271312	B1	20010807	US 1997-984121	19971203
PRIORITY APPLN. INFO.: JP 1994-78828 A 19940418				
JP 1995-143710 A 19950609				

L56 ANSWER 38 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



RN 208453-74-1 CAPLUS  
 CN 1H-Pyrrole-2,5-dione, 3,4-dichloro-1-(pentafluorophenyl)-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] and 2,2,2,2-tetrafluoro-


Page 393

L56 ANSWER 38 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 US 1995-553547 A2 19951215  
 EP 1995-915307 A3 19950412  
 JP 1995-86802 A3 19950412  
 WO 1995-JP715 W 19950412

IT 143376-23-2P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene anhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl)ether copolymer 143433-47-0P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene anhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl)ether copolymer, sru 208453-74-1P  
 RL: IMF (Industrial manufacture); PEP (Physical, engineering or chemical process); TEM (Technical or engineered material use); PREP (Preparation); PROC (Process); USES (Uses)  
 (heat-shrinkable sheath; graded-refractive-index optical perfluoropolymer plastic material, production thereof, and optical products with low transmission loss therefrom)

RN 143376-23-2 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8




CH 2  
 CRN 20115-19-9  
 CHF C12 H4 F8 N2 O

L56 ANSWER 38 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione], graft (9CI) (CA INDEX NAME)


CH 1  
 CRN 186958-58-7  
 CHF C10 C12 F5 N O2



CH 2  
 CRN 143363-91-1  
 CHF C22 F10 O8




CH 3  
 CRN 20115-19-9  
 CHF C12 H4 F8 N2 O



L56 ANSWER 38 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR  
 THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L56 ANSWER 39 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB Selected perfluorinated tertiary alcs. were reacted with  
 p-toluenesulfonyl  
 chloride to form their p-toluenesulfonyl esters C6F5(CF3)2COSO2C6H4CH3  
 (I), CF3C6F4OC6F4(C3F7)(C8F17)COSO2C6H4CH3 and  
 (CF3C6F4OC6F4)2(C7F15)COSO2C6H4CH3. The absolute configuration of I is  
 established by X-ray diffraction.  
 ACCESSION NUMBER: 1998:210377 CAPIUS  
 DOCUMENT NUMBER: 128:321424  
 TITLE: p-Toluenesulfonyl esters of perfluorinated tertiary  
 alcohols: crystal structure determination of the  
 absolute configuration of C6F5(CF3)2COSO2C6H4CH3  
 AUTHOR(S): Krumm, Burkhard; Vij, Ashwani; Kirchmeier, Robert L.;  
 Shreeve, Jean'ne M.  
 CORPORATE SOURCE: Dep. of Chem., Univ. of Idaho, Moscow, ID,  
 83844-2343,  
 SOURCE: USA  
 Journal of Fluorine Chemistry (1998), 89(1), 19-22  
 CODEN: JFLCAR; ISSN: 0022-1139  
 PUBLISHER: Elsevier Science S.A.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 128:321424  
 IT 207233-54-3  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal  
 structure determination of the absolute configuration of  
 C6F5(CF3)2COSO2C6H4CH3)  
 RN 207233-54-3 CAPIUS  
 CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentadecafluoroheptyl)-  
 $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]- (9CI) (CA  
 INDEX  
 NAME)


L56 ANSWER 39 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




IT 207233-53-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal  
 structure determination of the absolute configuration of  
 C6F5(CF3)2COSO2C6H4CH3)  
 RN 207233-53-2 CAPIUS  
 CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentadecafluoroheptyl)-  
 $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]-,  
 4-methylbenzenesulfonate (9CI) (CA INDEX NAME)

L56 ANSWER 39 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR  
 THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

AB Title linear polyethers consist of (A) bisphenol units having 2 benzene rings, each of which is directly linked to O and (B) aryl-type units for linking the bisphenol units. The polymers are useful as precursors of intermediate elec. insulator films in semiconductor devices and the resulting films can be subjected to high temperature reactive ion etching and they leave no Si oxide residue after etching. The polyethers are manufactured by polymerizing aromatic monomers corresponding to the unit A and compds. having 21 benzene rings which are directly linked to 22 halogens in the presence of basic catalysts. Thus, 28.6 g 2,2'-binaphthol and 33.4 g perfluorobiphenyl were polymerized in N,N-dimethylacetamide in the presence of K2CO3 at 80° under N to give title polymer, 5 g of which was dissolved in 50 mL 2-methoxyethyl acetate, spin-coated on a Si substrate, heated in air at 200° for 30 min, and baked at 360° in N for 1 h to give an elec. insulator film having sp. dielec. constant 2.70, glass-transition temperature 250°, and no weight degradation at 400-500°.

ACCESSION NUMBER: 1998:176533 CAPLUS  
DOCUMENT NUMBER: 128:244969  
TITLE: Silicon-free bisphenol-type polyethers as electric insulator film precursors and their manufacture  
INVENTOR(S): Ito, Toshio; Kosuga, Maki  
PATENT ASSIGNEE(S): Oki Electric Industry Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
CODEN: JGQCAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10074751	A2	19980317	JP 1996-229893	19960830
JP 3195248	B2	20010806		

PRIORITY APPLN. INFO.: JP 1996-229893 19960830

IT 204910-65-6P  
RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(silicon-free bisphenol-type polyethers as precursors for elec. insulators in semiconductor devices)  
RN 204910-65-6 CAPLUS  
CN Poly[oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,2',3,3',5,5',6,6'-octafluoro[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

AB The devices consist of direction coupler switches or Y-splitters and Mach-Zehnder optical modulator where the coupler switches are of core-sheath type and have the core or/and the sheath made from fluoro polyimide resins for improving resistance to heat and moisture while having good transmission rate. In an example, a fluoro polyimide resin was obtained from a 6:4 mixture of 2,2-bis[3,4-dicarboxyphenyl]hexafluoropropane dianhydride-2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl copolymer and 2,2-bis[3,4-dicarboxyphenyl]hexafluoropropane dianhydride-4,4'-oxydianiline copolymer.


ACCESSION NUMBER: 1998:163257 CAPLUS  
DOCUMENT NUMBER: 128:276864  
TITLE: Waveguide-type optical devices with low transmission loss and good resistance to heat and moisture  
INVENTOR(S): Kobayashi, Junya; Maruno, Toru; Matsura, Toru; Sasaki, Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 14 pp.  
CODEN: JGQCAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10068829	A2	19980310	JP 1997-94360	19970331
			JP 1996-178683	19960620

PRIORITY APPLN. INFO.: JP 1996-178683 19960620


IT 143376-21-0P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine copolymer  
143376-23-2P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) ether copolymer  
143333-45-8P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-tetrafluoro-m-phenylenediamine copolymer, polyimide sru  
143333-47-0P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) ether copolymer, polyimide sru 148362-03-2P, 4,4'-2,2-Bis[4-

aminophenyl]hexafluoropropane-1,4-bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride copolymer, polyimide sru 148362-05-4P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-4,4'-oxydianiline copolymer, polyimide sru 148362-06-5P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl copolymer, polyimide sru 148446-36-0P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl copolymer 148446-37-1P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-4,4'-oxydianiline copolymer 188412-73-9P, 2,2-Bis[3-amino-4-hydroxyphenyl]hexafluoropropane-1,4-bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride copolymer 188412-75-1P, 2,2-Bis[3-amino-4-hydroxyphenyl]hexafluoropropane-1,4-bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride copolymer, polyimide sru 205451-27-0P, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride-3,4'-oxydianiline




copolymer 205451-28-1P, 4,4'-2,2-Bis[4-aminophenyl]hexafluoropropane-1,4-bis[3,4-dicarboxytrifluorophenoxy]tetrafluorobenzene dianhydride copolymer  
RL: DEV (Device component use); IMF (Industrial manufacture); PRP (Properties); PREP (Preparation); USES (Uses)  
(resins for manuf. of waveguide-type optical devices with low transmission loss and good resistance to heat and moisture)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CMF C22 F10 O8



CH 2  
CRN 1198-63-6  
CMF C6 H4 F4 N2




RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)  
CH 1  
CRN 143363-91-1




CN 2

CRN 20115-19-9  
 CNF C12 H4 F8 N2 O




RN 143433-45-8 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-

PAGE 1-A




PAGE 1-B




RN 148362-05-4 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenyleneoxy-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 148362-03-2 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148362-06-5 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2,2'-(bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diyl)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B

RN 148446-36-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8




CM 2

CRN 341-58-2  
 CMF C14 H10 F6 N2




RN 148446-37-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-




CM 2

CRN 83558-87-6  
 CMF C15 H12 F6 N2 O2




RN 188412-75-1 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[6-hydroxy-1,3-phenylene][2,2,2-trifluoro-1-(trifluoromethyl)ethylidene](4-hydroxy-1,3-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A




CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2

CRN 101-80-4  
 CMF C12 H12 N2 O




RN 188412-73-9 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
 CMF C22 F10 O8


PAGE 1-B



RN 205451-27-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 3-(4-aminophenoxy)benzenamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2

CRN 2657-87-6  
 CMF C12 H12 N2 O




RN 205451-28-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-

L56 ANSWER 41 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 phenylene)bis(oxy))bis[4,6,7-trifluoro-, polymer with 4,4'-[(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)bis(4,1-phenyleneoxy)]bis(benzenamine) (9CI) (CA INDEX NAME)


CH 1

CRN 143363-91-1  
 CMF C22 F10 O8




CH 2

CRN 69563-88-8  
 CMF C27 H20 F6 N2 O2



L56 ANSWER 42 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN GI



AB There is provided a low-birefringent organic optical component comprising a polymer prepared by polymerizing a racemic mixture of the monomer with an asym. spiro ring represented by general formula I wherein ring A represents a monocyclic or polycyclic organic group, wherein two ring As are mutually bound each other via a spiro bond to form a spiro ring which has a mol. asym. structure; n is an integer of 0 to 10; X1 and Y2 are binding groups;

X1 is a polymerization-active group. The component has excellent transparency, mech. strength and heat resistance and useful in manufacture of optical disks and optical lenses and prisms.

ACCESSION NUMBER: 1998:147152 CAPIUS  
 DOCUMENT NUMBER: 128:210916

TITLE: Optical component and spirobiindan polymer therefor  
 INVENTOR(S): Otsuji, Atsuo; Takuma, Keisuke; Suzuki, Rihoko; Urakami, Tatsuhiro; Motoshima, Toshihiro; Yamashita, Wataru; Yoshimura, Tomomi; Shibuya, Atsushi; Sakata, Yoshihiro; Oikawa, Hideaki; Ohta, Masahiro; Ajioka, Masanobu; Takagi, Masatoshi; et al.  
 SOURCE: Mitsui Toatsu Chemicals, Inc., Japan  
 Eur. Pat. Appl., 147 pp.  
 CODEN: EPOKDW

DOCUMENT TYPE: Patent  
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 822545	A2	19980204	EP 1997-305763	19970731
EP 822545	A3	19980624		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
US 6080833	A	20000627	US 1997-896008	19970717
JP 11071316	A2	19990316	JP 1997-196009	19970722
CN 1175600	A	19980311	CN 1997-120511	19970731
CN 1093147	B	20021023		
PRIORITY APPLN. INFO.:				
			JP 1996-201825	A 19960731
			JP 1996-204614	A 19960802
			JP 1996-204615	A 19960802
			JP 1996-331831	A 19961212


L56 ANSWER 42 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

JP 1996-331832 A 19961212  
 JP 1996-331833 A 19961212  
 JP 1997-138355 A 19970528  
 JP 1997-138356 A 19970528  
 JP 1997-138357 A 19970528  
 JP 1997-138358 A 19970528  
 JP 1997-138359 A 19970528  
 JP 1997-159421 A 19970617  
 JP 1997-159422 A 19970617  
 JP 1997-159423 A 19970617  
 JP 1997-159424 A 19970617

IT 203712-59-8P  
 RL: DEV (Device component use); SPN (Synthetic preparation); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses) (preparation and use as material for optical disks and lenses)  
 RN 203712-59-8 CAPIUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy))bis[4,6,7-trifluoro-, polymer with 4,4'-[(2,2',3,3'-tetrahydro-3,3,3',3'-tetramethyl-1,1'-spirobi[1H-indene]-6,6'-diyl)bis(oxy))bis(benzenamine) (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
 CMF C22 F10 O8



CH 2

CRN 112147-64-5  
 CMF C33 H34 N2 O2

L56 ANSWER 42 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)






L56 ANSWER 43 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB Mixing a resin and a certain phthalocyanine compound having the ability  
 to  
 absorb near IR rays, provides a resin composition which is useful as a  
 heat  
 radiation-shielding material which is semi-transparent or transparent for  
 visible light but blocks heat rays. C black may also be added to enhance  
 the heat-shielding effect. Thus, polycarbonate containing 0.003%  
 VOPc(BuNH)8F8 (Pc = phthalocyanine; substituent  
 octakis(butylamino)octafluorophthalocyanine) was molded into a film  
 having transmittance and heat transmittance (JIS R-3106) 78.9 and 62.4%  
 vs. 89.2 and 84.6, resp., for polycarbonate without the IR absorber.  
 ACCESSION NUMBER: 1998:79369 CAPIUS  
 DOCUMENT NUMBER: 128:141503  
 TITLE: Compositions for shields for heat radiation  
 INVENTOR(S): Kaleda, Osamu; Yodoshi, Takashi; Morita, Ken;  
 Matsura, Michio  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Japan  
 SOURCE: U.S., 15 pp., Cont.-in-part of U.S. Ser. No. 180,488,  
 abandoned.  
 CODEN: USXQAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5712332	A	19980127	US 1995-478739	19950607
PRIORITY APPL. INFO.:			JP 1993-4326	A 19930113
			US 1994-180488	B2 19940112


OTHER SOURCE(S): MARPAT 128:141505  
 IT 163464-84-4  
 RL: MQA (Modifier or additive use); USES (Uses)  
 (polymer compns. containing certain phthalocyanines for shields for  
 heat  
 radiation)  
 RN 163464-84-4 CAPIUS  
 CN Zinc, [N,N',N'',N'''-tetrahexyl-1,3,4,8,10,11,15,17,18,22,24,25-  
 dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-  
 tetraminato(2-)-κN29,κN30,κN31,κN32]-, (SP-4-1)-  
 (9CI) (CA INDEX NAME)

L56 ANSWER 43 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A




PAGE 2-A




L56 ANSWER 43 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 3-A



Me-(CH2)5-NH-R2

PAGE 4-A




REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR  
 THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L56 ANSWER 44 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The title device utilizes a low operation voltage and shows good  
 displaying quality for a long time period and excellent low energy  
 consumption properties, wherein the device utilizes liquid crystals  
 having a  
 threshold voltage of ≤1.5 V and a F-containing polymer orientation  
 layer.  
 ACCESSION NUMBER: 1997:736287 CAPIUS  
 DOCUMENT NUMBER: 128:28680  
 TITLE: Liquid crystal display device for handy information  
 terminal  
 INVENTOR(S): Kawagoe, Jun; Ihara, Satoru; Ooaku, Hitoshi  
 PATENT ASSIGNEE(S): Optrex Corp., Japan  
 SOURCE: Ger. Offen., 6 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19710698	A1	19971106	DE 1997-19710698	19970314
JP 09304776	A2	19971128	JP 1997-36601	19970220
US 5781263	A	19980714	US 1997-816895	19970313
PRIORITY APPL. INFO.:			JP 1996-57954	A 19960314

IT 143376-23-2 143433-47-0  
 RL: DEV (Device component use); USES (Uses)  
 (orientation layer of liquid crystal display device for handy  
 information  
 terminal)  
 RN 143376-23-2 CAPIUS  
 CN 1,3-bis(isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-  
 phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with  
 4,4'-oxybis[2,3,5,6-  
 tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)


CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8




CH 2



PAGE 1-A





PAGE 1-B



L56 ANSWER 45 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

The chemical structure shows three identical 2,3,5,6-tetrafluorophenyl groups linked sequentially by ether bonds (-O-) at their para positions. The first group is substituted at its ortho position with a 1H-imidazo[1,2-a]pyridin-7-yl ring, which has a nitro group (-NO<sub>2</sub>) at its 2-position.



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR  
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB The invention relates to a heat-resistant polymer optical waveguide, suited for use in applications requiring higher degree of optical integration, wherein  $\geq 3\%$  of refractive index difference between the cladding and the core is achieved by selecting appropriate combination of polyimides for the cladding and the core. The core may be prepared from

the polyimide having the structural repeating unit represented by I or II (R1 is represented by III, IV, and V; R2 = (p-C6H4)2O, (p-C6H4)2CH2, (p-C6H4)2SO2, and (m-C6H4)2O2). The cladding may be prepared from the polyimide having the structural repeating unit represented by VI (R3 = 2,2'-bis(trifluorophenyl)biphenyl-4,4'-diyl, (p-C6H4)2C(CF3)2, (p-C6H4)2O,

and pentafluorophenoxy phenyl-2,4-diyl).

ACCESSION NUMBER: 1997:633003 CAPLUS

DOCUMENT NUMBER: 127:339030

TITLE: Heat-resistant polymer optical waveguide and its

production method

INVENTOR(S): Matsuura, Toru; Maruno, Toru; Kobayashi, Junya;

Sakata, Tomomi; Sasaki, Shigekuni

PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 11 pp.

CODEN: JKOQAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09251113	A2	19970922	JP 1996-85964	19960315

PRIORITY APPL. INFO.:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09251113	A2	19970922	JP 1996-85964	19960315

IT 143376-21-OP 143376-23-2P 143433-45-8P

143433-47-OP, 1,4-Bis[3,4-dicarboxytrifluorophenoxy]tetrafluoroben

zene dianhydride-bis[2,3,5,6-tetrafluoro-4-aminophenyl]ether

copolymer, sru

148362-03-2P 148362-05-4P 148362-06-5P

148446-36-OP 148446-37-1P 148446-40-6P

108412-73-9P 108412-75-1P

RL: DEV (Device component use); PMU (Preparation, unclassified); PREP

(Preparation); USES (Uses)

(heat-resistant polymer optical waveguide and its production method)

RN 143376-21-0 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-

phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with

2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CH 1

CRN 143363-91-1

CHF C22 F10 O8

L56 ANSWER 46 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


CRN 20115-19-9  
CHF C12 H4 F8 N2 O



RN 143433-45-8 CAPLUS

CN

Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)


L56 ANSWER 46 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



CH 2

CRN 1198-63-6

CHF C6 H4 F4 N2




RN 143376-23-2 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)

CH 1

CRN 143363-91-1


CHF C22 F10 O8




CH 2

L56 ANSWER 46 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B




RN 148362-03-2 CAPLUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 148362-05-4 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenyleneoxy-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148362-06-5 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diyl]] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148446-36-0 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2


CRN 341-58-2  
 CMF C14 H10 F6 N2



RN 148446-37-1 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2


CRN 101-80-4  
 CMF C12 H12 N2 O




RN 148446-40-6 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[benzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
 CMF C22 F10 O8




CM 2


CRN 1095-78-9  
CMF C15 H12 F6 N2

RN 188412-73-9 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,4'-(2,2,2-trifluoro-1-(trifluoromethyl)ethylidene)bis[2-aminophenyl] (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
CMF C22 F10 O8

CM 2



CRN 83558-87-6  
CMF C15 H12 F6 N2 O2

RN 188412-75-1 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(6-hydroxy-1,3-phenylene)[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene](4-hydroxy-1,3-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



AB Phthalic acids I (R = CO<sub>2</sub>H; X = halo) are prepared by heating phthalonitriles I (R = cyano; X = halo) in acidic aqueous media, (A) dissolving the products in organic solvent-containing media, (B) separating the products, and (C) treating the isolated products in acidic aqueous media under heating. Then, the above processes A to C are repeated. ≥1 time(s). I (R = cyano, X = F) (10 g) was refluxed with aqueous H<sub>2</sub>SO<sub>4</sub> for 6 h,

diluted with H<sub>2</sub>O, filtered, and the crude product was dissolved in Me<sub>2</sub>CO. The solution was treated with H<sub>2</sub>O to give 10.8 g I (R = CO<sub>2</sub>H, X = F) (II) with purity 91.8%, which was refluxed with H<sub>2</sub>SO<sub>4</sub> for 6 h and similarly treated. The process was repeated to give 9.8 g II with 99.2% purity.

ACCESSION NUMBER: 1997:377463 CAPLUS  
DOCUMENT NUMBER: 127:33992  
TITLE: Preparation of phthalic acids as intermediates for polyimides from phthalonitriles  
INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Maeda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
CODEN: JXOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110785	A2	19970428	JP 1995-266007	19951013
PRIORITY APPLN. INFO.:			JP 1995-266007	19951013


OTHER SOURCE(S): MARPAT 127:33992

IT 143363-92-2P


RL: IMP (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)  
(preparation and purification of phthalic acids as intermediates for polyimides by acid hydrolysis of phthalonitriles)

RN 143363-92-2 CAPLUS


CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




IT 143376-50-5  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation and purification of phthalic acids as intermediates for  
 polyimides by acid hydrolysis of phthalonitriles)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-(2,3,5,6-tetrafluoro-1,4-  
 phenylene)bis(oxy)bis[3,5,6-trifluoro-9CI] (CA INDEX NAME)




L56 ANSWER 48 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of phthalic acids as intermediates for polyimides by  
hydrolysis of phthalonitriles and heating with alkalis)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[2,3,5,6-tetrafluoro-1,4-  
phenylene]bis(oxy)]bis[3,5,6-trifluoro- (SCI) (CA INDEX NAME)



L56 ANSWER 48 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
GI



AS Title compds. I (X = halo; Y = CO<sub>2</sub>H) are prepared by heating I (Y = cyano) in acidic aqueous media, heating the product-containing media with alkaline substances, and separating I (Y = CO<sub>2</sub>H) from the media by mixing with acidic substances. An aqueous solution containing 10 g I (X = F, Y = cyano) and H<sub>2</sub>SO<sub>4</sub> was refluxed for 6 h, mixing with H<sub>2</sub>O and NaOH (pH 11.8), refluxed for 1 h, and mixed with aqueous H<sub>2</sub>SO<sub>4</sub> to give 10.8 g I (X = F, Y = CO<sub>2</sub>H) with 99.2%.

ACCESSION NUMBER: 1997:377462 CAPLUS  
 DOCUMENT NUMBER: 127:33986  
 TITLE: Preparation of phthalic acids as intermediates for  
 polyimides  
 INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Kaieda, Osamu  
 PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110784	A2	19970428	JP 1995-266006	19951013
PRIORITY APPLN. INFO.:			JP 1995-266006	19951013

OTHER SOURCE(S): MARPAT 127:33986  
IT 143363-92-2P  
RL: IMF (Industrial manufacture); PUR (Purification or recovery); SPN  
(Synthetic preparation); PREP (Preparation)  
REMARKS: Preparation of phthalic acids as intermediates for polyimides by  
hydrolysis  
of phthalonitriles and heating with alkalis)  
143363-92-2 CAPLUS  
CN 1,2-Benzidenedicarboxylic acid, 4,4'-(2,3,5,6-tetrafluoro-1,4-  
phenylenebis(alkoxy))bis[3,5,6-trifluoro-9(ICI) (CA INDEX NAME)

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS ON STN  
AB The invention relates to a polyimide optical waveguide comprising  
polyimide core and cladding layers fabricated on a substrate, wherein the  
nTE - nTM is  $\leq 0.003$  at  $1.3 \mu\text{m}$  in both core and cladding layers  
for improving the polarization properties of the waveguide, where the nTE  
and nTM are the refractive indexes of the polyimide in the directions  
parallel and perpendicular to the substrate surface, resp.

ACC. NO. 1977-250653 CAPLUS  
 ASSIGNMENT NUMBER: 126:24464:  
 DOCUMENT NUMBER: 1977-250653  
 TITLE: Polyimide optical waveguide  
 INVENTOR(S): Matsura, Toru; Koshobu, Nobutake; Maruno, Toru;  
 Ando, Shinji; Sakata, Tomomi; Sasaki, Shigekuni;  
 Kobayashi, Junya  
 PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
 CODEN: YKOGAK  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09021920	A2	19970121	JP 1995-194040	19950707
PRIORITY APPLN. INFO.:			JP 1995-194040	19950707

IT 143376-21-0 143376-23-2 143433-47-0  
140362-05-4 140362-06-5 140446-36-0  
140446-37-1 108412-73-9 108412-75-1  
108442-26-4  
RL: DEV (Device component use); USES (Uses)  
(polyimide optical waveguide)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzoxafurandione, 5,5'-(2,3,5,6-tetra-  
phenylene)bis(oxy)bis(4,6,7-trifluoro-, po-  
2,4,5,6-tetrafluoro-  
1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1

CMF C22 F10 08



CH 2


L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 CRN 1198-63-6  
 CMF C6 H4 F4 N2



RN 143376-23-2 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2


CRN 20115-19-9  
 CMF C12 H4 F8 N2 O




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A




PAGE 1-B




RN 148362-06-5 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)[2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diyl]] (9CI) (CA INDEX NAME)

PAGE 1-A




L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 1,3-dioxo-2H-isindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



RN 148362-05-4 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)-1,4-phenyleneoxy-1,4-phenylene] (9CI) (CA INDEX NAME)

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


PAGE 1-B



RN 148446-36-0 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CMF C22 F10 O8



CM 2

CRN 341-58-2  
 CMF C14 H10 F6 N2




RN 148446-37-1 CAPLUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
phenylene)bis(oxy)bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis(benzenamine) (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8



CM 2

CRN 101-80-4  
CMF C12 H12 N2 O




RN 188412-73-9 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis(4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
CMF C22 F10 O8

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




CM 2

CRN 83558-87-6  
CMF C15 H12 F6 N2 O2




RN 188412-75-1 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(6-hydroxy-1,3-phenylene)[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene](4-hydroxy-1,3-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


PAGE 1-B



RN 188442-26-4 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis(4,6,7-trifluoro-, polymer with ar,ar'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[benzenamine] (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1  
CMF C22 F10 O8



CM 2

CRN 30606-80-5  
CMF C15 H12 F6 N2  
CCI IDS

L56 ANSWER 49 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)





L56 ANSWER 50 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Moldings contain amorphous fluoropolymers such as fluoro polyethers, aromatic polyesters, aromatic carbonates, and polyimides containing no C-H bonds and 21 material which has refractive index difference  $\geq 0.001$  with the fluoropolymers and is distributed with a concentration gradient along a specified direction. Thus, optical fibers contained a perfluoro polyether triazine and 1,3-dibromotetrafluorobenzene and had n decreasing from the center to the periphery and optical transmission 380 dB/km at 780 nm and 250 dB/km at 1550 nm.


ACCESSION NUMBER: 1997:165106 CAPLUS  
 DOCUMENT NUMBER: 126:158497  
 TITLE: Graded-index optical resin materials  
 INVENTOR(S): Koike, Yasuhiro; Murofushi, Hidenobu; Sugiyama, Tokuhide  
 PATENT ASSIGNEE(S): Koike Yasuhiro, Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
 CODEN: JKO0AF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08337723	A2	19961224	JP 1995-143710	19950609
JP 3419960	B2	20030623		
US 5760139	A	19980602	US 1996-659867	19960607
			JP 1994-78828	A 19940418
			JP 1995-143710	A 19950609
			US 1995-553547	A2 19951215


IT 143376-23-2 143433-47-0  
 RL: POF (Polymer in formulation); PRP (Properties); TEM (Technical or engineered material use); USES (Uses)  
 (graded-index optical fibers containing amorphous fluoropolymers and additives)  
 RN 143376-23-2 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis(2,3,5,6-tetrafluorobenzeneamine) (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8

L56 ANSWER 50 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




CH 2  
 CRN 20115-19-9  
 CHF C12 H4 F8 N2 O




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)


PAGE 1-A



L56 ANSWER 50 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 PAGE 1-B



L56 ANSWER 51 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI




AB The title bisphthalonitriles I (X, Y = O, S; R = Z, Z1), useful as intermediates for optical materials, circuit board materials, photosensitive materials, liquid crystals, etc., are prepared by treatment of 3,4,5,6-tetrafluorophthalonitrile (II) with HXRYH in organic solvents being slightly soluble in H2O in the presence of basic substances, followed by addition of H2O to the reaction products and separation of the organic layer. An AcOEt solution of tetrafluorohydroquinone was added dropwise to a mixture of II, KF, and AcOEt under reflux over 1 h, and the reaction mixture was further stirred under reflux for 8 h, cooled, and then washed with H2O to give 99% I (X = Y = O, R = Z).

ACCESSION NUMBER: 1997:154684 CAPLUS  
 DOCUMENT NUMBER: 126:157295  
 TITLE: Preparation of bis(fluorophthalonitriles) from tetrafluorophthalonitrile and fluorohydroquinone or fluorodiphenol  
 INVENTOR(S): Okumura, Yasunori; Kaieda, Osamu  
 PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
 CODEN: JKO0AF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08333322	A2	19961217	JP 1995-143593	19950609
			JP 1995-143593	19950609

OTHER SOURCE(S): CASREACT 126:157295; MARPAT 126:157295  
 IT 143376-50-5P  
 RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP

L56 ANSWER 51 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
(Preparation)  
(prepn. of bis(fluorophthalonitriles) from tetrafluorophthalonitrile  
and fluorohydroquinone, diphenol, or their sulfur analog using  
slightly water-sol. solvents and basic substances)  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-  
phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



L56 ANSWER 52 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
GI




AB Title tapes are obtained by laminating UV-curable adhesive layers on  
21 side of fluorinated polyimide support films. The tapes show  
good productivity and dimensional stability and are useful for fixation  
of  
lead pins at lead frames. Thus, XE 17A5537 (silicone-based UV-curable  
adhesive) was applied on a polyimide I film to give an adhesive tape,  
which was bonded with Ni-Fe alloy and UV-cured to give a test piece

having  
peeling strength 255 g/15 mm.  
ACCESSION NUMBER: 1997:61080 CAPIUS  
DOCUMENT NUMBER: 126:105097  
TITLE: Adhesive tapes for fixation of electronic parts in  
semiconductor devices  
INVENTOR(S): Tanaka, Yoshiharu; Taguchi, Yoshio  
PATENT ASSIGNEE(S): Bando Chemical Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 4 pp.  
CODEN: JN0047  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08283677	A2	19961029	JP 1995-112470	19950412
PRIORITY APPL. INFO.:			JP 1995-112470	19950412

IT 143376-21-0 143433-45-8  
RL: TEM (Technical or engineered material use); USES (Uses)  
(adhesive tapes with good productivity and dimensional stability for  
fixation of electronic devices)  
RN 143376-21-0 CAPIUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-  
phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with  
2,4,5,6-tetrafluoro-  
1,3-benzenediamine (9CI) (CA INDEX NAME)  
CM 1  
CRN 143363-91-1  
CHF C22 F10 08


L56 ANSWER 52 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




CM 2  
CRN 1198-63-6  
CHF C6 H4 F4 N2



RN 143433-45-8 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-  
tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-  
isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA  
INDEX NAME)



L56 ANSWER 53 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
GI




AB Polyimide optical materials comprise a perfluorinated polyimide having a  
perfluorinated repeating unit represented by the general formula I (R1 =  
a  
tetravalent perfluorinated organic group; and R2 = a divalent  
perfluorinated  
organic group).

ACCESSION NUMBER: 1995:854348 CAPIUS  
DOCUMENT NUMBER: 123:354214  
TITLE: Polyimide optical material  
INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni;  
Yamamoto, Fumio  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corporation, Japan  
SOURCE: U.S., 26 pp. Cont.-in-part of U.S. Ser. No. 54,973,  
abandoned.  
CODEN: USXXGM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5449741	A	19950912	US 1993-140982	19931025
JP 09031014	A2	19970204	JP 1996-207475	19910913
US 5233018	A	19930803	US 1991-765672	19910926
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 5750731	A	19980512	US 1996-718208	19960920
US 5849934	A	19981215	US 1998-20573	19980128
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		
PRIORITY APPL. INFO.:			JP 1990-256843	A 19900928
			JP 1991-106552	A 19910412
			JP 1991-106554	A 19910412
			JP 1991-106557	A 19910412
			US 1991-765672	A3 19910926
			US 1993-54973	B2 19930430

L56	ANSWER 53 OF 89	CAPLUS	COPYRIGHT 2005 ACS on STM	(Continued)
			JP 1991-235020	A3 19910913
			CA 1991-2052368	A3 19910926
			US 1993-140482	A3 19931025
			US 1993-140982	A3 19931025
			US 1995-451465	A1 19950526
			US 1996-718208	A3 19960920
			US 1998-20573	A3 19980128

IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(polyimide optical materials)  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




IT	143363-91-1P, 1,4-Bis-(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (polyimide optical materials)
RN	143363-91-1 CAPLUS
CN	1,3-Isobenzofurandione, 5,5'-(2,3,5,6-tetrafluoro-1,4-phenylenebis(oxy))bis[4,6,7-trifluoro- (GCI) (CA INDEX NAME)



IT 143376-21-0P 143376-22-1P 143376-23-2P


L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
143376-24-32 143433-45-8P 143433-46-2P  
143433-47-0P 143433-48-1P  
RL: SPN (Synthetic preparation); TEM (Technical or engineered material  
use); PREP (Preparation); USES (Uses)  
(polyimide optical materials)  
RN 143376-21-0 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-(1,2,3,5,6-tetrafluoro-4,  
phenylene)bis(oxy)|bis[4,6,7-trifluoro-, polymer with  
2,4,5,6-tetrafluoro-  
1,3-benzenediamine (9CI) (CA INDEX NAME)  
CN 1  
CRN 143363-91-1  
CHF C22.F10.08



CM 2

CRN 1198-63-6

CMF C6 H4 F4 N2



```


RN      143376-22-1  CAPLWS
CN      1,3-isobenzofurandione, 5,5'-[ (2,3,5,6-tetrafluoro-1,4-
        phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with
        2,3,5,6-tetrafluoro-
        1,4-benzenediamine (9CI)  (CA INDEX NAME)

CM      1

CRN     143363-91-1
CMF     C22 F10 08

```


L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



CM 2

CRN 1198-64-7

CMF C6 H4 F4 N2




```

RN      143376-23-2  CAPLUS
CM      1,3-Isobenzoxfurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-
        phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with
        4,4'-oxybis[2,3,5,6-
        tetrafluorobenzeneamine] (9CI)  (CA INDEX NAME)

CM      1

CRN     143363-91-1
CMF     C22 F10 O8


```



CM 2

CRN 20115-19-9

L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CMF C12 H4 F8 N2 O




```

RN      143376-24-3  CAPIWS
CN      1,3-bis(methylthio)furan-2-one, 5,5'-[(2,3,5,6-tetrafluoro-1,4-
        phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with
        4,4'-thiobis[2,3,5,6-
        tetrafluorobenzenamine] (9CI)  (CA INDEX NAME)

CH      1

CRM     143363-91-1
CMF     C22 F10 O8


```



CM 2


CRN 61907-46-8

CMF C12 H4 F8 N2 S




RN 143433-45-8 CAPIUS  
CN  
Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl) (2,4,5,6-tetrafluoro-1,3-phenylene) (4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-

L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA  
 INDEX NAME)



RN 143433-46-9 CAPLUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B




RN 143433-48-1 CAPLUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)thio(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A



PAGE 1-B




IT 143363-92-2

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reg: polyimide optical materials)

RN 143363-92-2 CAPLUS

CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(3,5,6-trifluoro- (9CI) (CA INDEX NAME)

L56 ANSWER 53 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



L56 ANSWER 54 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB High resolution fluorine-19 NMR spectra of polyfluoroarom. compds.  
 dissolved  
 in deuterated DMSO were measured and substituent shielding parameters  
 were


derived. These parameters were compared with the values observed in  
 deuterated chloroform solns. and used to identify fluoroarom. compds.  
 related to perfluorinated polyimides and poly(amic acids)s. Average  
 differences of  $\Delta\delta$  between the solvents are 1.1, 1.0 and 2.1  
 ppm for o-, m- and p-fluorines of monosubstituted pentafluorobenzenes,  
 resp. The significant difference for p-fluorine of NH<sub>2</sub> is important in  
 identifying perfluorinated diamines because they are source materials for  
 perfluorinated polyimides. Substituent shielding parameters for meta and  
 para substitution increase as the resp. Hammett  $\sigma$  consts. increase,  
 which indicates that fluorine-19 NMR chemical shift is primarily

determined by  
 electron d.  
 ACCESSION NUMBER: 1995:736766 CAPIUS  
 DOCUMENT NUMBER: 123:227514  
 TITLE: Substituent shielding parameters of fluorine-19 NMR  
 on

polyfluoroaromatic compounds dissolved in dimethyl  
 sulfoxide-d6  
 AUTHOR(S): Ando, Shinji; Matsuura, Tohru  
 CORPORATE SOURCE: NTT Interdisciplinary Research Laboratories, Tokyo,  
 180, Japan  
 SOURCE: Magnetic Resonance in Chemistry (1995), 33(8), 639-45  
 CODEN: MRCHG; ISSN: 0749-1581


PUBLISHER: Wiley  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143363-91-1 143363-92-2 143376-21-0  
 143376-50-5 143433-45-8 168417-33-2  
 RL: PRP (Properties)  
 (substituent shielding parameters in fluorine-19 NMR study of  
 polyfluoroarom. compds.)

RN 143363-91-1 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-  
 phenylene]bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)




RN 143363-92-2 CAPIUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-[2,3,5,6-tetrafluoro-1,4-  
 phenylene]bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)


L56 ANSWER 54 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




RN 143433-45-8 CAPIUS  
 CN  
 Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-  
 tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-  
 isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA  
 INDEX NAME)



RN 168417-33-2 CAPIUS  
 CN 1,2-Benzenedicarboxylic acid, 3,4,6-trifluoro-5-[2,3,5,6-tetrafluoro-4-  
 (4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-5-isobenzofuranyl)oxy]phenoxy]-  
 (9CI) (CA INDEX NAME)




L56 ANSWER 54 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




RN 143376-21-0 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-  
 phenylene]bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CMF C22 F10 O8




CH 2  
 CRN 1198-63-6  
 CMF C6 H4 F4 N2



RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[2,3,5,6-tetrafluoro-1,4-  
 phenylene]bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)

L56 ANSWER 55 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 GI




AB The mixts. contain a transparent resin [e.g., polycarbonate, poly(Me  
 methacrylate), PVC, or poly(ethylene terephthalate)], a near-IR-absorbing  
 phthalocyanine compound I (Z1-16 = SRI, OR2, H, halo, MHY; Z1 of  
 Z1-16 = MHY; R1-2 = Ph, substituted Ph, Cl-20 alkyl; Y = Ph, substituted  
 Ph, Cl-8 alkyl; M = VO, Zn, Cu, SnCl<sub>2</sub>, Co, etc.), and, optionally, carbon  
 black. Replacing part of the I with carbon black does not decrease the  
 heat-radiation-shielding ability of the mixts. The mixts. are useful as  
 moldings which transmit visible light while blocking near-IR radiation  
 (i.e., they absorb the heat from sunlight).

ACCESSION NUMBER: 1995:550904 CAPIUS  
 DOCUMENT NUMBER: 122:316019  
 TITLE: Heat-radiation-shielding mixtures of polymers and  
 near-IR-absorbing phthalocyanine compounds  
 INVENTOR(S): Kaieda, Osamu; Yodoshi, Takashi; Morita, Ken;  
 Matsuura, Michio  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
 SOURCE: Eur. Pat. Appl., 28 pp.  
 CODEN: EPOKDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 607031	A1	19940720	EP 1994-300205	19940112
EP 607031	B1	19991006		
R: BE, DE, ES, FR, GB, IT, NL				
JP 06264050	A2	19940920	JP 1993-352128	19931229
ES 2136700	T3	19991201	ES 1994-300205	19940112

OTHER SOURCE(S): MARPAT 122:316019  
 IT 163464-84-4  
 RL: MOA (Modifier or additive use); POF (Polymer in formulation); PRP (Properties); TEM (Technical or engineered material use); USES (Uses) (near-IR absorber; heat-radiation-shielding compns. containing polymers and)  
 RN 163464-84-4 CAPIUS  
 CN Zinc, [N,N',N'',N'''-tetrahexyl-1,3,4,6,10,11,15,17,18,22,24,25-dodecakis(2,3,5,6-tetrafluorophenoxy)-29H,31H-phthalocyanine-2,9,16,23-tetraminato(2-)-<N29,<N30,<N31,<N32]-, (5P-4-1)-(9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 2-A



PAGE 3-A




PAGE 4-A




L56 ANSWER 56 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB In the presence of fluoride ion, per- or polyfluoroarom. siloxanes C6F5OSi(CH3)3 (1), (CH3)3SiOC6F4OSi(CH3)3 (2), 1-fluoro-2,3-bis((trimethylsilyl)oxy)benzene (3), and (CH3)3SiOC(CF3)2C6F4C(CF3)2OSi(CH3)3 (4) are useful transfer reagents for the synthesis of mono- and diethers. Siloxane 1 forms RfOCF2Br (5), (RfO)3CF (6), RfOCH2ORf (7), (RfO)2CHCH(ORf)2 (8), (RfO)2SO (9), (RfO)2CO (10), C3N3(ORf)3 (11), RfOC(O)(CF2)3C(O)ORf (12), CF3SO2ORf (13), [cyclic] CF2C(ORf):C(ORf)CF2 (14), and [cyclic] CF2C(ORf):C(F)CF2 (15) (Rf = C6F5) with CF2Br2, CF3Br, CH2Br2, Br2CHCHBr2, SOF2, COF2, (CNF)3, ClC(O)(CF2)3C(O)Cl, CF3SO2F, 1,2-dichlorotetrafluorocyclobutene, and perfluorocyclobutene, resp. Compound 5 is readily converted to C6F5OCF2Si(CH3)3 (16) with hexaethylphosphorous triamide and (CH3)3SiCl in benzonitrile. With C6F5CN, CF3C6F5, C5F5N, CH3I, perfluorocyclobutene, CF3C(O)Cl and (CNF)3, 2 forms diethers p-CNC6F4OC6F4OC6F4CN-p (17), p-CF3C6F4OC6F4OC6F4CF3-p (18), NC5F4OC6F4OC5F4N (19), CH3OC6F4OCH3 (20), and C2F5C(O)OC6F4OC(O)C2F5 (21), resp. Reaction of 3 with 1,2-dilodotetrafluorobenzene in diglyme gives 1,4,9-trifluoro-2,3-dilodophenoxine (22). Disiloxane 4 with C6H5CH2Br, CH3I, C6F5CH2Br, and COF2 results in C6H5CH2OC(CF3)2C6F4C(CF3)2OCH2C6H5 (23), CH3OC(CF3)2C6F4C(CF3)2OCH3 (24), C6F5CH2OC(CF3)2C6F4C(CF3)2OCH2C6F5 (25), and [cyclic] C(O)OC(CF3)2C6F4C(CF3)2OC(O)OC(CF3)2C6F4C(CF3)2O (26), resp. These materials are thermally and hydrolytically stable and are formed in high yields.

ACCESSION NUMBER: 1995:269201 CAPIUS  
 DOCUMENT NUMBER: 122:133262  
 TITLE: Per- and Polyfluoroaryl Mono- and Disiloxanes as Transfer Reagents in the Synthesis of Highly Fluorinated Mono- and Diethers  
 AUTHOR(S): Patel, Nimesh R.; Chen, Jianguo; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
 CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 83844-2343, USA  
 SOURCE: Inorganic Chemistry (1995), 34(1), 13-17  
 CODEN: INOCAJ; ISSN: 0020-1669  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 122:133262  
 IT 15053-71-1P 15077-30-2P  
 RL: SYN (Synthetic preparation); PREP (Preparation) (preparation of)  
 RN 15053-71-1 CAPIUS  
 CN Benzonitrile,  
 4,4'-((2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy))bis(2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



RN 15077-30-2 CAPLUS  
CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)



\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I, useful as material for hardeners for fluorinated epoxy resins (no data), is prepared via II [R = cyano, CO2H]. Thus, a mixture of

tetrafluorophthalonitrile, tetrafluorohydroquinone, and Et3N in DMF was heated at 35° for 30 to give 21% II [R = cyano], which was treated with 60% H2SO4 at 150° for 5 h to give 26% II [R = CO2H], which was refluxed with Ac2O for 2 h to 52% I.

ACCESSION NUMBER: 1994:630662 CAPLUS

DOCUMENT NUMBER: 121:230662

TITLE: preparation of a perfluorinated hexacarboxylic acid as

material for hardeners for fluorinated epoxy resins

INVENTOR(S): Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji

PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan

SOURCE: Jpn. Kokai Tokyo Koho, 5 pp.

CODEN: JI00KAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06157501	A2	19940603	JP 1992-340986	19921130
PRIORITY APPLN. INFO.:				
			JP 1992-340986	19921130

OTHER SOURCE(S): CASREACT 121:230662


IT 158394-12-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and conversion into trianhydride)

RN 158394-12-8 CAPLUS

CN 1,2-Benzenedicarboxylic acid, 4,5-bis[4-(3,4-dicarboxy-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)




IT 158394-11-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and hydrolysis of)

RN 158394-11-7 CAPLUS

CN 1,2-Benzenedicarbonitrile, 4,5-bis[4-(3,4-dicyano-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)



IT 158394-13-9P


RL: IMP (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)


(preparation of a perfluorinated hexacarboxylic acid as material)

RN 158394-13-9 CAPLUS

CN 1,3-Isobenzofurandione, 4,7-difluoro-5,6-bis[2,3,5,6-tetrafluoro-4-[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-5-isobenzofuran-2-yl)oxy]phenoxy]- (9CI)

(CA INDEX NAME)





AB The material comprises a perfluoropolyamic acid I or a perfluoropolyimide II [R1 = groups derived from (per)fluorinated aromatic tetracarboxylic acids;

R2 = groups derived from (per)fluorinated aromatic diamines] containing a nonlinear optical material. The manufacturing process employs a DC-field

polling at temperature greater than the glass transition, or an imidization of the

polyamide precursor in a corona-discharge field. The material is suited for use as a high-conversion 2nd harmonic generator for a near IR input.

ACCESSION NUMBER: 1994:521361 CAPLUS

DOCUMENT NUMBER: 121:121361

TITLE: Second harmonic generating materials and manufacture thereof

INVENTOR(S): Amano, Michuki; Kaino, Toshikuni; Sasaki, Shigekuni;

PATENT ASSIGNEE(S): Ando, Shinji; Matsura, Toru

SOURCE: Nippon Telegraph & Telephone, Japan

Jpn. Kokai Tokkyo Koho, 10 pp.

CODEN: JKOQAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06003713	A2	19940114	JP 1991-123054	19910426

PRIORITY APPLN. INFO.: JP 1991-123054 19910426

IT 157044-23-0P 157044-24-1P 157044-25-2P

RL: PREP (Preparation)  
(prepare and use of, as second harmonic generating materials)

RN 157044-23-0 CAPLUS

CN 1-Naphthalenesulfonic acid,  
3-hydroxy-4-[(1-hydroxy-2-naphthalenyl)azo]-7-nitro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine and

5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1

CMF C22 F10 O8


L56 ANSWER 58 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1


CMF C22 F10 O8



CM 2

CRN 25747-08-4

CMF C20 H13 N3 O7 S




CM 3

CRN 20115-19-9

CMF C12 H4 F8 N2 O


L56 ANSWER 58 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



CM 2

CRN 25747-08-4


CMF C20 H13 N3 O7 S



CM 3

CRN 1198-63-6

CMF C6 H4 F4 N2




RN 157044-24-1 CAPLUS

CN 1-Naphthalenesulfonic acid,

3-hydroxy-4-[(1-hydroxy-2-naphthalenyl)azo]-7-nitro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] and

5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

L56 ANSWER 58 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 157044-25-2 CAPLUS

CN 1-Naphthalenesulfonic acid,


3-hydroxy-4-[(1-hydroxy-2-naphthalenyl)azo]-7-nitro-, polymer with 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] and

4,4'-thiobis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1


CMF C22 F10 O8



CM 2

CRN 61907-46-8

CMF C12 H4 F8 N2 S




CM 3


CRN 25747-08-4

CMF C20 H13 N3 O7 S






LS6 ANSWER 59 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Perfluorinated polyimides with high glass temperature (>270°) and high optical transparency over the entire wave lengths of optical communications were prepared. Their high thermal stability and optical transparency were due to the fully aromatic mol. structure and the absence of H atoms. These polymers had low dielec. consts., low refractive indexes, and low birefringence.  
 ACCESSION NUMBER: 1994:484109 CAPLUS  
 DOCUMENT NUMBER: 121:84109  
 TITLE: Synthesis of perfluorinated polyimides for optical applications  
 AUTHOR(S): Ando, Shinji; Matsuura, Tohru; Sasaki, Shigekuni  
 CORPORATE SOURCE: Interdiscipl. Res. Lab., NTT, Musashino, 180, Japan  
 SOURCE: ACS Symposium Series (1994), 537(Polymer for Microelectronics), 304-22  
 CODEN: ACSMCS; ISSN: 0097-6156  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143363-91-1P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and polymerization of, with diamines)  
 RN 143363-91-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro- (9CI) (CA INDEX NAME)





IT 143376-21-0P 143433-45-8P  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (preparation and properties of, for optical applications)  
 RN 143376-21-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 143363-91-1  
 CMF C22 F10 O8



CM 2  
 CRN 1198-63-6  
 CMF C6 H4 F4 N2



RN 143433-45-8 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




AB The title derivs. I (X = O, S; Y = O, S, Q1, Q2) are prepared by treating 28 mol 3,4,5,6-tetrafluorophthalonitrile (II) with 1 mol phenols or thiophenols III in the presence of basic substances in organic solvents.  
 A mixture of 0.22 mol II and KF in MeCN was treated dropwise with a solution of 0.011 mol III (X = Y = O) in MeCN under reflux over 1 h, then refluxed for 4 h to give 99% I.

ACCESSION NUMBER: 1994:435030 CAPLUS  
 DOCUMENT NUMBER: 121:35030  
 TITLE: Preparation of fluorine-containing phthalonitrile derivatives as intermediates for fluorine-containing polyimides  
 INVENTOR(S): Kaieda, Osamu; Okumura, Yasunori; Ito, Hideki  
 PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
 CODEN: JKKKAR  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06016615	A2	19940125	JP 1992-176633	19920703
JP 08000803	B4	19960110		

PRIORITY APPLN. INFO.: JP 1992-176633 19920703


OTHER SOURCE(S): CASREACT 121:35030; MARPAT 121:35030  
 IT 143376-50-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of, as intermediate for polyimides)  
 RN 143376-50-5 CAPLUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-




L56 ANSWER 61 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Refractive indexes at 589.3 nm of 19 polyimides were measured using an Abbe refractometer and calculated from their Van der Waals vols. and mol. polarizabilities. The calculated refractive index when the packing coefficient was assumed to be 0.681 shows a linear relationship with the measured index (n) with a square correlation coefficient of 0.900. However, the slope considerably deviates from 1.0, which indicates that the mol. packing of polyimides changes according to their mol. structure. Packing coeffs. (Kp) of polyimides are estimated by comparing n with calculated parameter  $\phi_0$ . The mol. chains of the polyimides with high n are densely packed, and a planar structure of pyromellitic dianhydrides or ether linkages lead to high Kp. In contrast, the mol. chains of polyimides with low n are loosely packed, and trifluoromethyl groups cause a decrease of interchain interaction and an intra-chain steric hindrance that inhibits mol.

packing.  
ACCESSION NUMBER: 1994:299799 CAPLUS  
DOCUMENT NUMBER: 120:299799  
TITLE: Calculation of refractive indices of polyimides and their molecular packing  
AUTHOR(S): Ando, Shinji  
CORPORATE SOURCE: NTT Interdiscip. Res. Lab., Musashino, 180, Japan  
SOURCE: Kobunshi Ronbunshu (1994), 51(4), 251-7  
CODEN: KBRBA3; ISSN: 0386-2186  
DOCUMENT TYPE: Journal  
LANGUAGE: Japanese  
IT 148362-05-4 148362-06-5 148446-36-0  
148446-37-1  
RL: PRP (Properties)  
(Van der Waals volume and mol. polarizabilities and refractive indexes of, calcn. and measuring of)  
RN 148362-05-4 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenyleneoxy-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148362-06-5 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diyl)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 148446-36-0 CAPLUS

L56 ANSWER 61 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CN 1,3-isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis(4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diamine (9CI) (CA INDEX NAME)

CH 1  
CRN 143363-91-1  
CHF C22 F10 O8




CH 2  
CRN 341-58-2  
CHF C14 H10 F6 N2





RN 148446-37-1 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis(benzenamine) (9CI) (CA INDEX NAME)

CH 1  
CRN 143363-91-1  
CHF C22 F10 O8



CH 2

CRN 101-80-4  
CMF C12 H12 N2 O

AB The title compds. (I; R1, R2 = F, C1-12 linear or branched alkoxy, C1-12 alkylthio, C1-12 alkylamino, aryloxy, arylthio, arylamino, Q - Q2; X = O, S; Y1 = CO2CO), useful as intermediates for drugs, agrochems., optical materials, printed circuit board materials, photosensitive materials, and liquid crystal materials, are prepared by heating F-containing phthalic acid

derivs. (II; R1, R2, X = same as above; Y, Y1 = CO2H) in a solvent selected from SOCl2, POCl3, or AcCl at 40-105°. The process is safe and gives perfluorophthalic anhydride derivs. I of high purity in high yields. Thus, 70.5 g SOCl2 was added to 30.0 g tetrafluorophthalic acid and the resulting mixture was allowed to react at 70° for .apprx.3 h to give 97% tetrafluorophthalic anhydride.

ACCESSION NUMBER: 1994:298457 CAPLUS  
DOCUMENT NUMBER: 120:298457  
TITLE: Preparation of fluorine-containing phthalic anhydride derivatives  
INVENTOR(S): Okumura, Yasunori; Ito, Hideki; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
CODEN: JKOKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE

JP	A2	19940125	JP	1992-175450	19920702
JP 06016656					
JP 3130653	B2	20010131			

PRIORITY APPLN. INFO.: JP 1992-175450 19920702


OTHER SOURCE(S): CASREACT 120:298457; MARPAT 120:298457

IT 143363-92-2

RL: PROC (Process)  
(conversion of, into acid anhydride)

RN 143363-92-2 CAPLUS

CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)





IT 143363-91-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 143363-91-1 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)



AB The liquid crystal orientation film comprises a fluorinated polyimide with a repeating unit I [A = tetravalent aromatic; B = divalent aromatic; ≥1 of A and B is substituted with a F-containing substituent]. The title LCD using said orientation film was also claimed.

ACCESSION NUMBER: 1994:257539 CAPLUS  
DOCUMENT NUMBER: 120:257539  
TITLE: Liquid crystal orientation film providing large pretilt angle and bistability and liquid-crystal display (LCD) device using same

INVENTOR(S): Ishibashi, Shigeki; Hirayama, Misako; Matsura, Tooru  
PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKOKAF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 05150245	A2	19930618	JP 1991-136951	19910514


PRIORITY APPLN. INFO.: JP 1990-129146 19900521

IT 154455-33-1P



RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation and use of, liquid-crystal display orientation film from)  
154455-33-1 CAPLUS  
RN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 3,3'-dimethyl[1,1'-biphenyl]-4,4'-diamine (9CI) (CA INDEX NAME)

CH 1

CRN 143363-91-1  
CMF C22 F10 O8



CH 2

CRN 119-93-7  
CHF C14 H16 N2

AB An optical part comprises a perfluoropolyimide I (R1 = C6Rf4 or directly bonded 2-3 C6Rf4 rings; R2 = C6Rf4 or (in)directly bonded C6Rf4; Rf = F, perfluoroalkyl), wherein the optical part (lenses and prisms) is formed by

cast-molding a polyamide precursor.

ACCESSION NUMBER: 1994:41680 CAPIUS  
DOCUMENT NUMBER: 120:41680  
TITLE: Optical parts and manufacture thereof  
INVENTOR(S): Takeshima, Mikio; Ando, Shinji; Matsura, Tooru; Sasaki, Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 11 pp.  
CODEN: JYOGAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 05113501	A2	19930507	JP 1991-122960	19910426
JP 3050334	B2	20000612		


PRIORITY APPLN. INFO.: JP 1991-122960 19910426

IT 143376-21-0 143376-23-2 143376-24-3


RL: USES (Uses)  
(cast-molded lenses from)

RN 143376-21-0 CAPIUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
CHF C22 F10 O8

CH 2


CRN 1198-63-6  
CHF C6 H4 F4 N2

RN 143376-23-2 CAPIUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
CHF C22 F10 O8

CH 2


CRN 20115-19-9  
CHF C12 H4 F8 N2 O


RN 143376-24-3 CAPIUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-thiobis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

CH 1

CRN 143363-91-1  
CHF C22 F10 O8

CH 2

CRN 61907-46-8  
CHF C12 H4 F8 N2 S



AB The title liquid-crystal display device comprises a liquid crystal layer and a pair of transparent electrode-bearing plastic flexible film substrates, in

which the plastic film is made of a completely fluorinated polyimide film I [R1 = specified benzene-type moiety; R2 = specified moiety including benzene-type moiety; and n = 1-10].

ACCESSION NUMBER: 1994:19348 CAPLUS

DOCUMENT NUMBER: 120:19348

TITLE: Flexible liquid-crystal display device having improved

moisture-proof and optical properties  
INVENTOR(S): Takahashi, Kazuo; Ando, Shinji; Matsura, Toru; Sasaki,

Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKOXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328524	A2	19921117	JP 1991-122939	19910426
PRIORITY APPLN. INFO.:			JP 1991-122939	19910426

IT 143433-45-8 143433-47-0

RL: USES (Uses)  
(transparent substrates from, for flexible liquid-crystal display devices)

RN 143433-45-8 CAPLUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS



CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



AB An IR-transmitting ridge waveguide, formed typically on a SiO<sub>2</sub>-passivated Si wafer, contains I (X1 = II, III, IV; X2 = V, VI, VII; R1 = F, perfluoroalkyl; X3 = O, CO, SO<sub>2</sub>, C(R2)2, S, X4, OX4, X4O, (OX4)<sub>n</sub>, (OX4)<sub>n</sub>O, OCOX4COO; R2 = F, perfluoroalkyl; X4 = perfluoroalkylene; n = 1-10). The waveguide withstands high temperature and is suited for use in a large-area optoelectronic integrated circuit.

ACCESSION NUMBER: 1994:18766 CAPLUS

DOCUMENT NUMBER: 120:18766

TITLE: Perfluoropolyimide waveguide

INVENTOR(S): Matsura, Toru; Ando, Shinji; Sasaki, Shigekuni;

Shimokawa, Fusao

PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.

CODEN: JKOXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328504	A2	19921117	JP 1991-125450	19910426
PRIORITY APPLN. INFO.:			JP 1991-125450	19910426


IT 143376-21-0 143376-23-2 143433-45-8

143433-47-0  
RL: USES (Uses)

L56 ANSWER 66 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 (IR-transmitting ridge waveguides from)  
 RN 143376-21-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)  
 CH 1  
 CRN 143363-91-1  
 CMF C22 F10 O8




CH 2  
 CRN 1198-63-6  
 CMF C6 H4 F4 N2




RN 143376-23-2 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis(2,3,5,6-tetrafluorobenzenamine) (9CI) (CA INDEX NAME)  
 CH 1  
 CRN 143363-91-1  
 CMF C22 F10 O8


L56 ANSWER 66 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



CH 2  
 CRN 20115-19-9  
 CMF C12 H4 F8 N2 O




RN 143433-45-8 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

L56 ANSWER 66 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)


PAGE 1-A




PAGE 1-B



L56 ANSWER 67 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI




AB I was prepared and copolymd. with tetrafluoro-m-phenylenediamine to give an aromatic fluoropolymer-polyether-polyimide (II) with glass temperature 301°, initial decomposition temperature 407° (under N), and dielec. constant 2.8 (10 kHz). No substantial absorption peak was noted for II over the range 0.8-1.6 μm.  
 ACCESSION NUMBER: 1993:603979 CAPLUS  
 DOCUMENT NUMBER: 119:203979  
 TITLE: Synthesis of perfluorinated polyimides for optical applications  
 AUTHOR(S): Ando, S.; Matsuura, T.; Sasaki, S.  
 CORPORATE SOURCE: Interdiscip. Res. Lab., NTT, Musashino, 180, Japan  
 SOURCE: Polymeric Materials Science and Engineering (1992), 66, 200-1  
 CODEN: PMSENG; ISSN: 0743-0515  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143363-91-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and polycondensation of)  
 RN 143363-91-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro- (9CI) (CA INDEX NAME)




IT 143376-21-0P 143433-45-8P

L56 ANSWER 67 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and thermal, dielec. and spectral absorption properties of)  
 RN 143376-21-0 CAPIUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)  
 CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8




CH 2  
 CRN 1198-63-6  
 CHF C6 H4 F4 N2



RN 143433-45-8 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl) (2,4,5,6-tetrafluoro-1,3-phenylene) (4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)

L56 ANSWER 68 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 GI



AB A heat-resistant flexible printed circuit substrate comprises a totally fluorinated polyimide I (R1 = II and R2 = III (Rf = F, perfluoroalkylene, etc.)). The manufacture comprises laminating the substrate with a metal foil

(e.g., Al).  
 ACCESSION NUMBER: 1993:572383 CAPIUS  
 DOCUMENT NUMBER: 119:172383  
 TITLE: Heat-resistant flexible printed circuit substrate and manufacture thereof  
 INVENTOR(S): Matsuura, Toru; Ando, Shinji; Sasaki, Shigekuni  
 PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328891	A2	19921117	JP 1991-125449	19910426
PRIORITY APPLN. INFO.:			JP 1991-125449	19910426

IT 143376-21-0P 143376-23-2P 143433-45-8P  
 143433-47-0P  
 RL: PREP (Preparation)  
 (preparation of, heat-resistant flexible printed circuit substrate from)


RN 143376-21-0 CAPIUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8


L56 ANSWER 67 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)



L56 ANSWER 68 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)




CH 2  
 CRN 1198-63-6  
 CHF C6 H4 F4 N2




RN 143376-23-2 CAPIUS  
 CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8



CH 2



RN 143433-45-8 CAPLUS  
 CN  
 Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 143433-47-0 CAPLUS  
 CN  
 Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)


L56 ANSWER 69 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB The thin films are prepared by evaporation and polymerization of perfluorinated tetracarboxylic acids with perfluorinated diamines on a support. Thus, 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene anhydride and 2,4,5,6-tetrafluoro-1,3-phenylenediamine were evaporated at 80° and 70°, resp. and 1 + 10<sup>-7</sup> torr onto a support to grow a 2000-Å film followed by polymerization at 200° to give a film with dielec. constant ≤2.7 at 10 kHz.  
 ACCESSION NUMBER: 1993:540736 CAPLUS  
 DOCUMENT NUMBER: 119:140736  
 TITLE: Fluorinated polyimide resin thin films with low dielectric constant  
 INVENTOR(S): Maruo, Yoko; Ando, Shinji; Sasaki, Shigekuni; Matsura, Toru  
 PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JIOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328133	A2	19921117	JP 1991-123061	19910426
JP 3080183	B2	20000821		
PRIORITY APPLN. INFO.:			JP 1991-123061	19910426


IT 143376-21-OP 143376-23-2P 143433-45-8P  
 143433-47-OP  
 RL: PREP (Preparation)  
 (thin films, with good adhesion and small dielec. constant, manufacture of)  
 RN 143376-21-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
 CMF C22 F10 O8



PAGE 1-A




PAGE 1-B



CH 2


CRN 1198-63-6  
 CMF C6 H4 F4 N2



RN 143376-23-2 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis(2,3,5,6-tetrafluorobenzeneamine) (9CI) (CA INDEX NAME)


CH 1

CRN 143363-91-1  
 CMF C22 F10 O8



CH 2

CRN 20115-19-9  
 CMF C12 H4 F8 N2 O






RN 143433-45-8 CAPIUS

CN


Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 143433-47-0 CAPIUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



AB The title polyimides are prepared from perfluoro or perfluoroalkyl-substituted aromatic components with a specified structure. Thus, a DMF solution of 1,4-difluoropyromellitic dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer was spread on an Al plate and baked at 70-350° for 4.5 h to give a polyimide film with water absorption ≤0.2%, vs. 2.0 % for a film prepared from pyromellitic anhydride-oxidianiline copolymer.

ACCESSION NUMBER: 1993:519002 CAPIUS

DOCUMENT NUMBER: 119:119002

TITLE: Heat- and moisture-resistant polyimide electric insulators

INVENTOR(S): Sasaki, Shigekuni; Ando, Shinji; Matsura, Toru

PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKOXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04325580	A2	19921113	JP 1991-96622	19910426
PRIORITY APPLN. INFO.:			JP 1991-96622	19910426

IT 143376-21-0 143376-23-2 143433-45-8

143433-47-0

RL: USES (Uses)  
(elec. insulating coatings, with low water absorption)


RN 143376-21-0 CAPIUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1

CMF C22 F10 O8





CM 2

CRN 1198-63-6

CMF C6 H4 F4 N2

PAGE 1-B




RN 143376-23-2 CAPIUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1


CMF C22 F10 O8



CM 2

CRN 20115-19-9


CMF C12 H4 F8 N2 O



RN 143433-45-8 CAPIUS


CN

Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



RN 143433-47-0 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



L56 ANSWER 71 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
AB Precursors are prepared from perfluoro-substituted aromatic tetracarboxylic acid dianhydrides, perfluoro-substituted aromatic diamines, and 2-hydroxyethyl methacrylate (I) which introduces photosensitive groups. Thus, 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride was esterified with I to give a dicarboxylic acid diester, treated with thionyl chloride to give the corresponding acid chloride, and condensed with 2,4,5,6-tetrafluoro-1,3-phenylenediamine to prepare a precursor.

ACCESSION NUMBER: 1993:518573 CAPIUS  
DOCUMENT NUMBER: 119:118573  
TITLE: Photosensitive perfluoro polyimide precursors  
INVENTOR(S): Ichino, Toshihiro; Matsura, Toru; Ando, Shinji; Sasaki, Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
CODEN: JKOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328127	A2	19921117	JP 1991-122946	19910426
JP 2320887	B2	19990719		



PRIORITY APPL. INFO.: JP 1991-122946 19910426

IT 143433-45-8P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer, SRU 143433-47-0P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) ether copolymer, SRU 143433-48-1P, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-bis(2,3,5,6-tetrafluoro-4-aminophenyl) sulfide copolymer, SRU

RL: PREP (Preparation)  
(manufacture of, from photocured methacrylate group-containing polyamic acid esters)


RN 143433-45-8 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)

PAGE 1-B




RN 143433-47-0 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




RN 143433-48-1 CAPLUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)thio(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



IT 143363-91-1, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reaction of, with hydroxyethyl methacrylate and thionyl chloride)  
 RN 143363-91-1 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)

GI



AB The title polymers, having low dielec. constant, and also useful for interlayer insulating films for semiconductors, are prepared by polycondensation of perfluoro aromatic tetracarboxylic acid dianhydrides with perfluoro aromatic diamines. Thus, stirring 20.0 mmol I with 20.0 mmol 2,4,5,6-tetrafluoro-1,3-phenylenediamine in 86 g AcNMe2 at room temperature for 3 days, spin coating this solution on an Al sheet and heating gave a film having dielec. constant 2.6 at 1 kHz, 10% weight loss temperature 501°, and water absorption 0.15% at 23°.

ACCESSION NUMBER: 1993:451450 CAPLUS  
 DOCUMENT NUMBER: 119:51450

TITLE: Heat- and moisture-resistant polyimides for surface protective films for semiconductor devices  
 Matura, Toru; Sasaki, Shigekuni; Ando, Shinji  
 Nippon Telegraph and Telephone Corp., Japan  
 Jpn. Kokai Tokkyo Koho, 7 pp.  
 CODEN: JROKAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328126	A2	19921117	JP 1991-122985	19910426
PRIORITY APPLM. INFO.:			JP 1991-122985	19910426

IT 143376-21-0P 143376-23-2P 143433-45-8P

143433-47-0P 148691-71-8P

RL: PREP (Preparation)

(preparation of, for protective or insulating films for semiconductor devices, heat- and moisture-resistant)



RN 143376-21-0 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1

CRN 143363-91-1


CHF C22 F10 08



CH 2

CRN 1198-63-6

CHF C6 H4 F4 N2




RN 143376-23-2 CAPLUS

CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)


CH 1

CRN 143363-91-1


CHF C22 F10 08




CH 2




RN 143433-45-8 CAPLUS  
 CN  
 Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)




CH 2  
 CRN 46861-94-3  
 CHF C10 F2 O6




CH 3  
 CRN 1198-63-6  
 CHF C6 H4 F4 N2



PAGE 1-A



PAGE 1-B



RN 148691-71-8 CAPLUS  
 CN 1H,3H-Benzo[1,2-c:4,5-c']difuran-1,3,5,7-tetrone, 4,8-difluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine and 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-1,3-isobenzofurandione] (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8


L56 ANSWER 73 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB The title fibers, with good transmittance in near IR light, are comprised of (per)fluorotetracarboxy acid or dianhydride and (per)fluoro diamines. Thus, a 50-μm-diameter fiber prepared from 1:1 copolymer of 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride and 2,4,5,6-tetrafluoro-1,3-phenylenediamine showed no weight loss after 2 h at 500° and light loss 0.1 dB at wave length 1.3 μm.  
 ACCESSION NUMBER: 1993:450901 CAPLUS  
 DOCUMENT NUMBER: 119:50901  
 TITLE: Heat-resistant polyamide-polyimide optical fibers  
 INVENTOR(S): Ando, Shinji; Matsura, Toru; Sasaki, Shigekuni  
 PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
 CODEN: JKOQAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328503	A2	19921117	JP 1991-122988	19910426
JP 2940645	B2	19990825		


PRIORITY APPLN. INFO.: JP 1991-122988 19910426

IT 143376-21-0 143376-23-2 143433-45-8  
 143433-47-0 148780-36-3  
 RL: USES (Uses)  
 (optical fiber, heat-resistant)  
 RN 143376-21-0 CAPLUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1  
 CRN 143363-91-1  
 CHF C22 F10 O8




CH 2  
 CRN 1198-63-6  
 CHF C6 H4 F4 N2



RN 143376-23-2 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8




CM 2

CRN 20115-19-9  
CMF C12 H4 F8 N2 O



RN 143433-45-8 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



RN 148780-36-3 CAPLUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] and 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8




CM 2

CRN 20115-19-9  
CMF C12 H4 F8 N2 O



CM 3

CRN 1198-63-6  
CMF C6 H4 F4 N2



L56 ANSWER 74 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB The title films showing good light transmittance at 0.8-1.7  $\mu\text{m}$  comprise a polyimide layer and a metal layer, where the polyimide layer contains an aromatic polyimide with fluorine- or perfluoroalkyl-substituted phenylene groups. Stirring 11.644 g 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride and 3.602 g 2,4,5,6-tetrafluoro-1,3-phenylenediamine in 86 g ACNMe<sub>2</sub> for 3 days, coating the polyamic solution onto an Al foil, curing at 70°-350° for 4.5 h, and soaking in 10% HCl gave a polyimide film with dielec. const 2.7. Vapor deposition of Ag onto the film gave a composite film.


ACCESSION NUMBER: 1993:450896 CAPLUS  
DOCUMENT NUMBER: 119:50896  
TITLE: Polyimide-metal composite films with low dielectric constant and improved light transmission  
INVENTOR(S): Ando, Shinji; Matsura, Toru; Sasaki, Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp;  
CODEN: JIQQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04325236	A2	19921113	JP 1991-119108	19910424
JP 3001119	B2	20000124		


PRIORITY APPLN. INFO.: JP 1991-119108 19910424

IT 143376-21-0 143376-23-2 143433-45-8  
143433-47-0 148780-36-3  
RL: USES (Uses)  
(metal film laminate, with low dielec. constant and good optical transmission)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)  
CH 1  
CRN 143363-91-1  
CHF C22 F10 O8

L56 ANSWER 74 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




CH 2  
CRN 1198-63-6  
CHF C6 H4 F4 N2



RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CHF C22 F10 O8




CH 2

L56 ANSWER 74 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

CRN 20115-19-9  
CHF C12 H4 F8 N2 O




RN 143433-45-8 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

L56 ANSWER 74 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

PAGE 1-A




PAGE 1-B




RN 148780-36-3 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] and 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CHF C22 F10 O8



CM 2


CRN 20115-19-9  
CMF C12 H4 F8 N2 O

CM 3


CRN 1198-63-6  
CMF C6 H4 F4 N2CRN 1198-63-6  
CMF C6 H4 F4 N2

RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis(2,3,5,6-tetrafluorobenzenamine) (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 O8

CM 2

CRN 20115-19-9  
CMF C12 H4 F8 N2 O

RN 143433-45-8 CAPLUS  
CN

Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-

L56 ANSWER 75 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Films having good light transmittance at wavelength 0.8-1.7  $\mu$ m comprise a layer of a perfluoropolyimide and an ITO layer. Thus, a transparent 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine polyimide film was prepared and sputter-coated to give a film having surface resistance 30  $\Omega$ /box.

ACCESSION NUMBER: 1993:429590 CAPLUS  
DOCUMENT NUMBER: 119:29590  
TITLE: Polyimide-indium tin oxide electrically conductive films and their manufacture  
INVENTOR(S): Ando, Shinji; Matsura, Toru; Sasaki, Shigekuni  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JKOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04323232	A2	19921112	JP 1991-119109	19910424
JP 3105571	B2	20001106		

PRIORITY APPLN. INFO.: JP 1991-119109 19910424

IT 143376-21-0, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer 143376-23-2 143433-45-8, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer, SRU 143433-47-0  
RL: USES (Uses)


(films, ITO-coated, transparent and elec. conductive)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 O8

CM 2


L56 ANSWER 75 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
tetrafluoro-1,3-phenylene) (4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy (9CI) (CA INDEX NAME)





RN 143433-47-0 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



AB The title polyimides suitable for optical materials for optoelectronic integrated circuits contain repeating units I (R = divalent organic group containing 21 H). 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride was polymerized with 2,2'-bis(trifluoromethyl)-4,4'-diaminobiphenyl in Ac<sub>2</sub>Ome<sub>2</sub>, and the resulting polyamic acid solution was coated on a Si wafer and subjected to thermal imidation to give a polyimide film with very low absorption of light in the 0.8-2.0 μm wavelength range.


ACCESSION NUMBER: 1993:428840 CAPIUS  
DOCUMENT NUMBER: 119:28840  
TITLE: Fluorine-containing polyamic acids, fluorine-containing polyimides, and manufacture thereof  
INVENTOR(S): Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokyo Koho, 7 pp.  
CODEN: JJOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04314732	A2	19921105	JP 1991-106484	19910412
JP 2827058	B2	19981118		


PRIORITY APPLN. INFO.: JP 1991-106484 19910412

IT 148362-03-2P 148362-04-3P 148362-05-4P  
148362-06-5P 148362-07-6P 148446-36-0P  
148446-37-1P 148446-38-2P 148446-39-3P  
148446-40-4P 148446-41-5P 148446-42-6P  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(manufacture of, for optical materials for optoelectronic circuits)  
RN 148362-03-2 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 148362-04-3 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B




RN 148362-05-4 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)-1,4-phenylene[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]-1,4-phenylene] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B






RN 148362-06-5 CAPIUS

CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diyl]] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B

RN 148383-89-5 CAPIUS


CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-

diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)[2-(trifluoromethyl)-1,4-phenylene]oxy[1,1'-biphenyl]-4,4'-diyl]oxy[3-(trifluoromethyl)-1,4-phenylene]] (9CI) (CA INDEX NAME)

CM 2

CRN 341-58-2

CMF C14 H10 F6 N2




RN 148446-37-1 CAPIUS

CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis(benzenamine)] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1


CMF C22 F10 O8



CM 2

CRN 101-80-4

CMF C12 H12 N2 O




RN 148446-38-2 CAPIUS


CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-methylenebis(benzenamine)] (9CI) (CA INDEX NAME)

CM 1

PAGE 1-A



PAGE 1-B




RN 148446-36-0 CAPIUS

CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)[1,1'-biphenyl]-4,4'-diamine] (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1

CMF C22 F10 O8



CRN 143363-91-1


CMF C22 F10 O8



CM 2

CRN 101-77-9

CMF C13 H14 N2




RN 148446-39-3 CAPIUS

CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[(1,1'-biphenyl)-4,4'-diyl]bis(oxy)]bis[2-(trifluoromethyl)benzenamine] (9CI) (CA INDEX NAME)

CM 1


CRN 143363-91-1

CMF C22 F10 O8




CM 2

CRN 94517-23-4  
CHF C26 H18 F6 N2 O2




RN 148446-40-6 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[benzenamine] (9CI) (CA INDEX NAME)

CH 1  
CRN 143363-91-1  
CHF C22 F10 O8




CH 2  
CRN 1095-78-9  
CHF C15 H12 F6 N2




RN 148446-41-7 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diamine and 5,5'-[2,2,2-trifluoro-


CH 1  
CRN 143363-91-1  
CHF C22 F10 O8



CH 2  
CRN 341-58-2  
CHF C14 H10 F6 N2




CH 3  
CRN 101-80-4  
CHF C12 H12 N2 O




1-(trifluoromethyl)ethylidene]bis[1,3-isobenzofurandione] (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CHF C22 F10 O8



CH 2  
CRN 1107-00-2  
CHF C19 H6 F6 O6



CH 3  
CRN 341-58-2  
CHF C14 H10 F6 N2



RN 148446-42-8 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,2'-bis(trifluoromethyl)(1,1'-biphenyl)-4,4'-diamine and 4,4'-

AB The filter, comprising a dielec. multilayer, is characterized in that the substrate consists of a thin-film perfluoropolyimide or a blend or a copolymer thereof.


ACCESSION NUMBER: 1993:417687 CAPLUS  
DOCUMENT NUMBER: 119:17687  
TITLE: IR longpass dielectric interference filter  
INVENTOR(S): Yamamoto, Fumio; Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji; Oguchi, Taisuke  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JKOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328502	A2	19921117	JP 1991-125451	19910426

PRIORITY APPLN. INFO.: JP 1991-125451 19910426


IT 143433-45-8 148273-08-9  
RL: USES (Uses)  
(fluorinated polyimide blends containing, IR longpass filters containing)

RN 143433-45-8 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)





RN 148273-08-9 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1  
CRN 143363-92-2  
CHF C22 H4 F10 O10




CM 2


CRN 1198-63-6  
CMF C6 H4 F4 N2CRN 1198-64-7  
CMF C6 H4 F4 N2

RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)

CM 1

CRN 143363-91-1  
CMF C22 F10 O8

CM 2

CRN 20115-19-9  
CMF C12 H4 F8 N2 O


RN 143433-46-9 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-

L56 ANSWER 78 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Polyamic acids are prepared from perfluoroarom. carboxylic acid dianhydrides and perfluoroarom. diamines, mixed with fluoropolymers, and thermally cyclized. Thus, 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene dianhydride-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer (polyamic acid) was prepared, mixed (4 g) with cyclohexanone, Lublon 0.4, and Fluorad 431 (a surfactant) 0.1 g, spin-coated on Al, and imidized to give a film having dielec. constant <2.5 at 1 kHz.  
ACCESSION NUMBER: 1993:410132 CAPLUS  
DOCUMENT NUMBER: 119:10132  
TITLE: Fluoropolymer-containing polyamic acids and polyimides  
and films therefrom with low dielectric constants  
INVENTOR(S): Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JIQQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328161	A2	19921117	JP 1991-122850	19910426

PRIORITY APPLN. INFO.:  
IT 143376-22-1 143376-23-2 143433-46-9  
143433-47-0  
RL: USES (Uses)  
(films, containing PTFE, with low dielec. constant)  
RN 143376-22-1 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,3,5,6-tetrafluoro-1,4-benzenediamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
CMF C22 F10 O8


CM 2

tetrafluoro-1,4-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)  
[Chemical structure of the repeating unit of the polyamic acid, showing a 2,3,5,6-tetrafluoro-1,4-phenylene ring connected via ether linkages to a 4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl ring, which is further connected to another 2,3,5,6-tetrafluoro-1,4-phenylene ring via ether linkages. The structure is enclosed in brackets with a subscript 'n'.]  
RN 143433-47-0 CAPLUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




L56 ANSWER 79 OF 89 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB A perfluorinated polyimide that has glass temperature >300° and high optical transparency over the entire optical communication wavelengths was synthesized. The high thermal stability and optical transparency in near-IR region are due to the fully aromatic mol. structure and the absence of C-H bonds. The use of a diamine with relatively high reactivity and a new perfluorinated dianhydride, which has flexible structure, makes it possible to obtain a tough and flexible perfluorinated polyimide film.

In addition, this polymer has a low dielec. constant Perfluorinated polyimides are promising for use as optical-electronic materials.  
 ACCESSION NUMBER: 1992:592808 CAPIUS  
 DOCUMENT NUMBER: 117:192808  
 TITLE: Perfluorinated polyimide synthesis  
 AUTHOR(S): Ando, Shinji; Matsuura, Tohru; Sasaki, Shigekuni  
 CORPORATE SOURCE: Interdiscip. Res. Lab., NTT, Musashino, 180, Japan  
 SOURCE: Macromolecules (1992), 25(21), 5858-60  
 CODEN: MAMOBX; ISSN: 0024-9297  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143376-21-0P 143433-45-8P  
 RL: PEP (Physical, engineering or chemical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)  
 (preparation and characterization of, for optical materials)  
 RN 143376-21-0 CAPIUS  
 CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)


CM 1

CRN 143363-91-1  
 CHF C22 F10 O8




CM 2


CRN 1198-63-6  
 CHF C6 H4 F4 N2



RN 143433-45-8 CAPIUS  
 CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,4,5,6-tetrafluoro-1,3-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)



IT 143376-50-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and hydrolysis of)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143363-91-1P  
 RL: PEP (Physical, engineering or chemical process); SPN (Synthetic preparation); PREP (Preparation); PROC (Process)  
 (preparation and polymerization of)

L56 ANSWER 80 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB The title polymers, useful as waveguides in optoelectronic integrated circuits or optoelec. boards, are prepared from perfluoro tetracarboxylic acid deriva. and perfluoro diamines, the preparation of which is described.

Polymerization of 4,4'-(tetrafluoro-p-phenylene)bis(3,5,6-trifluorophthalic anhydride) with tetrafluoro-1,3-phenylenediamine gave a polymer with no substantial absorption at optical communication wavelengths.

ACCESSION NUMBER: 1992:531752 CAPLUS

DOCUMENT NUMBER: 117:131752

TITLE: Monomers for preparation of perfluorinated polyamic acids and polyimides

INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni;

SOURCE: Yamasato, Fumio

PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan

EUR. Pat. Appl., 39 pp.

CODEN: EPOXDW

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 480266	A2	19920415	EP 1991-116553	19910927
EP 480266	A3	19930901		
EP 480266	B1	19960626		
R: DE, FR, GB				
JP 05001148	A2	19930108	JP 1991-235020	19910913
JP 2851019	B2	19990127		
JP 09031014	A2	19970204	JP 1996-207475	19910913
CA 2052368	AA	19920329	CA 1991-2052368	19910926
CA 2052368	C	19980915		
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		
PRIORITY APPLN. INFO.:				
			JP 1990-256843	A 19900928
			JP 1991-106552	A 19910412
			JP 1991-106554	A 19910412
			JP 1991-106557	A 19910412
			JP 1991-235020	A3 19910913
			CA 1991-2052368	A3 19910926
			US 1991-765672	A3 19910926
			US 1993-54973	B2 19930430
			US 1993-140482	A3 19931025
			US 1995-451465	B1 19950526
			US 1996-718208	A3 19960920

L56 ANSWER 80 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
US 1998-20573 A3 19980128

IT 143376-50-5P  
RL: RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and hydrolysis of)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




IT 143363-91-1P 143363-92-2P  
RL: PREP (Preparation)  
(preparation of)  
RN 143363-91-1 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro- (9CI) (CA INDEX NAME)



RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)


L56 ANSWER 80 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



IT 143376-21-0P 143376-22-1P 143376-23-2P  
143376-24-3P 143433-45-8P 143433-46-9P  
143433-47-0P 143433-48-1P  
RL: PREP (Preparation)  
(preparation of, for use in optical communications)  
RN 143376-21-0 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CH 1


CRN 143363-91-1  
CMF C22 F10 O8



CH 2

CRN 1198-63-6  
CMF C6 H4 F4 N2


L56 ANSWER 80 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 143376-22-1 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 2,3,5,6-tetrafluoro-1,4-benzenediamine (9CI) (CA INDEX NAME)


CH 1

CRN 143363-91-1  
CMF C22 F10 O8




CH 2

CRN 1198-64-7  
CMF C6 H4 F4 N2




RN 143376-23-2 CAPLUS  
CN 1,3-Isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[4,6,7-trifluoro-, polymer with 4,4'-oxybis[2,3,5,6-tetrafluorobenzeneamine] (9CI) (CA INDEX NAME)

CH 1  
CRN 143363-91-1  
CHF C22 F10 O8




CH 2  
CRN 20115-19-9  
CHF C12 H4 F8 N2 O




RN 143376-24-3 CAPIUS  
CN 1,3-isobenzofurandione, 5,5'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis(4,6,7-trifluoro-, polymer with 4,4'-thiobis[2,3,5,6-tetrafluorobenzenamine] (9CI) (CA INDEX NAME)


CH 1  
CRN 143363-91-1  
CHF C22 F10 O8




CH 2  
CRN 61907-46-8  
CHF C12 H4 F8 N2 S



RN 143433-45-8 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-46-9 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy] (9CI) (CA INDEX NAME)




RN 143433-47-0 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A






PAGE 1-B



RN 143433-48-1 CAPIUS  
CN Poly[(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-2,5-diyl)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)oxy(4,6,7-trifluoro-1,3-dihydro-1,3-dioxo-2H-isoindole-5,2-diyl)(2,3,5,6-tetrafluoro-1,4-phenylene)thio(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A





I

AB The compds. have the general formula I (each R1 = alkyl, alkoxyalkyl, acyloxyalkyl, (un)substituted Ph, cycloalkyl, benzyl; 21 of R2 = alkoxy, alkoxyalkoxy, acyloxyalkoxy, hydroxyalkoxy, aminoalkoxy, cyanoalkoxy, alkoxycarbonylalkoxy, haloalkoxy, (un)substituted phenoxy, cycloalkoxy, benzyloxy, the remaining R2 being F; M = Zn, Cu, Pb, VO, TiO, TixZ, SnX2, AlX, InX; X = halogen) and are prepared by reacting I (R2 = F) with alkoxide, alc, or phenol derivs. in an organic solvent in the optional presence of alkali. I (R1 = Bu; R2 = F; M = VO) was treated with NaOEt in DMF under reflux for 2 h to give 1 (R1 = Bu; R2 = EtO; M = VO) in 75.1% yield,  $\lambda_{\text{max}}$  (DMF) 760 nm.

ACCESSION NUMBER: 1992:428767 CAPLUS

DOCUMENT NUMBER: 117:28767

TITLE: Phthalocyanines, their manufacture, and use as near-IR

absorbers  
INVENTOR(S): Yoshitoshi, Koji; Kaleda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKOGAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

IT 142111-01-1P

RL: PREP (Preparation)  
(manufacture of near-IR-absorbing)

RN 142111-01-1 CAPLUS

CN Titanium, (2,3,9,10,16,17,23,24-octakis(butylthio)-1,4,8,11,15,18,22,25-

octakis(2-fluorophenoxy)-29H,31H-phthalocyaninato(2-)-N29,N30,N31,N32)oxo-  
, (SP-5-12)- (9CI) (CA INDEX NAME)


L56 ANSWER 82 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN

AB The title polymers, e.g., polyether-polyketones, polyarylates, polyimides, and polyesters, have good heat resistance and low attenuation and are useful as optical transmitting systems, e.g., for controlling ignition timing and fuel metering systems for internal combustion engines in automobiles. Thus, an optical fiber comprised a core of amorphous PEEK and a sheath of poly(2,2,2-trifluoroethyl methacrylate).

ACCESSION NUMBER: 1992:130739 CAPLUS  
DOCUMENT NUMBER: 116:130739  
TITLE: Amorphous polymers for optical transmitting systems and optical members and their use  
INVENTOR(S): Takezawa, Yoshitaka; Ohara, Shuichi; Tanno, Seikichi; Takekuni, Noriaki; Shimura, Masato  
PATENT ASSIGNEE(S): Hitachi, Ltd., Japan  
SOURCE: Eur. Pat. Appl., 31 pp.  
CODEN: EPMOXM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 454165	A2	19911030	EP 1991-106851	19910426
EP 454165	A3	19930120		
R: DE, FR, GB, IT, NL				
JP 04009805	A2	19920114	JP 1990-112511	19900427
JP 2854669	B2	19990203		
US 5093888	A	19920303	US 1991-686997	19910418
PRIORITY APPL. INFO.:			JP 1990-112511	A 19900427

IT 138687-03-3  
RL: USES (Uses)  
(optical fibers, heat-resistant, for engine control systems)  
RN 138687-03-3 CAPLUS  
CN Poly(oxy(2,6-dimethyl-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)) (9CI) (CA INDEX NAME)




L56 ANSWER 83 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN


(Continued)



RN 93207-48-8 CAPLUS  
CN Cyclopropanecarboxylic acid, 3-(2,2-dichloroethenyl)-2,2-dimethyl-, cyano[4-fluoro-3-[4-(4-fluorophenoxy)phenoxy]phenyl]methyl ester (9CI) (CA INDEX NAME)



L56 ANSWER 83 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
GI



AB Benzyl cyclopropanecarboxylates I (R = H, F, Cl, Br; R1 = H, cyano, C.tpbond.CH; R2 = H, Me; R3 = CH:CCl2, CHBrCBrCl2, CHBrCBr3, Me), having insecticidal and acaricidal activities at 500 ppm with low fish toxicity, were prepared e.g. from II. Thus, 6.0 mmol NaCN and 0.88 mmol PhCH2NEt3Cl


in H2O was treated dropwise with 4.0 mmol II and 4.2 mmol di-cis,trans-2,2-dimethyl-3-(2,2-dichlorovinyl)cyclopropanecarboxyl chloride in PhMe to give 1.93 g I (R = R2 = H, R1 = cyano, R3 = CH:CCl2).

ACCESSION NUMBER: 1984:610760 CAPLUS  
DOCUMENT NUMBER: 101:210760  
TITLE: Insecticidal cyclopropanecarboxylate esters  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKXKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59134749	A2	19840802	JP 1983-8536	19830120
PRIORITY APPL. INFO.:			JP 1983-8536	19830120

IT 90928-31-7P 93207-48-8P  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation and insecticidal activity of)  
RN 90928-31-7 CAPLUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2,2-tetrabromoethyl)-, 1-[4-fluoro-3-[4-(4-fluorophenoxy)phenoxy]phenyl]-2-propynyl ester (9CI) (CA INDEX NAME)

L56 ANSWER 84 OF 89 CAPLUS COPYRIGHT 2005 ACS on STN  
GI




AB Twelve title esters (I; R = H, halo; R1 = H, F; R2 = H, cyano, HC.tpbond.C, R3 = haloethyl, haloethyl), effective insecticides and miticides at 500 ppm, were prepared. Thus, 5.0 mmol II and 5.25 mmol (1)-III in MePh were added to a solution of 7.5 mmol NaCN and 1.1 mmol PhCH2NEt3Cl in H2O at room temperature to give 2.07 g (2)-I (R = R1 = H; R2 = cyano, R3 = Cl2C:CH).

ACCESSION NUMBER: 1984:438168 CAPLUS  
DOCUMENT NUMBER: 101:38168  
TITLE: Cyclopropanecarboxylate esters as insecticides and miticides with low toxicity  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
CODEN: JKXKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59042342	A2	19840308	JP 1982-153029	19820901
PRIORITY APPL. INFO.:			JP 1982-153029	19820901

IT 90928-31-7P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)  
RN 90928-31-7 CAPLUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2,2-tetrabromoethyl)-, 1-[4-fluoro-3-[4-(4-fluorophenoxy)phenoxy]phenyl]-2-propynyl ester (9CI) (CA INDEX NAME)





GI For diagram(s), see printed CA Issue.

AB The PbO<sub>2</sub> oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl

ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).

ACCESSION NUMBER: 1974:520135 CAPIUS

DOCUMENT NUMBER: 81:120135

TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol

AUTHOR(S): Denivelle, Leon; Huynh Anh Hoa

CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conservatoire Natl. Arts

SOURCE: Metiers, Paris, Fr.

Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 487-90

CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal


LANGUAGE: French

IT 53279-71-3P 53359-93-6P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 53279-71-3 CAPIUS

CN 1,4-Benzenediol, 2,3,5,6-tetrakis(pentafluorophenoxy)- (9CI) (CA INDEX NAME)



RN 53359-93-6 CAPIUS

CN Phenol, 2,3,5,6-tetrafluoro-4-[(2,3,5,6-tetrafluoro-4-(pentafluorophenoxy)phenoxy)- (9CI) (CA INDEX NAME)

<-----User Break----->

GI For diagram(s), see printed CA Issue.

AB The controlled oxidation of perhalo p-bromophenols (I), (II), and (III) gives

the corresponding perhalo polyethers (IV); the perchloro 4-aryloxycyclohexa-2,5-dienone (V) is obtained from pentachlorophenol (VI). Similarly, VII gives VIII and 4-chloro-2,3,5,6-tetrabromophenol gives IX.

ACCESSION NUMBER: 1971:111698 CAPIUS

DOCUMENT NUMBER: 74:111698

TITLE: Oxidation of pentahalobenzenic phenols

AUTHOR(S): Denivelle, Leon; Lampel, Alfred

CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conserv. Natl. Arts

Metiers, Paris, Fr.

SOURCE: Comptes Rendus des Seances de l'Academie des

Sciences,

Serie C: Sciences Chimiques (1971), 272(7), 653-6

CODEN: CHDCAQ; ISSN: 0567-6541

DOCUMENT TYPE: Journal

LANGUAGE: French


IT 31404-37-2P 31404-38-3P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 31404-37-2 CAPIUS

CN 2,5-Cyclohexadien-1-one, 4-bromo-4-[4-(4-bromo-2,3,5,6-

tetrafluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          430.68      4422.25

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          -63.51      -329.23
```

FILE 'REGISTRY' ENTERED AT 20:32:56 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
 DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
 information enter HELP PROP at an arrow prompt in the file or refer  
 to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          0.86      4423.11

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          0.00      -329.23
```

FILE 'REGISTRY' ENTERED AT 20:34:08 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
 DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004


Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10718532.str



chain nodes :

1 13 23 25 26 28 29 30 31 32 33 35

ring nodes :

2 3 4 5 6 7 8 9 10 11 12 16 17 18 19 20 21 22

chain bonds :

1-4 8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31  
 22-35

ring bonds :

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
 18-19 19-20 20-21 21-22

exact/norm bonds :

8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31

exact bonds :

1-4 22-35

normalized bonds :

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
 18-19 19-20 20-21 21-22

G1:N,X

G2:X,Ak,H


Match level :

1:CLASS 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
 11:Atom 12:Atom 13:CLASS 14:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom  
 21:Atom 22:Atom 23:CLASS 25:CLASS 26:CLASS 28:CLASS 29:CLASS 30:CLASS  
 31:CLASS 32:CLASS 33:CLASS 35:CLASS

L57 STRUCTURE UPLOADED

=> d query

L57 STR



G1 N,X  
G2 X,Ak,H

Structure attributes must be viewed using STN Express query preparation.

=> s l57

SAMPLE SEARCH INITIATED 20:34:27 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 110 TO ITERATE

100.0% PROCESSED 110 ITERATIONS 3 ANSWERS  
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 1571 TO 2829  
PROJECTED ANSWERS: 3 TO 163

L58 3 SEA SSS SAM L57

=> s l57 full

FULL SEARCH INITIATED 20:34:32 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 2122 TO ITERATE

100.0% PROCESSED 2122 ITERATIONS 29 ANSWERS  
SEARCH TIME: 00.00.01

L59 29 SEA SSS FUL L57

=> fil caplus

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	161.33	4584.44

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-329.23

FILE 'CAPLUS' ENTERED AT 20:34:35 ON 12 JAN 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.


FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 159

L60            32 L59

=> d l60 1-32 abs ibib hitstr



AB This invention pertains to a method for producing bis(fluorine-containing phthalic anhydride) derivs. I [wherein m = 1-3; n = 1-3; Z = a single bond or a spacer], which comprises reacting a fluorine-containing dinitrile with a fluorine-containing hydroquinone, followed by hydrolysis and dehydration reaction. For example, 3,4,5,6-tetrafluorophthalonitrile was reacted with tetrafluorohydroquinone in Me iso-Bu ketone in the presence of KF to give the bisphthalonitrile (97%). The bisphthalonitrile was hydrolyzed in propionic acid in the presence of H2SO4 to afford the bis(phthalic acid) (75%). The bis(phthalic acid) was treated with acetic anhydride to provide the bis(phthalic anhydride) (98%).

ACCESSION NUMBER: 2004:1127358 CAPIUS  
TITLE: Process for preparation of bis(fluorine-containing phthalic anhydride)  
INVENTOR(S): Masuda, Go; Okumura, Yasunori; Nishimae, Shinji; Nareya, Toshihiro  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: PCT Int. Appl., 39 pp.  
CODEN: PIXKD2  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004111026	A1	20041223	WO 2004-JP8829	20040617
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, ID, IL, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZH, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SE, SZ, TZ, UG, ZH, ZW, AM, AZ, BY, BG, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			JP 2003-172046	A 20030617
			JP 2003-201123	A 20030724

IT 143363-92-2P 143376-50-5P  
RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic)

AB The compns. containing repeating units of AD(R) [A = F-substituted (O- or S-containing) hydrocarbon; D = F-substituted (O- or S-containing) trivalent hydrocarbon group; R = Si(OR)<sub>1</sub>(OR)<sub>2</sub>R<sub>3</sub>; R<sub>1</sub>, R<sub>2</sub> = hydrocarbyl (each b.p. of R<sub>1</sub>OH and R<sub>2</sub>OH under normal pressure ≤250°); R<sub>3</sub> = (F-substituted) double bond-terminated hydrocarbyl] show n and dielec. constant of their cured products 1.350-1.600 and 2.00-4.00, resp. The compns. give cured products with adjustable n and dielec. constant, and improved mech. properties and solvent resistance.

ACCESSION NUMBER: 2004:411677 CAPIUS  
DOCUMENT NUMBER: 140:431126  
TITLE: Manufacture of silicon-containing curable polymer compositions for planar optical waveguides and wiring boards  
INVENTOR(S): Florence, Corey Nawarage  
PATENT ASSIGNEE(S): Fujitsu Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
CODEN: JPOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004143280	A2	20040520	JP 2002-309280	20021024
PRIORITY APPLN. INFO.:			JP 2002-309280	20021024

IT 691906-05-SDP, reaction products with allyltrimethoxysilane and ethanol  
RL: DEV (Device component use); IMF (Industrial manufacture); PREP (Preparation); USES (Uses)  
(crosslinked; manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)


RN 691906-05-5 CAPIUS  
CN 1,3-benzenedicarbonyl dichloride, 5-bromo-, polymer with 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)

CN 1

CRN 691906-04-4  
CMF C20 R4 C12 F8 O4




RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT


CN 2

CRN 83558-87-6  
CMF C15 H12 F6 N2 O2




CN 3


CRN 57863-69-1  
CMF C8 H3 Br C12 O2




IT 691906-02-2P 691906-03-3P 691906-04-4P  
691906-05-5P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation);  
RACT (Reactant or reagent)  
(manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)  
RN 691906-02-2 CAPIUS  
CN Benzonitrile, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



L60 ANSWER 2 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 RN 691906-03-3 CAPLUS  
 CN Benzoic acid, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI)  
 (CA INDEX NAME)




RN 691906-04-4 CAPLUS  
 CN Benzoyl chloride, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI)  
 (CA INDEX NAME)



RN 691906-05-5 CAPLUS  
 CN 1,3-Benzenedicarbonyl dichloride, 5-bromo-, polymer with  
 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and  
 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol]  
 (9CI) (CA INDEX NAME)

CM 1

CRN 691906-04-4  
 CMF C20 H4 Cl2 F8 O4



CM 2


L60 ANSWER 2 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

CRN 83558-87-6  
 CMF C15 H12 F6 N2 O2




CM 3

CRN 57863-69-1  
 CMF C8 H3 Br Cl2 O2



L60 ANSWER 3 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI



AB The title phthalic acid derivs. (Markush structure given) are prepared by  
 reaction of tetrafluorophthalic acid derivs. with MAM (A = divalent  
 organic  
 moiety; M = H, etc.). Thus, I was prepared I is a raw material for the  
 manufacture of fluorinated polyimides. The title phthalic acid derivs.  
 are

useful as intermediates for optical materials, liquid crystals, etc.

ACCESSION NUMBER: 2003:823334 CAPLUS

DOCUMENT NUMBER: 139:330642

TITLE: Phthalic acid derivatives as intermediates for  
 halogenated polyimides, liquid crystals, etc., and  
 process for manufacturing them

INVENTOR(S): Okumura, Yasunori; Kuwahara, Masayoshi; Masuda,  
 Takeshi

PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKKOAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003300934	A2	20031021	JP 2002-108095	20020410
PRIORITY APPLN. INFO.:				
			JP 2002-108095	20020410

OTHER SOURCE(S): MARPAT 139:330642


IT 615263-86-0P  
 RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP  
 (Preparation)  
 (phthalic acid derivs. as intermediates for halogenated polyimides,  
 optical materials, liquid crystals, and process for manufacturing  
 them)


RN 615263-86-0 CAPLUS

CN 1,2-Benzenedicarboxylic acid, 4,4'-[2,3,5,6-tetrafluoro-1,4-  
 phenylene]bis(oxy)]bis[3,5,6-trifluoro-, tetraethyl ester (9CI) (CA

INDEX  
 NAME)

L60 ANSWER 3 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)







AB The compds. are prepared by hydrolysis of aromatic tetranitriles I (X, X' = halo; n, n' = 1-3; Z = single bond, O, S, O-p-C6H4-mYmO, etc; Y = halo; m = 1-4) in the presence of acids in organic solvents. 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene was treated with H2SO4 in propionic acid under reflux for 6 h to give 95.7% 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene.

ACCESSION NUMBER: 2002:886125 CAPIUS  
DOCUMENT NUMBER: 137:384650  
TITLE: Preparation of halogen-containing aromatic carboxylic acids  
INVENTOR(S): Kuwahara, Masayoshi; Yokoo, Junko; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKOQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002332253	A2	20021122	JP 2001-142027	20010511
JP 3490407	B2	20040126		
PRIORITY APPLN. INFO.:			JP 2001-142027	20010511

OTHER SOURCE(S): MARPAT 137:384650  
IT 143363-92-2P 474805-31-7P  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143363-92-2 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]


L60 ANSWER 4 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
RN 474805-29-3 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



RN 474805-31-7 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



IT 143376-50-5, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



IT 474805-29-3P, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrachlorobenzene  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)


L60 ANSWER 5 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN  
AB The present invention relates to halogen-containing aromatic compounds and methods thereof. The present invention relates to a halogen-containing aromatic dianhydride, halogen-containing aromatic tetranitrile compound, halogen-containing m-phenylenediamine compound and fluorine compound, and a method thereof. Tetrafluorophthalonitrile was reacted with tetrachlorohydroquinone in the presence of potassium fluoride and acetonitrile to give 1,4-bis(3,4-dicyanotrifluorophenoxy)tetrachlorobenzene.

ACCESSION NUMBER: 2002:866679 CAPIUS  
DOCUMENT NUMBER: 137:354701  
TITLE: Halogen-containing aromatic compound  
INVENTOR(S): Kuwahara, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan; Nippon Catalytic Chem. Ind.  
SOURCE: Eur. Pat. Appl., 38 pp.  
CODEN: EPXKDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1256564	A1	20021113	EP 2002-253088	20020501
EP 1256564	B1	20040728		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2002332268	A2	20021122	JP 2001-142028	20010511
JP 2002332281	A2	20021122	JP 2001-142029	20010511
JP 2002332264	A2	20021122	JP 2001-142031	20010511
JP 3563040	B2	20040908		
JP 2002332254	A2	20021122	JP 2001-142032	20010511
US 2003018204	A1	20030123	US 2002-133158	20020426
EP 1462436	A1	20040929	EP 2004-10371	20020501
R: DE, FR, GB				
CN 1385427	A	20021218	CN 2002-119176	20020513
PRIORITY APPLN. INFO.:			JP 2001-142028	A 20010511
			JP 2001-142029	A 20010511
			JP 2001-142031	A 20010511
			JP 2001-142032	A 20010511
			EP 2002-253088	A3 20020501

OTHER SOURCE(S): MARPAT 137:354701  
IT 474805-29-3P 474805-31-7P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
(halogen-containing aromatic compound)  
RN 474805-29-3 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]





RN 474805-31-7 CAPLUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-bis-[(2,3,5,6-tetrachloro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-1,3-phenylenediamine copolymer (9CI) (CA INDEX NAME)



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

AB The repeating units of polyimides for coating materials contain tetravalent aromatic organic groups and aromatic groups having no CH bonds. Thus,

a film of 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer had dielec. constant 2.6 at 1 MHz, compared with 3.5 for a com. polyimide.

ACCESSION NUMBER: 2002:139034 CAPLUS  
 DOCUMENT NUMBER: 136:185465  
 TITLE: Insulating coating materials for coil wire  
 INVENTOR(S): Tajiri, Kozo; Kuwahara, Masayoshi; Okumura, Yasunori  
 PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 16 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002056720	AZ	20020222	JP 2000-243309	20000810
PRIORITY APPL. INFO.:			JP 2000-243309	20000810


IT 148273-08-9, 1,4-Bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene-2,4,5,6-tetrafluoro-1,3-phenylenediamine copolymer  
 RL: PRP (Properties); TEM (Technical or engineered material use); USES (Uses)

(fluoro polyimide insulating coating materials for coil wire)

RN 148273-08-9 CAPLUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-bis-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-1,3-phenylenediamine copolymer (9CI) (CA INDEX NAME)


CH 1

CRN 143363-92-2  
 CHF C22 H4 F10 O10



CH 2

CRN 1198-63-6  
 CHF C6 H4 F4 N2



AB The comds. QCO-p-C6H4O-p-C6H4R (Q = 2,3,4,5,6-pentafluorophenyl; R = OH, COQ) are manufactured. Thus, 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone (prepared from 2,3,4,5,6-pentafluoro-4'-methoxybenzophenone) was

polymerized in dimethylacetamide at 160° to give a polymer showing good solubility in dimethylacetamide and m-cresol, 10% weight loss (in air) temperature 421°.

ACCESSION NUMBER: 2001:176773 CAPLUS  
 DOCUMENT NUMBER: 134:208664  
 TITLE: (2,3,4,5,6-pentafluorobenzoyl)diphenyl ethers and fluorine-containing aryl ether ketone polymers having high solubility and heat stability  
 INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Casiday, Rachel E.; Fitch, John W., III; Reddy, V. Sreenivasulu  
 PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
 CODEN: JKOXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001064226	AZ	20010313	JP 1999-183950	19990629
PRIORITY APPL. INFO.:			US 1998-106270	A 19980629
			JP 1999-180091	A 19990625

IT 213693-15-3P


RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation) (pentafluorobenzoyl)phenyl ether as monomers for F-containing aryl ether

ketone polymers having high solubility and heat stability)


RN 213693-15-3 CAPLUS

CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene) (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



L60 ANSWER 8 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN

AB The compns., useful for elec. insulating materials, contain (p-C6F4COC6H4-qxqOR1)m (R1 = C6H4-RX'z(CO-p-C6F4OR2)po; R2 = divalent aryl; X, X' = halo, lower alkyl, alkoxy; q, z = 0-4; m, p = 0, 1). Thus, 2,2-bis(4-(4-hydroxyphenyl)-1,1,1,3,3,3-hexafluoropropane was polymerized with 4,4'-bis(2,3,4,5,6-pentafluorobenzoyl)diphenyl ether to give a polymer showing dielec. constant 3.06 at 25°, 10% weight loss temperature 524° under N, and Tg 174°.


ACCESSION NUMBER: 2001:124290 CAPLUS  
DOCUMENT NUMBER: 134:179345  
TITLE: Low dielectric fluorinated aromatic polyether ketone compositions with good heat resistance  
INVENTOR(S): Kimura, Kunio; Yamashita, Yoshihiko; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 21 pp.  
CODEN: JG00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001049110	A2	20010220	JP 1999-226981	19990810
JP 3539897	B2	20040707		


PRIORITY APPLN. INFO.: JP 1999-226981 19990810

IT 213693-15-3P  
RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(low dielec. fluorinated aromatic polyether ketone compns. with good heat resistance)  
RN 213693-15-3 CAPLUS  
CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



L60 ANSWER 9 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN

AB Ketonic di-Ph ethers of p-R1COC6H4OC6H4R2-p type compds. (R1 = 2,3,4,5,6-pentafluorobenzoyl; R2 = OH, pentafluorobenzoyl group) and polyether-polyketone polymers containing tetrafluorophenylene and phenylene groups are provided which have good mech. strength, toughness, elec. property, thermal oxidative stability and solubility. Thus, heating 0.5 g 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone with 0.36 ground K carbonate, 2 mL N-methyl-2-pyrrolidone and 1 mL PhMe at 160° while stirring for 3 h gave a polymer at 85% yield and having viscosity 0.5 g/dL in AcNMe2.


ACCESSION NUMBER: 2001:25791 CAPLUS  
DOCUMENT NUMBER: 134:86663  
TITLE: (2,3,4,5,6-Pentafluorobenzoyl)diphenyl ether compound, and fluorine-containing aryl ether ketone polymer  
INVENTOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.;  
Fitch, John W., III; Reddy, V. Sreenivasulu  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: U.S., 22 pp., Cont.-in-part of U.S. Ser. No. 106,270, abandoned.  
CODEN: USXXAM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6172181 B1		20010109	US 1999-354976	19990716
			US 1998-106270	19980629


PRIORITY APPLN. INFO.: MARPAT 134:86663

OTHER SOURCE(S):  
IT 213693-15-3P  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(manufacture of (2,3,4,5,6-pentafluorobenzoyl)diphenyl ether compound and fluorine-containing aryl ether ketone polymer)  
RN 213693-15-3 CAPLUS  
CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT


AB Fluoropolymer-polyether-polyketones were obtained by homopolycondensation of 4-hydroxy-4'-(pentafluorobenzoyl)diphenyl ether and by copolycondensation of 4,4'-(pentafluorobenzoyl)diphenyl ether with benzenediols or bisphenols. The polymers have very good heat resistance, with 10% weight loss temps.  $\geq 500^\circ$ .

ACCESSION NUMBER: 1998:532324 CAPLUS  
 DOCUMENT NUMBER: 129:260978  
 TITLE: New polymers derived from 2,3,4,5,6-pentafluorobenzoic acid  
 AUTHOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.; Fitch, John W., III; Reddy, V. Sreenivasulu; Sakaguchi, Yoshimitsu  
 CORPORATE SOURCE: Faculty of Environmental Science and Technology, Okayama University, Okayama, 700-8530, Japan  
 SOURCE: Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1998), 39(2), 790-791  
 CODEN: ACPAY; ISSN: 0032-3934  
 PUBLISHER: American Chemical Society, Division of Polymer Chemistry  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 213693-15-3P, 4,4'-Bis(pentafluorobenzoyl)diphenyl ether-hydroquinone copolymer, SRU  
 RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of thermally stable fluoropolymer-polyether-polyketones)  
 RN 213693-15-3 CAPLUS  
 CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

AB Selected perfluorinated tertiary alcs. were reacted with p-toluenesulfonyl chloride to form their p-toluenesulfonyl esters C6F5(CF3)2COSO2C6H4CH3 (I), CF3C6F4OC6F4(C3F7)(C8F17)COSO2C6H4CH3 and (CF3C6F4OC6F4)2(C7F15)COSO2C6H4CH3. The absolute configuration of I is established by X-ray diffraction.

ACCESSION NUMBER: 1998:210377 CAPLUS  
 DOCUMENT NUMBER: 128:321424  
 TITLE: p-Toluenesulfonyl esters of perfluorinated tertiary alcohols: crystal structure determination of the absolute configuration of C6F5(CF3)2COSO2C6H4CH3  
 AUTHOR(S): Krumm, Burkhard; Vij, Ashwani; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
 CORPORATE SOURCE: Dep. of Chem., Univ. of Idaho, Moscow, ID, 83844-2343, USA  
 SOURCE: Journal of Fluorine Chemistry (1998), 89(1), 19-22  
 CODEN: JFLCAR; ISSN: 0022-1139  
 PUBLISHER: Elsevier Science S.A.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 128:321424  
 IT 207233-54-3  
 RL: RCT (Reactant); RACT (Reactant or reagent) (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal structure determination of the absolute configuration of C6F5(CF3)2COSO2C6H4CH3)  
 RN 207233-54-3 CAPLUS  
 CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentafluoroheptyl)- $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]- (9CI) (CA INDEX NAME)




IT 207233-53-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal structure determination of the absolute configuration of


RN 207233-53-2 CAPLUS

CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentadecafluoroheptyl)-  
 $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]-, 4-methylbenzenesulfonate (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L60 ANSWER 12 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN

AB Reactions of 4'-CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Li, generated in situ, with elements of group 16 (S, Se, Te) lead to CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SH (2), (CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Se)<sub>2</sub> (3), and (CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Te)<sub>2</sub> (4)/(CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Te)<sub>2</sub> (4a). The phenol derivative CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>OH (1) is obtained by reaction of CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Li with B(OMe)<sub>3</sub>/H<sub>2</sub>O<sub>2</sub>. The reaction of CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>Li with trimethylsilyl chloride or trimethyltin chloride gives CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>XMe<sub>3</sub> (X = Si (5), Sn (6)). Oxidation of 2 in the presence of bromine results in the formation of (CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SH)<sub>2</sub> (7) and CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SHO<sub>2</sub>Br (8). Mixed perfluoroaryloxy/thio ethers CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SCF<sub>4</sub>R (R = NO<sub>2</sub> (9), CN (10),

CF<sub>3</sub> (11)) and CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SCF<sub>4</sub>N (12) are obtained upon reaction of 2 with excess C<sub>6</sub>F<sub>5</sub>R and pentafluoropyridine in the presence of K<sub>2</sub>CO<sub>3</sub>. With 4-C<sub>6</sub>F<sub>5</sub>OC<sub>6</sub>F<sub>4</sub>NO<sub>2</sub>, a mixture of (2-CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SH) (4-C<sub>6</sub>F<sub>5</sub>OC<sub>6</sub>F<sub>4</sub>NO<sub>2</sub>) (13) and 9 is formed. Reaction of excess 2 with C<sub>6</sub>F<sub>5</sub>R gives the 2,4,6-substituted benzenes (CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>SH)<sub>2</sub> (R = NO<sub>2</sub> (14), CN (15)). The trimethylsilyl ether CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>OSiMe<sub>3</sub> (16) is prepared from the reaction

of 1 with hexamethyldisilazane. 16 is a convenient reagent for the preparation of the aryl ethers CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>OCF<sub>4</sub>R (R = NO<sub>2</sub> (17), CN (18)) and CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>OCF<sub>4</sub>N (19) upon reaction with C<sub>6</sub>F<sub>5</sub>R and C<sub>6</sub>F<sub>5</sub>N. The secondary alcs. CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>CH(C<sub>6</sub>H<sub>5</sub>)OH (20) and CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>CH(C<sub>6</sub>F<sub>5</sub>)OH (21) are synthesized by the reactions of 5 with benzaldehyde and pentafluorobenzaldehyde in the presence of tetrabutylammonium fluoride as a catalyst. In the synthesis of 21 the byproduct CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub>CH(C<sub>6</sub>F<sub>5</sub>)OC<sub>6</sub>F<sub>4</sub>CHO is also formed and isolated.

ACCESSION NUMBER: 1997:667252 CAPLUS  
 DOCUMENT NUMBER: 127:293323  
 TITLE: Synthesis and Chemistry of CF<sub>3</sub>CF<sub>3</sub>CF<sub>4</sub>CF<sub>4</sub> Group 14/16 Derivatives  
 AUTHOR(S): Krumm, Burkhard; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
 CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 83844-2343, USA  
 SOURCE: Inorganic Chemistry (1997), 36(23), 5222-5230  
 CODEN: INOCAL; ISSN: 0020-1669  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 127:293323


IT 197150-21-3P 197150-22-4P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)

RN 197150-21-3 CAPLUS


CN Benzene, 1,2,4,5-tetrafluoro-3-(2,3,5,6-tetrafluoro-4-nitrophenoxy)-6-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy)- (9CI) (CA INDEX NAME)

L60 ANSWER 12 OF 32 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 197150-22-4 CAPLUS

CN Benzonitrile, 2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT




AB Phthalic acids I (R = CO<sub>2</sub>H; X = halo) are prepared by heating phthalonitriles I (R = cyano; X = halo) in acidic aqueous media, (A) dissolving the products in organic solvent-containing media, (B) separating the products, and (C) treating the isolated products in acidic aqueous media under heating. Then, the above processes A to C are repeated. ≥1 time(s). I (R = cyano, X = F) (10 g) was refluxed with aqueous H<sub>2</sub>SO<sub>4</sub> for 6 h, diluted with H<sub>2</sub>O, filtered, and the crude product was dissolved in Me<sub>2</sub>CO. The solution was treated with H<sub>2</sub>O to give 10.8 g I (R = CO<sub>2</sub>H, X = F) (II) with purity 91.8%, which was refluxed with H<sub>2</sub>SO<sub>4</sub> for 6 h and similarly treated. The process was repeated to give 9.8 g II with 99.2% purity.

ACCESSION NUMBER: 1997:377463 CAPLUS  
DOCUMENT NUMBER: 127:33992  
TITLE: Preparation of phthalic acids as intermediates for polyimides from phthalonitriles  
INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
CODEN: JMOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110785	A2	19970428	JP 1995-266007	19951013

PRIORITY APPLN. INFO.: JP 1995-266007 19951013

OTHER SOURCE(S): MARPAT 127:33992  
IT 143363-92-2P  
RL: IMF (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)  
(preparation and purification of phthalic acids as intermediates for polyimides by acid hydrolysis of phthalonitriles)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation and purification of phthalic acids as intermediates for polyimides by acid hydrolysis of phthalonitriles)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




AB Title compds. I (X = halo; Y = CO<sub>2</sub>H) are prepared by heating I (Y = cyano) in acidic aqueous media, heating the product-containing media with alkaline substances, and separating I (Y = CO<sub>2</sub>H) from the media by mixing with acidic substances. An aqueous solution containing 10 g I (X = F, Y = cyano) and H<sub>2</sub>SO<sub>4</sub> was refluxed for 6 h, mixing with H<sub>2</sub>O and NaOH (pH 11.8), refluxed for 1 h, and mixed with aqueous H<sub>2</sub>SO<sub>4</sub> to give 10.8 g I (X = F, Y = CO<sub>2</sub>H) with 99.2% purity.

ACCESSION NUMBER: 1997:377462 CAPLUS  
DOCUMENT NUMBER: 127:33986  
TITLE: Preparation of phthalic acids as intermediates for polyimides  
INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JMOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110784	A2	19970428	JP 1995-266006	19951013


PRIORITY APPLN. INFO.: JP 1995-266006 19951013

OTHER SOURCE(S): MARPAT 127:33986  
IT 143363-92-2P  
RL: IMF (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of phthalic acids as intermediates for polyimides by hydrolysis of phthalonitriles and heating with alkalies)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of phthalic acids as intermediates for polyimides by hydrolysis of phthalonitriles and heating with alkalies)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)





AB The title bisphthalonitriles I (X, Y = O, S; R = Z, Z1), useful as intermediates for optical materials, circuit board materials, photosensitive materials, liquid crystals, etc., are prepared by treatment of 3,4,5,6-tetrafluorophthalonitrile (II) with HORYH in organic solvents being slightly soluble in H2O in the presence of basic substances, followed by addition of H2O to the reaction products and separation of the organic layer. An AcOEt solution of tetrafluorohydroquinone was added dropwise to a mixture of II, KF, and AcOEt under reflux over 1 h, and the reaction mixture was further stirred under reflux for 8 h, cooled, and then washed with H2O to give 99% I (X = Y = O, R = Z).



ACCESSION NUMBER: 1997:154684 CAPIUS  
DOCUMENT NUMBER: 126:157295  
TITLE: Preparation of bis(fluorophthalonitriles) from tetrafluorophthalonitrile and fluorohydroquinone or fluorodiphenol  
INVENTOR(S): Okumura, Yasunori; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JI00AF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08333322	A2	19961217	JP 1995-143593	19950609

PRIORITY APPLN. INFO.: JP 1995-143593 19950609

OTHER SOURCE(S): CASREACT 126:157295; MARPAT 126:157295  
IT 143376-50-59  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP

L60 ANSWER 15 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
(Preparation)  
[prepn. of bis(fluorophthalonitriles) from tetrafluorophthalonitrile and fluorohydroquinone, diphenol, or their sulfur analog using slightly water-sol. solvents and basic substances]  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



AB Polyimide optical materials comprise a perfluorinated polyimide having a perfluorinated repeating unit represented by the general formula I (R1 = a tetraavalent perfluorinated organic group; and R2 = a divalent perfluorinated organic group).


ACCESSION NUMBER: 1995:854348 CAPIUS  
DOCUMENT NUMBER: 123:354214  
TITLE: Polyimide optical material  
INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni; Yamamoto, Fumio  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corporation, Japan  
SOURCE: U.S., 26 pp. Cont.-in-part of U.S. Ser. No. 54,973, abandoned.  
CODEN: USXXAM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5449741	A	19950912	US 1993-140982	19931025
JP 09031014	A2	19970204	JP 1996-207475	19910913
US 5233018	A	19930803	US 1991-765672	19910926
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 5750731	A	19980512	US 1996-718208	19960920
US 5849934	A	19981215	US 1998-20573	19980128
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		


PRIORITY APPLN. INFO.: JP 1990-256843 A 19900928  
JP 1991-106552 A 19910412  
JP 1991-106554 A 19910412  
JP 1991-106557 A 19910412  
US 1991-765672 A3 19910926  
US 1993-54973 B2 19930430  
JP 1991-235020 A3 19910913

L60 ANSWER 16 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
CA 1991-2052368 A3 19910926  
US 1993-140482 A3 19931025  
US 1993-140982 A3 19931025  
US 1995-451465 A1 19950526  
US 1996-718208 A3 19960920  
US 1998-20573 A3 19980128

IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(polyimide optical materials)  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143363-92-2  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reg. polyimide optical materials)  
RN 143363-92-2 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)





\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I, useful as material for hardeners for fluorinated epoxy resins (no data), is prepared via II [R = cyano, CO<sub>2</sub>H]. Thus, a mixture of

tetrafluorophthalonitrile, tetrafluorohydroquinone, and Et<sub>3</sub>N in DMF was heated at 35° for 30 to give 21% II [R = cyano], which was treated with 60% H<sub>2</sub>SO<sub>4</sub> at 150° for 5 h to give 26% II [R = CO<sub>2</sub>H], which was refluxed with Ac<sub>2</sub>O for 2 h to 52% I.

ACCESSION NUMBER: 1994:630662 CAPIUS

DOCUMENT NUMBER: 121:230662  
TITLE: preparation of a perfluorinated hexacarboxylic acid as

material for hardeners for fluorinated epoxy resins

INVENTOR(S): Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji

PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKOQAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06157501	A2	19940603	JP 1992-340986	19921130
PRIORITY APPLN. INFO.:			JP 1992-340986	19921130


OTHER SOURCE(S): CASREACT 121:230662

IT 158394-12-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and conversion into trianhydride)

RN 158394-12-8 CAPIUS

CN 1,2-Benzenedicarboxylic acid, 4,5-bis[4-(3,4-dicarboxy-2,5,6-trifluorophenoxy)-3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)





IT 158394-11-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation and hydrolysis of)

RN 158394-11-7 CAPIUS

CN 1,2-Benzenedicarbonitrile, 4,5-bis[4-(3,4-dicyano-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)



AB The title derivs. I (X = O, S; Y = O, S, Q1, Q2) are prepared by treating 28 mol 3,4,5,6-tetrafluorophthalonitrile (II) with 1 mol phenols or thiophenols III in the presence of basic substances in organic solvents.

A mixture of 0.22 mol II and KF in MeCN was treated dropwise with a solution of 0.011 mol III (X = Y = O) in MeCN under reflux over 1 h, then refluxed for

4 h to give 99% I.

ACCESSION NUMBER: 1994:435030 CAPIUS

DOCUMENT NUMBER: 121:35030  
TITLE: Preparation of fluorine-containing phthalonitrile derivatives as intermediates for fluorine-containing polyimides

INVENTOR(S): Kaieda, Osamu; Okumura, Yasunori; Ito, Hideki

PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKOQAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06016615	A2	19940125	JP 1992-176633	19920703
JP 08000803	B4	19960110		
PRIORITY APPLN. INFO.:			JP 1992-176633	19920703

OTHER SOURCE(S): CASREACT 121:35030; MARPAT 121:35030


IT 143376-50-5P

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation of, as intermediate for polyimides)

RN 143376-50-5 CAPIUS

CN 1,2-Benzenedicarbonitrile, 4,4'-bis[4-(2,3,5,6-tetrafluoro-1,4-







AB The title compds. (I; R1, R2 = F, C1-12 linear or branched alkoxy, C1-12 alkylthio, C1-12 alkylamino, aryloxy, arylthio, arylamino, Q - Q2; X = O, S; Y1 = CO2CO), useful as intermediates for drugs, agrochems., optical materials, printed circuit board materials, photosensitive materials, and liquid crystal materials, are prepared by heating F-containing phthalic acid

derivs. (II; R1, R2, X = same as above; Y, Y1 = CO2H) in a solvent selected from SOCl2, POCl3, or AcCl at 40-105°. The process is safe and gives perfluorophthalic anhydride derivs. I of high purity in high yields. Thus, 70.5 g SOCl2 was added to 30.0 g tetrafluorophthalic acid and the resulting mixture was allowed to react at 70° for .apprx.3 h to give 97% tetrafluorophthalic anhydride.

ACCESSION NUMBER: 1994:298457 CAPIUS  
DOCUMENT NUMBER: 120:298457  
TITLE: Preparation of fluorine-containing phthalic anhydride derivatives  
INVENTOR(S): Okumura, Yasunori; Ito, Hideki; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
CODEN: JIQQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328502	A2	19921117	JP 1991-125451	19910426

AB The filter, comprising a dielec. multilayer, is characterized in that the substrate consists of a thin-film perfluoropolyimide or a blend or a copolymer thereof.


ACCESSION NUMBER: 1993:417687 CAPIUS  
DOCUMENT NUMBER: 119:17687  
TITLE: IR longpass dielectric interference filter  
INVENTOR(S): Yamamoto, Fumio; Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji; Oguchi, Taisuke  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JIQQAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328502	A2	19921117	JP 1991-125451	19910426

IT 148273-08-9  
RL: USES (Uses)  
(fluorinated polyimide blends containing, IR longpass filters containing)  
RN 148273-08-9 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1

CRN 143363-92-2  
CMF C22 H4 F10 O10



CM 2

CRN 1198-63-6  
CMF C6 H4 F4 N2



JP 06016656	A2	19940125	JP 1992-175450	19920702
JP 3130653	B2	20010131		

PRIORITY APPLN. INFO.: JP 1992-175450 19920702

OTHER SOURCE(S): CASREACT 120:298457; MARPAT 120:298457

IT 143363-92-2  
RL: PROC (Process)  
(conversion of, into acid anhydride)

RN 143363-92-2 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




L60 ANSWER 23 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB A perfluorinated polyimide that has glass temperature >300° and high optical transparency over the entire optical communication wavelengths was synthesized. The high thermal stability and optical transparency in near-IR region are due to the fully aromatic mol. structure and the absence of C-H bonds. The use of a diamine with relatively high reactivity and a new perfluorinated dianhydride, which has flexible structure, makes it possible to obtain a tough and flexible perfluorinated polyimide film.

In addition, this polymer has a low dielec. constant Perfluorinated polyimides are promising for use as optical-electronic materials.


ACCESSION NUMBER: 1992:592808 CAPIUS  
 DOCUMENT NUMBER: 117:192808  
 TITLE: Perfluorinated polyimide synthesis  
 AUTHOR(S): Ando, Shinji; Matsuura, Tohru; Sasaki, Shigekuni  
 CORPORATE SOURCE: Interdiscip. Res. Lab., NTT, Musashino, 180, Japan  
 SOURCE: Macromolecules (1992), 25(21), 5858-60  
 CODEN: MAMOBX; ISSN: 0024-9297  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

IT 143376-50-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (Reactant or reagent)  
 (preparation and hydrolysis of)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]




L60 ANSWER 24 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 US 1998-20573 A3 19980128

IT 143376-50-5P  
 RL: RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
 (Reactant or reagent)  
 (preparation and hydrolysis of)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



IT 143363-92-2P  
 RL: PREP (Preparation)  
 (Preparation of)  
 RN 143363-92-2 CAPIUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



L60 ANSWER 24 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The title polymers, useful as waveguides in optoelectronic integrated circuits or optoelec. boards, are prepared from perfluoro tetracarboxylic acid derivs. and perfluoro diamines, the preparation of which is described.

Polymerization of 4,4'-(tetrafluoro-p-phenylene)bis[3,5,6-trifluorophthalic anhydride] with tetrafluoro-1,3-phenylenediamine gave a polymer with no substantial absorption at optical communication wavelengths.

ACCESSION NUMBER: 1992:531752 CAPIUS  
 DOCUMENT NUMBER: 117:131752  
 TITLE: Monomers for preparation of perfluorinated polyamic acids and polyimides  
 INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni; Yamamoto, Fumio  
 PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
 SOURCE: Eur. Pat. Appl., 39 pp.  
 CODEN: EPOXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 480266	A2	19920415	EP 1991-116553	19910927
EP 480266	A3	19930901		
EP 480266	B1	19960626		
R: DE, FR, GB				
JP 05001148	A2	19930108	JP 1991-235020	19910913
JP 2851019	B2	19950127		
JP 09031014	A2	19970204	JP 1996-207475	19910913
CA 2052368	AA	19920329	CA 1991-2052368	19910926
CA 2052368	C	19980915		
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		
PRIORITY APPL. INFO.:				
			JP 1990-256843	A 19900928
			JP 1991-106552	A 19910412
			JP 1991-106554	A 19910412
			JP 1991-106557	A 19910412
			JP 1991-235020	A3 19910913
			CA 1991-2052368	A3 19910926
			US 1991-765672	A3 19910926
			US 1993-54973	B2 19930430
			US 1993-140482	A3 19931025
			US 1995-451465	B1 19950526
			US 1996-718208	A3 19960920


L60 ANSWER 25 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The title polymers, e.g., polyether-polyketones, polyarylates, polyimides, and polyesters, have good heat resistance and low attenuation and are useful as optical transmitting systems, e.g., for controlling ignition timing and fuel metering systems for internal combustion engines in automobiles. Thus, an optical fiber comprised a core of amorphous PEEK and a sheath of poly(2,2,2-trifluoroethyl methacrylate).

ACCESSION NUMBER: 1992:130739 CAPIUS  
 DOCUMENT NUMBER: 116:130739  
 TITLE: Amorphous polymers for optical transmitting systems and optical members and their use  
 INVENTOR(S): Takekawa, Yoshitaka; Ohara, Shuichi; Tanno, Seikichi; Taketani, Noriaki; Shimura, Masato  
 PATENT ASSIGNEE(S): Hitachi, Ltd., Japan  
 SOURCE: Eur. Pat. Appl., 31 pp.  
 CODEN: EPOXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 454165	A2	19911030	EP 1991-106851	19910426
EP 454165	A3	19930120		
R: DE, FR, GB, IT, NL				
JP 04009805	A2	19920114	JP 1990-112511	19900427
JP 2854669	B2	19990203		
US 5093888	A	19920303	US 1991-686997	19910418
PRIORITY APPL. INFO.:				
			JP 1990-112511	A 19900427

IT 138687-03-3  
 RL: USES (Uses)  
 (optical fibers, heat-resistant, for engine control systems)  
 RN 138687-03-3 CAPIUS  
 CN Poly[oxy(2,6-dimethyl-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)]





AB Benzyl cyclopropanecarboxylates I (R = H, F, Cl, Br; R1 = H, cyano, C.tplbond, CH; R2 = H, Me; R3 = CH:CCl2, CHBrCBrCl2, CHBrCBr3, Me), having insecticidal and acaricidal activities at 500 ppm with low fish toxicity, were prepared e.g. from II. Thus, 6.0 mmol NaCN and 0.88 mmol


PhCH2NEt3Cl in H2O was treated dropwise with 4.0 mmol II and 4.2 mmol dl-cis,trans-2,2-dimethyl-3-(2,2-dichlorovinyl)cyclopropanecarbonyl chloride in PhMe to give 1.93 g I (R = R2 = H, R1 = cyano, R3 = CH:CCl2).

ACCESSION NUMBER: 1984:610760 CAPLUS  
DOCUMENT NUMBER: 101:210760  
TITLE: Insecticidal cyclopropanecarboxylate esters  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKQXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59134749	A2	19840802	JP 1983-8536	19830120
PRIORITY APPLN. INFO.: JP 1983-8536 19830120				

IT 90928-31-7P 93207-48-8P  
RI: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation and insecticidal activity of)

RN 90928-31-7 CAPLUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2,2-tetrabromoethyl)-, 1-[4-fluoro-3-(4-fluorophenoxy)phenoxy]phenyl-2-propynyl ester (9CI) (CA INDEX NAME)



RN 93207-48-8 CAPLUS  
CN Cyclopropanecarboxylic acid, 3-(2,2-dichloroethyl)-2,2-dimethyl-, cyano[4-fluoro-3-(4-(4-fluorophenoxy)phenoxy]phenylmethyl ester (9CI) (CA INDEX NAME)



AB Twelve title esters (I; R = H, halo; R1 = H, F; R2 = H, cyano, HC.tplbond, C, R3 = haloethyl, haloethyl), effective insecticides and miticides at 500 ppm, were prepared. Thus, 5.0 mmol II and 5.25 mmol (±)-III in MePh were added to a solution of 7.5 mmol NaCN and 1.1 mmol PhCH2NEt3 Cl- in H2O at room temperature to give 2.07 g (±)-I (R = R1 =


H; R2 = cyano, R3 = Cl2C:CH).

ACCESSION NUMBER: 1984:438168 CAPLUS  
DOCUMENT NUMBER: 101:38168  
TITLE: Cyclopropanecarboxylate esters as insecticides and miticides with low toxicity  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
CODEN: JKQXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59042342	A2	19840308	JP 1982-153029	19820901
PRIORITY APPLN. INFO.: JP 1982-153029 19820901				

IT 90928-31-7P  
RI: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 90928-31-7 CAPLUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2,2-tetrabromoethyl)-, 1-[4-fluoro-3-(4-(4-fluorophenoxy)phenoxy]phenyl-2-propynyl ester (9CI) (CA INDEX NAME)



L60 ANSWER 28 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The PbO<sub>2</sub> oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).

ACCESSION NUMBER: 1974:520135 CAPIUS

DOCUMENT NUMBER: 81:120135

TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol

AUTHOR(S): Denivelle, Leon; Huynh Anh Hoa

CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conservatoire Natl. Arts

SOURCE: Metiers, Paris, Fr. Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 487-90

CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal


LANGUAGE: French

IT 53359-93-6P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 53359-93-6 CAPIUS

CN Phenol, 2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



L60 ANSWER 29 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The controlled oxidation of perhalo p-bromophenols (I), (II), and (III) gives

the corresponding perhalo polyethers (IV): the perchloro 4-aryloxycyclohexa-2,5-dienone (V) is obtained from pentachlorophenol (VI). Similarly, VII gives VIII and 4-chloro-2,3,5,6-tetrabromophenol gives IX.

ACCESSION NUMBER: 1971:111698 CAPIUS

DOCUMENT NUMBER: 74:111698

TITLE: Oxidation of pentahalobenzenic phenols

AUTHOR(S): Denivelle, Leon; Lampel, Alfred

CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conserv. Natl. Aris

SOURCE: Metiers, Paris, Fr. Comptes Rendus des Seances de l'Academie des

Sciences,

Serie C: Sciences Chimiques (1971), 272(7), 653-6

CODEN: CHDCAQ; ISSN: 0567-6541

DOCUMENT TYPE: Journal


LANGUAGE: French

IT 31404-37-2P 31404-38-3P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)


RN 31404-37-2 CAPIUS

CN 2,5-Cyclohexadien-1-one, 4-bromo-4-[4-(4-bromo-2,3,5,6-tetrafluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



RN 31404-38-3 CAPIUS

CN Phenol, 4-[4-[4-(4-bromo-2,3,5,6-tetrafluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



L60 ANSWER 29 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

(Continued)

L60 ANSWER 30 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

AB A number of perfluorinated and partially fluorinated phenyl and polyphenyl

ethers were synthesized, characterized for thermal stability, fire resistance, and viscosity, and compared with their H analogs to assess

the potential use of this class of compds. as functional fluids. Without exception, polyfluorination and perfluorination lower thermal stability; the decrease in stability depends on the position and number of fluorine substituents. The autoignition temperature and fire resistance are not

improved over the H analog, and viscosity is degraded. These data coupled with

the comparatively high melting points do not suggest a bright future for this class of compds. as useful functional fluids. 6 references.

ACCESSION NUMBER: 1968:77890 CAPIUS

DOCUMENT NUMBER: 68:77890

TITLE: Synthesis, thermal stability, flammability, and viscosity of some partially fluorinated and

perfluorinated aromatic and polyaromatic ethers

AUTHOR(S): Richardson, George Albert; Blake, Edward S.

CORPORATE SOURCE: Monsanto Res. Corp., Dayton, OH, USA

SOURCE: Industrial & Engineering Chemistry Product Research

and Development (1968), 7(1), 17-21

CODEN: IEPRA6; ISSN: 0196-4321

DOCUMENT TYPE: Journal


LANGUAGE: English

IT 6804-37-1P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 6804-37-1 CAPIUS

CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)- (7CI, 8CI) (CA INDEX NAME)



L60 ANSWER 31 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB 2,3,5,6-Tetrafluoro-O,O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]hydroquinones (I) and 2,4,5,6-tetrafluoro-O,O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]resorcinols (II) are prepared by the

treatment of the compds. of the general formulas C6F5X, where X is CN, CF3, CONH2, and CO2H, with 2,3,5,6-tetrafluorohydroquinone di-Na salt and 2,4,5,6-tetrafluororesorcinol di-Li salt (III), resp. 19F-N.M.R. and ir data for the prepared I and II are given. III is treated with decafluorocyclohexene to give a mixture of 2,4,5,6-tetrafluoro-O,O'-bis(perfluoro-1-cyclohexen-1-yl)resorcinol and 2,4,5,6-tetrafluoro-O-(perfluoro-1-cyclohexen-1-yl)-O'-(perfluoro-2-cyclohexen-1-yl)resorcinol.

ACCESSION NUMBER: 1968:68571 CAPIUS

DOCUMENT NUMBER: 68:68571

TITLE: Reactions of perfluorophenolates with substituted

pentafluorobenzenes and perfluorocyclohexene

AUTHOR(S): De Pasquale, Ralph J.; Tamborski, Christ

CORPORATE SOURCE: Wright-Patterson Air Force Base, Dayton, OH, USA

SOURCE: Journal of Organic Chemistry (1968), 33(2), 830-3

CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 14796-02-2P 14796-03-3P 15053-71-1P


15077-30-2P

RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

RN 14796-02-2 CAPIUS

CN Benzamide, 4,4'-[(2,3,5,6-tetrafluoro-p-phenylene)dioxy]bis[2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



RN 14796-03-3 CAPIUS

CN Benzoic acid, 4,4'-[(2,3,5,6-tetrafluoro-p-phenylene)dioxy]bis[2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)

L60 ANSWER 32 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

AB A mixture of 26.3 g. C6F6, 26.5 g. 85% KOH, and 75 ml. H2O was heated in a

sealed bomb at 175° for 5 hrs. with agitation to give 33.1 g. C6F5OH, b. 144-5°. A mixture of 67 g. C6F5H, 21.6 g. KOH, 150 ml. pyridine, and 2 ml. H2O was refluxed 1 hr., treated with 21.6 KOH, and refluxed 24 hrs. to give 21 g. 2,3,5,6-tetrafluorophenol, b20 47°. Similarly refluxing 36 g. C6F5Me and 28 g. KOH in 300 ml. tert-BuOH gave 12 g. 2,3,5,6-tetrafluoro-p-cresol, m. 52°. Reaction of 20 g. C6F5I with 2 g. KOH and 1 ml. H2O in 100 ml. pyridine gave only one product, 2,3,5,6-tetrafluoro-4-iodophenol, m. 79-81°; benzoate m. 59-60.2°. Reaction of 66 g. C6F5Br with 28.5 g. KOH and 1 ml. H2O in 150 ml. pyridine, however, gave a mixture of products: 3.5 g. 2-bromo-3,4,5,6-tetrafluorophenol, m. 41-3° (3,5-dinitrobenzoate, m. 104-5°); and 11.5 g. 4-bromo-2,3,5,6-tetrafluorophenol (3,5-dinitrobenzoate m. 131-3°). Similarly, 100 g. 2-chlorotetrafluoro- $\alpha,\alpha,\alpha$ -trifluorotoluene on treatment with 5.6 g. KOH and 1 ml. H2O in 100 ml. pyridine gave 2.5 g. 2-chlorotrifluoro- $\alpha,\alpha,\alpha$ -trifluoro-o-cresol, b15 92-3°, n24D 1.4510; and 15 g. 2-chlorotrifluoro- $\alpha,\alpha,\alpha$ -trifluoro-p-cresol, b15 102-3°, n24D 1.4510. The reactions of polyfluorobenzenes were then studied with alkoxides. A solution of 123.5 g. C6F5Br in 70 ml. pyridine was treated

with a solution of 11.5 g. Na in 150 ml. MeOH during 1.5 hrs. and the mixture refluxed 15 hrs. and acidified with 1 l. 10% HCl to give 66 g. 4-bromo-2,3,5,6-tetrafluoroanisole, b5 79-81°, n25D 1.4812. Similarly, a mixture of 10 g. C6F5I in 50 ml. pyridine and 0.8 g. Na in

15 ml. MeOH on refluxing for 3 hrs. gave 1.5 g. unchanged C6F5I and 5.5 g. 2,3,5,6-tetrafluoro-4-iodoanisole (I), b20 113-15°, n22D 1.5229. Refluxing 1 g. I with 1 g. activated Cu powder for 12 min. gave 0.2 g. octafluoro-4,4'-dimethoxybiphenyl, m. 90-1.2°. To a cold solution of 9 g. Na in 250 ml. PhCH2OH was added 75 g. C6F6 and the mixture refluxed

24 hrs. to give 30 g. benzyl pentafluorophenyl ether (II), m. 44°. A better yield was obtained when a solution of 4.6 g. Na and 22 g. PhCH2OH in 250 ml. tert-BuOH was refluxed with 40 g. C6F6 for 40 hrs. to give 33 g. II. A solution of 8 g. C6F6 in 30 ml. HCONMe2 was treated with 5.28 g.


PhOK and the mixture refluxed 0.5 hr. to give 1 g. 2,3,5,6-tetrafluoro-1,4-diphenoxybenzene, m. 147-9°, and 3.5 g. 2,3,4,5,6-pentafluorophenyl phenyl ether, (III), m. 29°. III was also obtained by heating a mixture of 11 g. C6F5OK, 15 g. PhBr, and 1 g. Cu at 210° in a sealed bomb. A solution of 6 g. C6F5OK and 12.8 g. C6F6 in 30 ml. HCONMe2 was refluxed 14 hrs. to give 1.5 g. bis(perfluorophenyl) ether, m. 67-9°, and a second product, m. 145-8°, probably

p-bis(pentafluorophenoxy)2,3,5,6-tetrafluorobenzene. Similarly, a solution of 0.6 g. Na in 50 ml. EtOH refluxed with 5.3 g. C6F5NHMe2 2 hrs. gave 5.1 g. 4-ethoxy-2,3,5,6-tetrafluoro-N,N-dimethylaniline, b. 34°. Reactions with amines were next investigated. A mixture of 280 g. C6F6

and 400 ml. 28% aqueous NH3 was rocked in a sealed bomb for 2 hrs. at 235° to give 236 g. C6F5NH2, m. 34°, and 28 g. tetrafluorophenylenediamine (sublimed 75/1 mm.) shown by its nuclear magnetic resonance spectrum to be essentially the meta isomer mixed with a small amount of the para isomer. Similarly, heating a mixture of

L60 ANSWER 31 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN


(Continued)



RN 15053-71-1 CAPIUS


CN Benzonitrile,

4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



RN 15077-30-2 CAPIUS

CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]benzene (9CI) (CA INDEX NAME)



L60 ANSWER 32 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN

(Continued)

56 g. C6F6 and 110 ml. 30% aq. MeNH2 at 220° for 3 hrs. gave 59% C6F5NHMe, b. 170-2°, and 25% 2,3,5,6-tetrafluoro-N,N'-diphenylphenylenediamine, m. 94°. The reaction product obtained by heating 50 g. C6F6 and 110 ml. 25% aq. MeNH2 at 235° for 1 hr. was distd. at 1 mm. pressure and five fractions were collected. The first fraction (65%), b1 88°, was C6F5NHMe2. Fraction 2, b1 88-126°, was shown by vapor phase chromatography to be C6F5NHMe2 with 3 other compds. Fraction 3, b1 126-134°, consisted of 3 isomers of bis(dimethylamino)tetrafluorobenzene with the meta-isomer predominating. Fraction 4, b1 134-40°, contained equal ams. of the meta and para isomers. Fraction 5, b1 140-8°, was pure para isomer. The meta and para isomers could be sepd. by vapor phase chromatography. Similarly, heating a mixt. of 30 g. C6F5Br and 70 ml.

281

NH4OH at 200° for 2 hrs. gave 22 g. p-bromotetrafluoroaniline, m. 61°. Heating 16 g. C6F5I and 30 ml. 8% NH4OH at 165° for 2 hrs. gave 7.6 g. tetrafluoro-p-iodoaniline, m. 77°. Benzyl pentafluorophenyl ether (20 g.) was heated with large excess of 28% NH4OH to give 3 g. p-(benzyloxy)tetrafluoroaniline, m. 97°. Similarly, heating 50 g. 2-chlorotetrafluoro- $\alpha,\alpha,\alpha$ -trifluorotoluene and 120 ml. 28% NH4OH at 21° for 2 hrs. gave 22 g. 2-chlorotrifluoro- $\alpha,\alpha,\alpha$ -trifluoro-p-toluidine, which decompd. readily at room temp. in the presence of air. To 100 ml. anhyd. NH3 at -70° were added 0.1 g. Fe(NO3)3 and 2.99 g. Na and, after disappearance of the blue color, 25 g. C6F5OMe during 45 min. After 5 hrs. at -70° the reaction mixt. was worked up to give 7 g. unreacted C6F5OMe, 2.8 g. tetrafluoro-p-anisidine, m. 75-6.5°, 1.2 g. 4,4'-dimethoxyoctafluorodiphenylamine, m. 78-9°, and 2.2 g. 4,4',4''-trimethoxydecafluorotriphenylamine, b. 157-9°, n23D 1.5005. Diazotization of C6F5NH2 required concd. acids since the salts

of


the amine hydrolyzed very readily in dil. solns. In 48% HBr, diazotization of C6F5NH2 gave C6F5N+Br- (C6F5), probably owing to slow diazotization. The reaction was temp.-dependent, the diazaminobenzene being formed much faster at 10° than at -10°. The product decompd. in warm HBr to give 5.4% C6F5Br and a mixt. of o- and p-dibromotetrafluorobenzenes. In concd. H2SO4 the reaction was very slow even at 25°. Addition of HNO3 hastened it. Deamination with hypophosphorus acid gave a mixt. probably of C6F5H and C6H2F4. Better diazotization could be carried out in liquid HF and the diazo product underwent successful Sandmeyer reaction. A soln. of 20 g. C6F5NH2 in 75 ml. anhyd. HF at -20° was treated with 7.27 g. NaNO2 during 30 min. After stirring for 1 hr. at -10° the mixt. was treated with 17.6 g. KI during 30 min. and allowed to warm to 25° in 1 hr. to give 16.5 g. C6F5I, b35 77-9°. Use of 12 g. KBr and 15 g. Cu2Br2 instead of KI gave 35% C6F5Br. The reaction of diazotized amine with C6F5OLi gave C6F5N+Li-(O)C6F5, which decompd. on removal of solvent. The Sandmeyer nitrile synthesis was not successful. A soln. of 10 g. C6F5NH2 in 100

ml.

HOAc was oxidized with 25 ml. 30% H2O2 at 25° for 24 hrs. to give decafluoroazobenzene (IV), m. 53-4°. A mixt. of 5 g. IV, 15 g. Zn powder, 5 g. NH4Cl, 10 ml. H2O, and 75 ml. 95% EtOH refluxed 30 min. gave 2 g. decafluoroazobenzene, m. 57-9°. C6F6 reacted readily with organolithium compds. A soln. of MeLi, prepd. from 4.5 g. Li and 43 g. MeI in 50 ml. ether, was cooled to -10° to -20°, added dropwise to a soln. of 60 g. C6F6 in 250 ml. pentane, and stirred for 17 hrs. at room temp. to give 34 g. C6F5Me, b. 115°. Similarly, reaction of BuLi, prepd. from 1.86 g. Li and 18.3 g. BuBr in 30 ml. ether,

L60 ANSWER 32 OF 32 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 with 25.3 g. C6F6 in 25 ml. ether gave 10.5 g. unreacted C6F6, 7 g. C6F5Bu, b25 86-7°, n20D 1.4229, and 2.5 g. of a compd., b1 230°, n20D 1.4683, probably impure tributylidifluorobenzene. Similarly, 32.7 g. C6F6 in 150 ml. ether with 0.18 mole PhLi in 250 ml. ether gave 8.5 g. 2,3,5,6-tetrafluoro-p-terphenyl, m. 220°, and 33 g. C6F5Ph, m. 69°. A similar reaction of 18.6 g. C6F6 and isopropenyllithium prepd. from 12.1 g. 2-bromopropene gave 5 g. 2,3,4,5,6-pentafluoro- $\alpha$ -methylstyrene, b52 72-4°. With vinylolithium, prepd. from 0.1 mole PhLi and 0.025 mole tetravinyltin, 18.6 g. C6F6 gave 4 g. unreacted C6F6 and 20% C6F5CH:CH2, b25 34°. LiAlH4 redn. of 21 g. C6F6 in ether gave 17 g. of a mixt. of C6F6 and C6F5H which was sepd. by vapor phase chromatography to give 7.5 g. C6HF5. All the products in all above reactions were studied by infrared and nuclear magnetic resonance spectroscopy. The mechanism of reaction and the directional effects were discussed.

ACCESSION NUMBER: 1964:52434 CAPIUS  
 DOCUMENT NUMBER: 60:52434  
 ORIGINAL REFERENCE NO.: 60:9170b-h, 9171a-e  
 TITLE: Reactions of polyfluorobenzenes with nucleophilic reagents  
 AUTHOR(S): Wall, Leo A.; Pummer, Walter J.; Fearn, James E.; Antonucci, Joseph M.  
 SOURCE: J. Res. Natl. Bur. Std. (1963), 67A(5), 481-97  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 IT 6804-37-1, Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)-(7)  
 (preparation of)  
 RN 6804-37-1 CAPIUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)-(7CI, 8CI) (CA INDEX NAME)



=> fil reg  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
160.33	4744.77

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-23.36	-352.59

CA SUBSCRIBER PRICE

FILE 'REGISTRY' ENTERED AT 20:37:41 ON 12 JAN 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8


TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>  
Uploading C:\Program Files\Stnexp\Queries\10718532.str



chain nodes :

1 13 23 25 26 28 29 30 31 32 33 35

ring nodes :

2 3 4 5 6 7 8 9 10 11 12 16 17 18 19 20 21 22

chain bonds :

1-4 8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31  
22-35

ring bonds :

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
18-19 19-20 20-21 21-22

exact/norm bonds :

8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31

exact bonds :

1-4 22-35

normalized bonds :

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
18-19 19-20 20-21 21-22

G1:N,X

G2:X,Ak,H


Match level :

1:CLASS 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom  
11:Atom 12:Atom 13:CLASS 14:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom  
21:Atom 22:Atom 23:CLASS 25:CLASS 26:CLASS 28:CLASS 29:CLASS 30:CLASS  
31:CLASS 32:CLASS 33:CLASS 35:CLASS

L61 STRUCTURE UPLOADED

=> d query

L61 STR



G1 N,X

G2 X,Ak,H

Structure attributes must be viewed using STN Express query preparation.

=> s l61

SAMPLE SEARCH INITIATED 20:38:32 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 65 TO ITERATE

100.0% PROCESSED 65 ITERATIONS  
SEARCH TIME: 00.00.01

3 ANSWERS

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 817 TO 1783  
PROJECTED ANSWERS: 3 TO 163

L62 3 SEA SSS SAM L61



=> s l61 full  
FULL SEARCH INITIATED 20:38:37 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 1235 TO ITERATE

100.0% PROCESSED 1235 ITERATIONS  
SEARCH TIME: 00.00.01

27 ANSWERS

L63 27 SEA SSS FUL L61

=> fil caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
161.76	4906.53

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-352.59

CA SUBSCRIBER PRICE

FILE 'CAPLUS' ENTERED AT 20:38:41 ON 12 JAN 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)


This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l63  
L64 30 L63

=> d l65 1-30 abs ibib hitstr  
L65 NOT FOUND

The L-number entered has not been defined in this session, or it has been deleted. To see the L-numbers currently defined in this session, enter DISPLAY HISTORY at an arrow prompt (=>).

=> d l64 1-30 abs ibib hitstr



AB This invention pertains to a method for producing bis(fluorine-containing phthalic anhydride) derivs. I [wherein m = 1-3; n = 1-3; Z = a single bond or a spacer], which comprises reacting a fluorine-containing dinitrile with a fluorine-containing hydroquinone, followed by hydrolysis and dehydration reaction. For example, 3,4,5,6-tetrafluorophthalonitrile was reacted with tetrafluorohydroquinone in Me iso-Bu ketone in the presence of KF to give the bisphthalonitrile (97%). The bisphthalonitrile was hydrolyzed in propionic acid in the presence of H<sub>2</sub>SO<sub>4</sub> to afford the bis(phthalic acid) (73%). The bis(phthalic acid) was treated with acetic anhydride to provide the bis(phthalic anhydride) (98%).

ACCESSION NUMBER: 2004:1127358 CAPLUS  
TITLE: Process for preparation of bis(fluorine-containing phthalic anhydride)  
INVENTOR(S): Masuda, Go; Okumura, Yasunori; Nishimae, Shinji; Nacaya, Toshihiro  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: PCT Int. Appl., 39 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004111026	A1	20041223	WO 2004-JP8829	20040617
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OH, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:		JP 2003-172046	A	20030617
		JP 2003-201123	A	20030724

IT 143363-92-2P 143376-50-5P  
RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic)

AB The compns. containing repeating units of AD(R) [A = F-substituted (O- or S-containing) hydrocarbylene; D = F-substituted (O- or S-containing) trivalent hydrocarbon group; R = Si(OR<sub>1</sub>)(OR<sub>2</sub>)R<sub>3</sub>; R<sub>1</sub>, R<sub>2</sub> = hydrocarbyl (each b.p. of R<sub>1</sub>OH and R<sub>2</sub>OH under normal pressure ≤250°); R<sub>3</sub> = (F-substituted) double bond-terminated hydrocarbyl] show n and dielec. constant of their cured products 1.350-1.600 and 2.00-4.00, resp. The compns. give cured products with adjustable n and dielec. constant, and improved mech. properties and solvent resistance.

ACCESSION NUMBER: 2004:411677 CAPLUS  
DOCUMENT NUMBER: 140:431126  
TITLE: Manufacture of silicon-containing curable polymer compositions for planar optical waveguides and wiring boards  
INVENTOR(S): Florence, Corey Nawarage  
PATENT ASSIGNEE(S): Fujitsu Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 18 pp.  
CODEN: JIXXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004143280	A2	20040520	JP 2002-309280	20021024
PRIORITY APPLN. INFO.:		JP 2002-309280		20021024

IT 691906-05-5DP, reaction products with allyltrichlorosilane and ethanol  
RL: DEV (Device component use); IMF (Industrial manufacture); PREP (Preparation); USES (Uses)  
(crosslinked; manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)


RN 691906-05-5 CAPLUS  
CN 1,3-Benzenedicarbonyl dichloride, 5-bromo-, polymer with 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI) (CA INDEX NAME)

CH 1


CRN 691906-04-4  
CMF C20 H4 C12 F8 O4



L64 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
preparation); PREP (Preparation); RACT (Reactant or reagent)  
(intermediate; prepn. of bis(fluorine-contg. phthalic anhydride))  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[2,3,5,6-tetrafluoro-1,4-phenylene]bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT


CH 2

CRN 83558-87-6  
CMF C15 H12 F6 N2 O2




CH 3


CRN 57863-69-1  
CMF C8 H3 Br Cl2 O2




IT 691906-02-2P 691906-03-3P 691906-04-4P  
691906-05-5P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
(manufacture of silicon-containing curable polymer compns. for planar optical waveguides and wiring boards)  
RN 691906-02-2 CAPLUS  
CN Benzonitrile, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)



L64 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 RN 691906-03-3 CAPLUS  
 CN Benzoic acid, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI)  
 (CA INDEX NAME)




RN 691906-04-4 CAPLUS  
 CN Benzoyl chloride, 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI)  
 (CA INDEX NAME)



RN 691906-05-5 CAPLUS  
 CN 1,3-Benzenedicarbonyl dichloride, 5-bromo-, polymer with 4,4'-[1,4-phenylenebis(oxy)]bis[2,3,5,6-tetrafluorobenzoyl chloride] and 4,4'-[2,2,2-trifluoro-1-(trifluoromethyl)ethylidene]bis[2-aminophenol] (9CI)  
 (CA INDEX NAME)


CH 1

CRM 691906-04-4  
 CMF C20 H4 Cl2 F8 O4



CH 2

L64 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI



AB The title phthalic acid derivs. (Markush structure given) are prepared by reaction of tetrafluorophthalic acid derivs. with NAM [A = divalent organic moiety; H = H, et.]. Thus, I was prepared I is a raw material for the manufacture of fluorinated polyimides. The title phthalic acid derivs. are


useful as intermediates for optical materials, liquid crystals, etc.  
 ACCESSION NUMBER: 2003:823334 CAPLUS  
 DOCUMENT NUMBER: 139:330642  
 TITLE: Phthalic acid derivatives as intermediates for halogenated polyimides, liquid crystals, etc., and process for manufacturing them  
 INVENTOR(S): Okumura, Yasunori; Kuwahara, Masayoshi; Masuda, Takeshi  
 PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokyo Koho, 5 pp.  
 CODEN: JIOKAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003300934	A2	20031021	JP 2002-108095	20020410
PRIORITY APPLN. INFO.: JP 2002-108095 20020410				

OTHER SOURCE(S): MARPAT 139:330642  
 IT 615263-86-0P  
 RL: IMP (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
 (phthalic acid derivs. as intermediates for halogenated polyimides, optical materials, liquid crystals, and process for manufacturing them)  
 RN 615263-86-0 CAPLUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro-, tetraethyl ester (9CI)  
 INDEX NAME)


L64 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)

CRM 83558-87-6  
 CMF C15 H12 F6 N2 O2





CH 3

CRM 57863-69-1  
 CMF C8 H3 Br Cl2 O2



L64 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)






AB The compds. are prepared by hydrolysis of aromatic tetranitriles I (X, X' = halo; n, n' = 1-3; Z = single bond, O, S, O-p-C6H4-mYmO, etc; Y = halo; m = 1-4) in the presence of acids in organic solvents. 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene was treated with H2SO4 in propionic acid under reflux for 6 h to give 95.7% 1,4-bis(3,4-dicarboxytrifluorophenoxy)tetrafluorobenzene.

ACCESSION NUMBER: 2002:886125 CAPIUS  
DOCUMENT NUMBER: 137:384650  
TITLE: Preparation of halogen-containing aromatic carboxylic acids  
INVENTOR(S): Kuwahara, Masayoshi; Yokoo, Junko; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002332253	A2	20021122	JP 2001-142027	20010511
JP 3490407	B2	20040126		

PRIORITY APPLN. INFO.: JP 2001-142027 20010511


OTHER SOURCE(S): MARPAT 137:384650  
IT 143363-92-2P 474805-31-7P  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143363-92-2 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



RN 474805-31-7 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]




IT 143376-50-5, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)  
RN 143376-50-5 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



IT 474805-29-3P, 1,4-Bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of halogen-containing aromatic carboxylic acids)

L64 ANSWER 4 OF 30 CAPIUS COPYRIGHT 2005 ACS ON STN (Continued)  
RN 474805-29-3 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]



L64 ANSWER 5 OF 30 CAPIUS COPYRIGHT 2005 ACS ON STN  
AB The present invention relates to halogen-containing aromatic compounds and methods thereof. The present invention relates to a halogen-containing aromatic dianhydride, halogen-containing aromatic tetranitrile compound, halogen-containing m-phenylenediamine compound and fluorine compound, and a method thereof. Tetrafluorophthalonitrile was reacted with tetrachlorohydroquinone in the presence of potassium fluoride and acetonitrile to give 1,4-bis(3,4-dicyanotrifluorophenoxy)tetrafluorobenzene.


ACCESSION NUMBER: 2002:866679 CAPIUS  
DOCUMENT NUMBER: 137:354701  
TITLE: Halogen-containing aromatic compound  
INVENTOR(S): Kuwabara, Masayoshi; Okumura, Yasunori  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan; Nippon Catalytic Chem. Ind.  
SOURCE: Eur. Pat. Appl., 38 pp.  
CODEN: EPXXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1256564	A1	20021113	EP 2002-253088	20020501
EP 1256564	B1	20040728		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
JP 2002332268 A2 20021122 JP 2001-142028 20010511  
JP 2002332281 A2 20021122 JP 2001-142029 20010511  
JP 2002332264 A2 20021122 JP 2001-142031 20010511  
JP 3563040 B2 20040908  
JP 2002332254 A2 20021122 JP 2001-142032 20010511  
US 2003018204 A1 20030123 US 2002-133158 20020426  
EP 1462436 A1 20040929 EP 2004-10371 20020501  
R: DE, FR, GB  
CN 1385427 A 20021218 CN 2002-119176 20020513  
JP 2001-142028 A 20010511  
PRIORITY APPLN. INFO.: JP 2001-142029 A 20010511  
JP 2001-142031 A 20010511  
JP 2001-142032 A 20010511  
EP 2002-253088 A3 20020501

OTHER SOURCE(S): MARPAT 137:354701  
IT 474805-29-3P 474805-31-7P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
(halogen-containing aromatic compound)  
RN 474805-29-3 CAPIUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)]





AB The compns., useful for elec. insulating materials, contain (p-C6F4COC6H4-qQORlm)n (R1 = C6H4-rX'r(CO-p-C6F4OR2)pO; R2 = divalent aryl; X, X' = halo, lower alkyl, alkoxy; q, r = 0-4; m, p = 0, 1). Thus, 2,2-bis(4-hydroxyphenyl)-1,1,1,3,3,3-hexafluoropropane was polymerized with 4,4'-bis(2,3,4,5,6-pentafluorobenzoyl)diphenyl ether to give a polymer showing dielec. constant 3.06 at 25°, 10% weight loss temperature 524° under N<sub>2</sub> and T<sub>g</sub> 174°.

ACCESSION NUMBER: 2001:124290 CAPLUS  
DOCUMENT NUMBER: 134:179345  
TITLE: Low dielectric fluorinated aromatic polyether ketone compositions with good heat resistance  
INVENTOR(S): Yasunori Kimura, Kunio Yamashita, Yoshihiko Okumura,  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 21 pp.  
CODEN: JNOKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001049110	A2	20010220	JP 1999-226981	19990810
JP 3539897	B2	20040707		

PRIORITY APPLN. INFO.: JP 1999-226981 19990810

IT 213693-15-3P



RL: IMF (Industrial manufacture); PRP (Properties); TEM (Technical or engineered material use); PREP (Preparation); USES (Uses)  
(low dielec. fluorinated aromatic polyether ketone compns. with good heat

resistance)

RN 213693-15-3 CAPLUS

CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylenecarbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)) (9CI) (CA INDEX NAME)

PAGE 1-A



AB Ketonic di-Ph ethers of p-R1COC6H4OC6H4R2-p type compds. (R1 = 2,3,4,5,6-pentafluorobenzoyl; R2 = OH, pentafluorobenzoyl group) and polyether-polyketone polymers containing tetrafluorophenylene and phenylene groups are provided which have good mech. strength, toughness, elec. property, thermal oxidative stability and solubility Thus, heating 0.5 g 2,3,4,5,6-pentafluoro-4'-hydroxybenzophenone with 0.36 ground K carbonate, 2 mL N-methyl-2-pyrrolidone and 1 mL PhMe at 160° while stirring for 3 h gave a polymer at 85% yield and having viscosity 0.5 g/dL in AcNMe<sub>2</sub>.

ACCESSION NUMBER: 2001:25791 CAPLUS  
DOCUMENT NUMBER: 134:86663  
TITLE: (2,3,4,5,6-Pentafluorobenzoyl)diphenyl ether compound,  
and fluorine-containing aryl ether ketone polymer  
INVENTOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.;  
Fitch, John W., III; Reddy, V. Sreenivasulu  
PATENT ASSIGNEE(S): Nippon Shokubai Co., Ltd., Japan  
SOURCE: U.S., 22 pp., Cont.-in-part of U.S. Ser. No. 106,270, abandoned.  
CODEN: USXXAM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6172181 B1		20010109	US 1999-354976	19990716
			US 1998-106270	19980629

PRIORITY APPLN. INFO.: MARPAT 134:86663

OTHER SOURCE(S):

IT 213693-15-3P


RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(manufacture of (2,3,4,5,6-pentafluorobenzoyl)diphenyl ether compound

and


RN 213693-15-3 CAPLUS

CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylenecarbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)) (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT


L64 ANSWER 10 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Fluoropolymer-polyether-polyketones were obtained by homopolycondensation of 4-hydroxy-4'-(pentafluorobenzoyl)diphenyl ether and by copolycondensation of 4,4'-bis(pentafluorobenzoyl)diphenyl ether with benzenediols or bisphenols. The polymers have very good heat resistance, with 10% weight loss temps. >500°.

ACCESSION NUMBER: 1998:532324 CAPLUS  
DOCUMENT NUMBER: 129:260978  
TITLE: New polymers derived from 2,3,4,5,6-pentafluorobenzoic acid  
AUTHOR(S): Kimura, Kunio; Yamashita, Yuhiko; Cassidy, Patrick E.; Fitch, John W., III; Reddy, V. Sreenivasulu; Sakaguchi, Yoshinobu  
CORPORATE SOURCE: Faculty of Environmental Science and Technology, Okayama University, Okayama, 700-8530, Japan  
SOURCE: Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1998), 39(2), 790-791  
PUBLISHER: American Chemical Society, Division of Polymer Chemistry  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 213693-15-3P, 4,4'-Bis(pentafluorobenzoyl)diphenyl ether-hydroquinone copolymer, SRU  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of thermally stable fluoropolymer-polyether-polyketones)  
RN 213693-15-3 CAPLUS  
CN Poly(oxy-1,4-phenyleneoxy(2,3,5,6-tetrafluoro-1,4-phenylene)carbonyl-1,4-phenyleneoxy-1,4-phenylene)carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene) (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L64 ANSWER 11 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN  
AB Selected perfluorinated tertiary alcs. were reacted with p-toluenesulfonyl chloride to form their p-toluenesulfonyl esters C6F5(CF3)2COSO2C6H4CH3 (I), CF3C6F4OC6F4(C3F7)(C8F17)COSO2C6H4CH3 and (CF3C6F4OC6F4)2(C7F15)COSO2C6H4CH3. The absolute configuration of I is established by X-ray diffraction.

ACCESSION NUMBER: 1998:210377 CAPLUS  
DOCUMENT NUMBER: 128:321424  
TITLE: p-Toluenesulfonyl esters of perfluorinated tertiary alcohols: crystal structure determination of the absolute configuration of C6F5(CF3)2COSO2C6H4CH3  
AUTHOR(S): Krumm, Burkhard; Vij, Ashwani; Kirchmeier, Robert L.; Shreeve, Jean'ne M.  
CORPORATE SOURCE: Dep. of Chem., Univ. of Idaho, Moscow, ID, 83844-2343, USA  
SOURCE: Journal of Fluorine Chemistry (1998), 89(1), 19-22  
CODEN: JFLICAR; ISSN: 0022-1139  
PUBLISHER: Elsevier Science S.A.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 128:321424  
IT 207233-54-3  
RL: RCT (Reactant); RACT (Reactant or reagent) (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal structure determination of the absolute configuration of C6F5(CF3)2COSO2C6H4CH3)  
RN 207233-54-3 CAPLUS  
CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentafluorooheptyl)- $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy)phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy)phenoxy]- (9CI) (CA INDEX NAME)



IT 207233-53-2P


RL: SPN (Synthetic preparation); PREP (Preparation)  
 (p-Toluenesulfonyl esters of perfluorinated tertiary alcs.: crystal structure determination of the absolute configuration of

C6F5(CF3)2COSO2C6H4CH3


RN 207233-53-2 CAPLUS

CN Benzenemethanol, 2,3,5,6-tetrafluoro- $\alpha$ -(pentadecafluoroheptyl)- $\alpha$ -(2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]phenyl)-4-[2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]-, 4-methylbenzenesulfonate (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT

L64 ANSWER 12 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN

AB Reactions of 4'-CF3C6F4OC6F4Li, generated in situ, with elements of group 16 (S, Se, Te) lead to CF3C6F4OC6F4SH (2), (CF3C6F4OC6F4Se)2 (3), and (CF3C6F4OC6F4Te)2 (4)/(CF3C6F4OC6F4)2Te (4a). The phenol derivative CF3C6F4OC6F4OH (1) is obtained by reaction of CF3C6F4OC6F4Li with B(OMe)3/H2O2. The reaction of CF3C6F4OC6F4Li with trimethylsilyl chloride

or trimethyltin chloride gives CF3C6F4OC6F4XMe3 (X = Si (5), Sn (6)). Oxidation of 2 in the presence of bromine results in the formation of (CF3C6F4OC6F4S)2 (7) and CF3C6F4OC6F4SO2Br (8). Mixed perfluoroaryloxo/thio ethers CF3C6F4OC6F4SC6F4R (R = NO2 (9), CN (10),

CF3 (11)) and CF3C6F4OC6F4SC5F4N (12) are obtained upon reaction of 2 with excess C6F5R and pentafluoropyridine in the presence of K2CO3. With 4-C6F5OC6F4NO2, a mixture of (2-CF3C6F4OC6F4S) (4-C6F5O)C6F3NO2 (13) and

9 is formed. Reaction of excess 2 with C6F5R gives the 2,4,6-substituted benzenes (CF3C6F4OC6F4S)3C6F2R (R = NO2 (14), CN (15)). The trimethylsilyl ether CF3C6F4OC6F4OSiMe3 (16) is prepared from the

reaction of 1 with hexamethyldisilazane. 16 is a convenient reagent for the preparation

of the aryl ethers CF3C6F4OC6F4OC6F4R (R = NO2 (17), CN (18)) and CF3C6F4OC6F4OC5F4N (19) upon reaction with C6F5R and C5F5N. The

secondary alcs. CF3C6F4OC6F4CH(C6H5)OH (20) and CF3C6F4OC6F4CH(C6F5)OH (21) are synthesized by the reactions of 5 with benzaldehyde and pentafluorobenzaldehyde in the presence of tetrabutylammonium fluoride as a catalyst. In the synthesis of 21 the byproduct CF3C6F4OC6F4CH(C6F5)OC6F4CHO is also formed and isolated.

ACCESSION NUMBER: 1997:667252 CAPLUS

DOCUMENT NUMBER: 127:293323

TITLE: Synthesis and Chemistry of CF3C6F4OC6F4 Group 14/16

Derivatives

AUTHOR(S): Krumm, Burkhard; Kirchmeier, Robert L.; Shreeve, Jean'ne M.

CORPORATE SOURCE: Department of Chemistry, University of Idaho, Moscow, ID, 83844-2343, USA

SOURCE: Inorganic Chemistry (1997), 36(23), 5222-5230

CODEN: INOCAT; ISSN: 0020-1669

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 127:293323

IT 197150-21-3P 197150-22-4P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

RN 197150-21-3 CAPLUS


CN Benzene, 1,2,4,5-tetrafluoro-3-(2,3,5,6-tetrafluoro-4-nitrophenoxy)-6-[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)

L64 ANSWER 12 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)




RN 197150-22-4 CAPLUS

CN Benzonitrile, 2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]phenoxy]- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE  
 FORMAT






AB Phthalic acids I (R = CO<sub>2</sub>H; X = halo) are prepared by heating phthalonitriles I (R = cyano; X = halo) in acidic aqueous media, (A) dissolving the products in organic solvent-containing media, (B) separating the products, and (C) treating the isolated products in acidic aqueous media under heating. Then, the above processes A to C are repeated. 21 time(s). I (R = cyano, X = F) (10 g) was refluxed with aqueous H<sub>2</sub>SO<sub>4</sub> for 6 h, diluted with H<sub>2</sub>O, filtered, and the crude product was dissolved in Me<sub>2</sub>CO. The solution was treated with H<sub>2</sub>O to give 10.8 g I (R = CO<sub>2</sub>H, X = F) (II) with purity 91.8%, which was refluxed with H<sub>2</sub>SO<sub>4</sub> for 6 h and similarly treated. The process was repeated to give 9.8 g II with 99.2% purity.

ACCESSION NUMBER: 1997:377463 CAPLUS  
DOCUMENT NUMBER: 127:33992  
TITLE: Preparation of phthalic acids as intermediates for polyimides from phthalonitriles  
INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
CODEN: JIOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110785	A2	19970428	JP 1995-266007	19951013

PRIORITY APPLN. INFO.: JP 1995-266007 19951013

OTHER SOURCE(S): MARPAT 127:33992  
IT 143363-92-2P  
RL: IMF (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)  
(preparation and purification of phthalic acids as intermediates for polyimides by acid hydrolysis of phthalonitriles)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation and purification of phthalic acids as intermediates for polyimides by acid hydrolysis of phthalonitriles)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




AB Title compds. I (X = halo; Y = CO<sub>2</sub>H) are prepared by heating I (Y = cyano) in acidic aqueous media, heating the product-containing media with alkaline substances, and separating I (Y = CO<sub>2</sub>H) from the media by mixing with acidic substances. An aqueous solution containing 10 g I (X = F, Y = cyano) and H<sub>2</sub>SO<sub>4</sub> was refluxed for 6 h, mixing with H<sub>2</sub>O and NaOH (pH 11.8), refluxed for 1 h, and mixed with aqueous H<sub>2</sub>SO<sub>4</sub> to give 10.8 g I (X = F, Y = CO<sub>2</sub>H) with 99.2% purity.

ACCESSION NUMBER: 1997:377462 CAPLUS  
DOCUMENT NUMBER: 127:33986  
TITLE: Preparation of phthalic acids as intermediates for polyimides  
INVENTOR(S): Okumura, Yasunori; Yoshitoshi, Koji; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Shokubai Kagaku Kogyo Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JIOXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09110784	A2	19970428	JP 1995-266006	19951013


PRIORITY APPLN. INFO.: JP 1995-266006 19951013

OTHER SOURCE(S): MARPAT 127:33986  
IT 143363-92-2P  
RL: IMF (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); PREP (Preparation)  
(preparation of phthalic acids as intermediates for polyimides by hydrolysis of phthalonitriles and heating with alkalies)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of phthalic acids as intermediates for polyimides by hydrolysis of phthalonitriles and heating with alkalies)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)





AB The title bisphthalonitriles I (X, Y = O, S; R = Z, Z1), useful as intermediates for optical materials, circuit board materials, photosensitive materials, liquid crystals, etc., are prepared by treatment of 3,4,5,6-tetrafluorophthalonitrile (II) with HXRYH in organic solvents being slightly soluble in H<sub>2</sub>O in the presence of basic substances, followed by addition of H<sub>2</sub>O to the reaction products and separation of the organic layer. An AcOEt solution of tetrafluorohydroquinone was added dropwise to a mixture of II, KF, and AcOEt under reflux over 1 h, and the reaction mixture was further stirred under reflux for 8 h, cooled, and then washed with H<sub>2</sub>O to give 99% I (X = Y = O, R = Z).



ACCESSION NUMBER: 1997:154684 CAPLUS  
DOCUMENT NUMBER: 126:157295  
TITLE: Preparation of bis(fluorophthalonitriles) from tetrafluorophthalonitrile and fluorohydroquinone or fluorodiphenol  
INVENTOR(S): Okumura, Yasunori; Kaieda, Osamu  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JIOKAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 08333322	A2	19961217	JP 1995-143593	19950609

PRIORITY APPLN. INFO.: JP 1995-143593 19950609

OTHER SOURCE(S): CASREACT 126:157295; MARPAT 126:157295  
IT 143376-50-5P  
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP

L64 ANSWER 15 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
(Preparation)  
(prepn. of bis(fluorophthalonitriles) from tetrafluorophthalonitrile and fluorohydroquinone, diphenol, or their sulfur analog using slightly water-sol. solvents and basic substances)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



AB Polyimide optical materials comprise a perfluorinated polyimide having a perfluorinated repeating unit represented by the general formula I (R1 = a tetraivalent perfluorinated organic group; and R2 = a divalent perfluorinated organic group).


ACCESSION NUMBER: 1995:854348 CAPLUS  
DOCUMENT NUMBER: 123:354214  
TITLE: Polyimide optical material  
INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni; Yamamoto, Fumio  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corporation, Japan  
SOURCE: U.S., 26 pp. Cont.-in-part of U.S. Ser. No. 54,973, abandoned.  
CODEN: USXXAM  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5449741	A	19950912	US 1993-140982	19931025
JP 09031014	A2	19970204	JP 1996-207475	19910913
US 5233018	A	19930803	US 1991-765672	19910926
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 5750731	A	19980512	US 1996-718208	19960920
US 5049924	A	19981215	US 1998-20573	19980128
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		


PRIORITY APPLN. INFO.: JP 1990-256843 A 19900928  
JP 1991-106552 A 19910412  
JP 1991-106554 A 19910412  
JP 1991-106557 A 19910412  
US 1991-765672 A3 19910926  
US 1993-54973 B2 19930430  
JP 1991-235020 A3 19910913

L64 ANSWER 16 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
CA 1991-2052368 A3 19910926  
US 1993-140482 A3 19931025  
US 1993-140982 A3 19931025  
US 1995-451465 A1 19950526  
US 1996-718208 A3 19960920  
US 1998-20573 A3 19980128

IT 143376-50-5  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(polyimide optical materials)  
RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143363-92-2  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reg. polyimide optical materials)  
RN 143363-92-2 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)





\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. I, useful as material for hardeners for fluorinated epoxy resins (no data), is prepared via II [R = cyano, CO<sub>2</sub>H]. Thus, a mixture of

tetrafluorophthalonitrile, tetrafluorohydroquinone, and Et<sub>3</sub>N in DMF was heated at 35° for 30 to give 21% II [R = cyano], which was treated with 60% H<sub>2</sub>SO<sub>4</sub> at 150° for 5 h to give 26% II [R = CO<sub>2</sub>H], which was refluxed with Ac<sub>2</sub>O for 2 h to 52% I.

ACCESSION NUMBER: 1994:630662 CAPLUS  
DOCUMENT NUMBER: 121:230662  
TITLE: preparation of a perfluorinated hexacarboxylic acid as

material for hardeners for fluorinated epoxy resins  
INVENTOR(S): Sasaki, Shigekuni; Matsura, Tooru; Ando, Shinji  
PATENT ASSIGNEE(S): Nippon Telegraph & Telephone, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
CODEN: JKOXAF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06157501	A2	19940603	JP 1992-340986	19921130
PRIORITY APPLN. INFO.:			JP 1992-340986	19921130



OTHER SOURCE(S): CASREACT 121:230662

IT 158394-12-8P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and conversion into trianhydride)

RN 158394-12-8 CAPLUS  
CN 1,2-Benzenedicarboxylic acid, 4,5-bis[4-(3,4-dicarboxy-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)



IT 158394-11-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and hydrolysis of)  
RN 158394-11-7 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,5-bis[4-(3,4-dicyano-2,5,6-trifluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-3,6-difluoro- (9CI) (CA INDEX NAME)



AB The title derivs. I (X = O, S; Y = O, S, Q1, Q2) are prepared by treating 28 mol 3,4,5,6-tetrafluorophthalonitrile (II) with 1 mol phenols or thiophenols III in the presence of basic substances in organic solvents.

A mixture of 0.22 mol II and KF in MeCN was treated dropwise with a solution of 0.011 mol III (X = Y = O) in MeCN under reflux over 1 h, then refluxed for 4 h to give 99% I.

ACCESSION NUMBER: 1994:435030 CAPLUS  
DOCUMENT NUMBER: 121:35030  
TITLE: Preparation of fluorine-containing phthalonitrile derivatives as intermediates for fluorine-containing polyimides

INVENTOR(S): Kaieda, Osamu; Okumura, Yasunori; Ito, Hideaki  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.  
CODEN: JKOXAF

DOCUMENT TYPE: Patent  
LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 06016615	A2	19940125	JP 1992-176633	19920703
JP 08000803	B4	19960110		
PRIORITY APPLN. INFO.:			JP 1992-176633	19920703

OTHER SOURCE(S): CASREACT 121:35030; MARPAT 121:35030

IT 143376-50-5P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as intermediate for polyimides)

RN 143376-50-5 CAPLUS  
CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-





AB The title compds. (I; R1, R2 = F, C1-12 linear or branched alkoxy, C1-12 alkylthio, C1-12 alkylamino, aryloxy, arylthio, arylamino, Q - Q2; X = O, S; Y1 = CO2CO), useful as intermediates for drugs, agrochems., optical materials, printed circuit board materials, photosensitive materials, and liquid crystal materials, are prepared by heating F-containing phthalic acid derivs. (II; R1, R2, X = same as above; Y, Y1 = CO2H) in a solvent selected from SOCl2, POCl3, or AcCl at 40-105°. The process is safe and gives perfluorophthalic anhydride derivs. I of high purity in high yields. Thus, 70.5 g SOCl2 was added to 30.0 g tetrafluorophthalic acid and the resulting mixture was allowed to react at 70° for approx. 3 h to give 97% tetrafluorophthalic anhydride.

ACCESSION NUMBER: 1994:298457 CAPLUS  
DOCUMENT NUMBER: 120:298457  
TITLE: Preparation of fluorine-containing phthalic anhydride derivatives  
INVENTOR(S): Okumura, Yasunori; Ito, Hideki; Kaleda, Osamu  
PATENT ASSIGNEE(S): Nippon Catalytic Chem Ind, Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
CODEN: JKKXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328502	A2	19921117	JP 1991-125451	19910426

AB The filter, comprising a dielec. multibilayer, is characterized in that the substrate consists of a thin-film perfluoropolyimide or a blend or a copolymer thereof.


ACCESSION NUMBER: 1993:417687 CAPLUS  
DOCUMENT NUMBER: 119:17687  
TITLE: IR longpass dielectric interference filter  
INVENTOR(S): Yamamoto, Fumio; Sasaki, Shigekuni; Matsura, Toru; Ando, Shinji; Oguchi, Taisuke  
PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JKKXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 04328502	A2	19921117	JP 1991-125451	19910426

IT 148273-08-9  
RL: USES (Uses)  
(fluorinated polyimide blends containing, IR longpass filters containing)  
RN 148273-08-9 CAPLUS  
CN 1,2-benzenedicarboxylic acid, 4,4'-(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis[3,5,6-trifluoro-, polymer with 2,4,5,6-tetrafluoro-1,3-benzenediamine (9CI) (CA INDEX NAME)

CM 1



CRN 143363-92-2  
CMF C22 H4 F10 O10



CM 2

CRN 1198-63-6  
CMF C6 H4 F4 N2

OTHER SOURCE(S): CASREACT 120:298457; MARPAT 120:298457  
IT 143363-92-2  
RL: PROC (Process)  
(conversion of, into acid anhydride)  
RN 143363-92-2 CAPLUS  
CN 1,2-benzenedicarboxylic acid, 4,4'-(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




L64 ANSWER 23 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB A perfluorinated polyimide that has glass temperature >300° and high optical transparency over the entire optical communication wavelengths was synthesized. The high thermal stability and optical transparency in near-IR region are due to the fully aromatic mol. structure and the absence of C-H bonds. The use of a diamine with relatively high reactivity and a new perfluorinated dianhydride, which has flexible structure, makes it possible to obtain a tough and flexible perfluorinated polyimide film.

In addition, this polymer has a low dielec. constant Perfluorinated polyimides are promising for use as optical-electronic materials.


ACCESSION NUMBER: 1992:592808 CAPIUS  
 DOCUMENT NUMBER: 117:192808  
 TITLE: Perfluorinated polyimide synthesis  
 AUTHOR(S): Ando, Shinji; Matsuura, Tohru; Sasaki, Shigekuni  
 CORPORATE SOURCE: Interdiscip. Res. Lab., NTT, Musashino, 180, Japan  
 SOURCE: Macromolecules (1992), 25(21), 5858-60  
 CODEN: MAMOBX; ISSN: 0024-9297  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 143376-50-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and hydrolysis of)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)




L64 ANSWER 24 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN (Continued)  
 US 1998-20573 A3 19980128

IT 143376-50-5P  
 RL: RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
 (preparation and hydrolysis of)  
 RN 143376-50-5 CAPIUS  
 CN 1,2-Benzenedicarbonitrile, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



IT 143363-92-2P  
 RL: PREP (Preparation)  
 (preparation of)  
 RN 143363-92-2 CAPIUS  
 CN 1,2-Benzenedicarboxylic acid, 4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[3,5,6-trifluoro- (9CI) (CA INDEX NAME)



L64 ANSWER 24 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The title polymers, useful as waveguides in optoelectronic integrated circuits or optoelec. boards, are prepared from perfluoro tetracarboxylic acid derivs. and perfluoro diamines, the preparation of which is described.

Polymerization of 4,4'-[(tetrafluoro-p-phenylene)bis(3,5,6-trifluorophthalic anhydride)] with tetrafluoro-1,3-phenylenediamine gave a polymer with no substantial absorption at optical communication wavelengths.

ACCESSION NUMBER: 1992:531752 CAPIUS  
 DOCUMENT NUMBER: 117:131752  
 TITLE: Monomers for preparation of perfluorinated polyamic acids and polyimides  
 INVENTOR(S): Ando, Shinji; Matsuura, Toru; Sasaki, Shigekuni; Yamamoto, Fumio  
 PATENT ASSIGNEE(S): Nippon Telegraph and Telephone Corp., Japan  
 SOURCE: Eur. Pat. Appl., 39 pp.  
 CODEN: EPOKDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 480266	A2	19920415	EP 1991-116553	19910927
EP 480266	A3	19930901		
EP 480266	B1	19960626		
R: DE, FR, GB				
JP 05001148	A2	19930108	JP 1991-235020	19910913
JP 2851019	B2	19990127		
JP 09031014	A2	19970204	JP 1996-207475	19910913
CA 2052368	AA	19920329	CA 1991-2052368	19910926
CA 2052368	C	19980915		
CA 2199703	C	20001031	CA 1991-2199703	19910926
US 6048986	A	20000411	US 1998-98605	19980617
JP 11147955	A2	19990602	JP 1998-251741	19980803
JP 3085666	B2	20000911		
PRIORITY APPLN. INFO.:				
			JP 1990-256843	A 19900928
			JP 1991-106552	A 19910412
			JP 1991-106554	A 19910412
			JP 1991-106557	A 19910412
			JP 1991-235020	A3 19910913
			CA 1991-2052368	A3 19910926
			US 1991-765672	A3 19910926
			US 1993-54973	B2 19930430
			US 1993-140482	A3 19931025
			US 1995-451465	B1 19950526
			US 1996-718208	A3 19960920

L64 ANSWER 25 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN  
 AB The title polymers, e.g., polyether-polyketones, polyarylates, polyimides, and polyesters, have good heat resistance and low attenuation and are useful as optical transmitting systems, e.g., for controlling ignition timing and fuel metering systems for internal combustion engines in automobiles. Thus, an optical fiber comprised a core of amorphous PEEK and a sheath of poly(2,2,2-trifluoroethyl methacrylate).

ACCESSION NUMBER: 1992:130739 CAPIUS  
 DOCUMENT NUMBER: 116:130739  
 TITLE: Amorphous polymers for optical transmitting systems and optical members and their use  
 INVENTOR(S): Takezawa, Yoshitake; Ohara, Shuichi; Tanno, Seikichi; Taketani, Noriaki; Shimura, Masato  
 PATENT ASSIGNEE(S): Hitachi, Ltd., Japan  
 SOURCE: Eur. Pat. Appl., 31 pp.  
 CODEN: EPOKDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 454165	A2	19911030	EP 1991-106851	19910426
EP 454165	A3	19930120		
R: DE, FR, GB, IT, NL				
JP 04009805	A2	19920114	JP 1990-112511	19900427
JP 2854669	B2	19990203		
US 5093888	A	19920303	US 1991-686997	19910418
PRIORITY APPLN. INFO.:				
			JP 1990-112511	A 19900427

IT 138687-03-3  
 RL: USES (Uses)  
 (optical fibers, heat-resistant, for engine control systems)  
 RN 138687-03-3 CAPIUS  
 CN Poly[oxy(2,6-dimethyl-1,4-phenylene)oxy(2,3,5,6-tetrafluoro-1,4-phenylene)]carbonyl(2,3,5,6-tetrafluoro-1,4-phenylene)] (9CI) (CA INDEX NAME)



L64 ANSWER 26 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The PbO<sub>2</sub> oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).

ACCESSION NUMBER: 1974:520135 CAPIUS  
DOCUMENT NUMBER: 81:120135  
TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol  
AUTHOR(S): Denivelle, Leon; Huynh Anh Hoa  
CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conservatoire Natl. Arts

SOURCE: Metiers, Paris, Fr.  
Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 487-90

CODEN: BSCFAS; ISSN: 0037-8968


DOCUMENT TYPE: Journal  
LANGUAGE: French

IT 53359-93-6P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 53359-93-6 CAPIUS

CN Phenol, 2,3,5,6-tetrafluoro-4-[2,3,5,6-tetrafluoro-4-(pentafluorophenoxy)phenoxy]- (9CI) (CA INDEX NAME)



L64 ANSWER 27 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN

GI For diagram(s), see printed CA Issue.

AB The controlled oxidation of perchloro p-bromophenols (I), (II), and (III) gives

the corresponding perchloro polyethers (IV); the perchloro 4-aryloxycyclohexa-2,5-dienone (V) is obtained from pentachlorophenol (VI). Similarly, VII gives VIII and 4-chloro-2,3,5,6-tetrabromophenol gives IX.

ACCESSION NUMBER: 1971:111698 CAPIUS  
DOCUMENT NUMBER: 74:111698  
TITLE: Oxidation of perchloro benzenic phenols  
AUTHOR(S): Denivelle, Leon; Lampel, Alfred  
CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conserv. Natl. Aris

SOURCE: Metiers, Paris, Fr.  
Comptes Rendus des Seances de l'Academie des Sciences,

Serie C: Sciences Chimiques (1971), 272(7), 653-6

CODEN: CHDCAQ; ISSN: 0567-6541


DOCUMENT TYPE: Journal  
LANGUAGE: French

IT 31404-37-2P 31404-38-3P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)


RN 31404-37-2 CAPIUS

CN 2,5-Cyclohexadien-1-one, 4-bromo-4-[4-(4-bromo-2,3,5,6-tetrafluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



RN 31404-38-3 CAPIUS

CN Phenol, 4-[4-(4-bromo-2,3,5,6-tetrafluorophenoxy)-2,3,5,6-tetrafluorophenoxy]-2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)



L64 ANSWER 27 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN

(Continued)

L64 ANSWER 28 OF 30 CAPIUS COPYRIGHT 2005 ACS on STN

AB A number of perfluorinated and partially fluorinated phenyl and polyphenyl

ethers were synthesized, characterized for thermal stability, fire resistance, and viscosity, and compared with their H analogs to assess

the potential use of this class of compds. as functional fluids. Without exception, polyfluorination and perfluorination lower thermal stability; the decrease in stability depends on the position and number of fluorine substituents. The autoignition temperature and fire resistance are not

improved over the H analog, and viscosity is degraded. These data coupled with the

comparatively high melting points do not suggest a bright future for this class of compds. as useful functional fluids. 6 references.

ACCESSION NUMBER: 1968:77890 CAPIUS

DOCUMENT NUMBER: 68:77890

TITLE: Synthesis, thermal stability, flammability, and viscosity of some partially fluorinated and perfluorinated aromatic and polyaromatic ethers

AUTHOR(S): Richardson, George Albert; Blake, Edward S.

CORPORATE SOURCE: Monsanto Res. Corp., Dayton, OH, USA

SOURCE: Industrial & Engineering Chemistry Product Research and Development (1968), 7(1), 17-21

CODEN: IEPRA6; ISSN: 0196-4321


DOCUMENT TYPE: Journal  
LANGUAGE: English

IT 6804-37-1P

RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)


RN 6804-37-1 CAPIUS

CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)- (7CI, 8CI) (CA INDEX NAME)



L64 ANSWER 29 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN  
GI For diagram(s), see printed CA Issue.  
AB 2,3,5,6-Tetrafluoro-O,O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]hydroquinones (I) and 2,4,5,6-tetrafluoro-O'-bis[4-(X-substituted)-2,3,5,6-tetrafluorophenyl]resorcinols (II) are prepared by the treatment of the compds. of the general formulas C6F5X, where X is CN, CF3, CONH2, and CO2H, with 2,3,5,6-tetrafluorohydroquinone di-Na salt and 2,4,5,6-tetrafluororesorcinol di-Li salt (III), resp. 19F-N.M.R. and ir data for the prepared I and II are given. III is treated with decafluorocyclohexene to give a mixture of 2,4,5,6-tetrafluoro-O,O'-bis(perfluoro-1-cyclohexen-1-yl)resorcinol and 2,4,5,6-tetrafluoro-O-(perfluoro-1-cyclohexen-1-yl)-O'-(perfluoro-2-cyclohexen-1-yl)resorcinol.


ACCESSION NUMBER: 1968:68571 CAPLUS  
DOCUMENT NUMBER: 68:68571  
TITLE: Reactions of perfluorophenolates with substituted pentafluorobenzenes and perfluorocyclohexene  
AUTHOR(S): De Pasquale, Ralph J.; Tamborski, Christ  
CORPORATE SOURCE: Wright-Patterson Air Force Base, Dayton, OH, USA  
SOURCE: Journal of Organic Chemistry (1968), 33(2), 830-3  
CODEN: JOCEAH; ISSN: 0022-3263  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
IT 14796-02-2P 14796-03-3P 15053-71-1P  
15077-30-2P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 14796-02-2 CAPLUS  
CN Benzanilide, 4,4'-[(2,3,5,6-tetrafluoro-p-phenylene)dioxy]bis[2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)]




RN 14796-03-3 CAPLUS  
CN Benzoic acid, 4,4'-[(2,3,5,6-tetrafluoro-p-phenylene)dioxy]bis[2,3,5,6-tetrafluoro- (8CI) (CA INDEX NAME)]

L64 ANSWER 30 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN  
AB A mixture of 26.5 g. C6F6, 26.5 g. 85% KOH, and 75 ml. H2O was heated in a sealed bomb at 175° for 5 hrs. with agitation to give 33.1 g. C6F5OH, b. 144-5°. A mixture of 67 g. C6F5H, 21.6 g. KOH, 150 ml. pyridine, and 2 ml. H2O was refluxed 1 hr., treated with 21.6 KOH, and refluxed 24 hrs. to give 21 g. 2,3,5,6-tetrafluorophenol, b20 47°. Similarly refluxing 36 g. C6F5Me and 28 g. KOH in 300 ml. tert-BuOH gave 12 g. 2,3,5,6-tetrafluoro-p-cresol, m. 52°. Reaction of 20 g. C6F5I with 2 g. KOH and 1 ml. H2O in 100 ml. pyridine gave only one product, 2,3,5,6-tetrafluoro-4-iodophenol, m. 79-81°, benzoate m. 59-60.2°. Reaction of 66 g. C6F5Br with 28.5 g. KOH and 1 ml. H2O in 150 ml. pyridine, however, gave a mixture of products: 3.5 g. 2-bromo-3,4,5,6-tetrafluorophenol, m. 41-3° (3,5-dinitrobenzoate, m. 104-5°); and 11.5 g. 4-bromo-2,3,5,6-tetrafluorophenol (3,5-dinitrobenzoate m. 131-3°). Similarly, 100 g. 2-chlorotetrafluoro-o,a,a-trifluorotoluene on treatment with 5.6 g. KOH and 1 ml. H2O in 100 ml. pyridine gave 2.5 g. 2-chlorotrifluoro-o,a,a-trifluoro-o-cresol, b15 92-3°, n24D 1.4510; and 15 g. 2-chlorotrifluoro-o,a,a-trifluoro-p-cresol, b15 102-3°, n24D 1.4510. The reactions of polyfluorobenzenes were then studied with alkoxides. A solution of 123.5 g. C6F5Br in 70 ml. pyridine was treated with a solution of 11.5 g. Na in 150 ml. MeOH during 1.5 hrs. and the mixture refluxed 15 hrs. and acidified with 1 l. 10% HCl to give 66 g. 4-bromo-2,3,5,6-tetrafluoroanisole, b5 79-81°, n25D 1.4812. Similarly, a mixture of 10 g. C6F5I in 50 ml. pyridine and 0.8 g. Na in 150 ml. MeOH on refluxing for 3 hrs. gave 1.5 g. unchanged C6F5I and 5.5 g. 2,3,5,6-tetrafluoro-4-iodoanisole (I), b20 113-15°, n22D 1.5229. Refluxing 1 g. I with 1 g. activated Cu powder for 12 min. gave 0.2 g. octafluoro-4,4'-dimethoxybiphenyl, m. 90-1.2°. To a cold solution of 9 g. Na in 250 ml. PhCH2OH was added 75 g. C6F6 and the mixture refluxed 24 hrs. to give 30 g. benzyl pentafluorophenyl ether (II), m. 44°. A better yield was obtained when a solution of 4.6 g. Na and 22 g. PhCH2OH in 250 ml. tert-BuOH was refluxed with 40 g. C6F6 for 40 hrs. to give 33 g. III. A solution of 8 g. C6F6 in 30 ml. HCONMe2 was treated with 5.28 g. PhOK and the mixture refluxed 0.5 hr. to give 1 g. 2,3,5,6-tetrafluoro-1,4-diphenoxybenzene, m. 147-9°, and 3.5 g. 2,3,4,5,6-pentafluorophenyl phenyl ether, (III), m. 29°. III was also obtained by heating a mixture of 11 g. C6F5OK, 15 g. PhBr, and 1 g. Cu at 210° in a sealed bomb. A solution of 6 g. C6F5OK and 12.8 g. C6F6 in 30 ml. HCONMe2 was refluxed 14 hrs. to give 1.5 g. bis(perfluorophenyl) ether, m. 67-9°, and a second product, m. 145-8°, probably p-bis(pentafluorophenoxy)2,3,5,6-tetrafluorobenzene. Similarly, a solution of 0.6 g. Na in 50 ml. EtOH refluxed with 5.3 g. C6F5NMe2 2 hrs. gave 5.1 g. 4-ethoxy-2,3,5,6-tetrafluoro-N,N-dimethylaniline, b. 34°. Reactions with amines were next investigated. A mixture of 280 g. C6F6 and 400 ml. 28% aqueous NH3 was rocked in a sealed bomb for 2 hrs. at 235° to give 236 g. C6F5NH2, m. 34°, and 28 g. tetrafluorophenylenediamine (sublimed 75°/1 mm.) shown by its nuclear magnetic resonance spectrum to be essentially the meta isomer mixed with a small amount of the para isomer. Similarly, heating a mixture of


L64 ANSWER 29 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)



RN 15053-71-1 CAPLUS  
CN Benzonitrile,  
4,4'-[(2,3,5,6-tetrafluoro-1,4-phenylene)bis(oxy)]bis[2,3,5,6-tetrafluoro- (9CI) (CA INDEX NAME)]



RN 15077-30-2 CAPLUS  
CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis[2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenoxy]- (9CI) (CA INDEX NAME)]




L64 ANSWER 30 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
56 g. C6F6 and 110 ml. 30% aq. MeNH2 at 220° for 3 hrs. gave 59% C6F5NHMe, b. 170-2°, and 25% 2,3,5,6-tetrafluoro-N,N'-diphenylphenylenediamine, m. 94°. The reaction product obtained by heating 50 g. C6F6 and 110 ml. 25% aq. Me2NH at 235° for 1 hr. was distd. at 1 mm. pressure and five fractions were collected. The first fraction (65%), b1 88°, was C6F5NMe2. Fraction 2, b1 88-126°, was shown by vapor phase chromatography to be C6F5NMe2 with 3 other compds. Fraction 3, b1 126-134°, consisted of 3 isomers of bis(dimethylamino)tetrafluorobenzene with the meta-isomer predominating. Fraction 4, b1 134-40°, contained equal amts. of the meta and para isomers. Fraction 5, b1 140-8°, was pure para isomer. The meta and para isomers could be sepd. by vapor phase chromatography. Similarly, heating a mixt. of 30 g. C6F5Br and 70 ml. NH4OH at 200° for 2 hrs. gave 22 g. p-bromotetrafluoroaniline, m. 61°. Heating 16 g. C6F5I and 30 ml. 8% NH4OH at 165° for 2 hrs. gave 7.6 g. tetrafluoro-p-iodoaniline, m. 77°. Benzyl pentafluorophenyl ether (20 g.) was heated with large excess of 28% NH4OH to give 3 g. p-(benzyloxy)tetrafluoroaniline, m. 97°. Similarly, heating 50 g. 2-chlorotetrafluoro-o,a,a-trifluorotoluene and 120 ml. 28% NH4OH at 21° for 2 hrs. gave 22 g. 2-chlorotrifluoro-o,a,a-trifluoro-p-toluidine, which decompd. readily at room temp. in the presence of air. To 100 ml. anhyd. NH3 at -70° were added 0.1 g. Fe(NO3)3 and 2.99 g. Na and, after disappearance of the blue color, 25 g. C6F5OMe during 45 min. After 5 hrs. at -70° the reaction mixt. was worked up to give 7 g. unreacted C6F5OMe, 2.8 g. tetrafluoro-p-anisidine, m. 75-6.5°, 1.2 g. 4,4'-dimethoxyoctafluorodiphenylamine, m. 78-9°, and 2.2 g. 4,4',4''-trimethoxydecafluorotriphenylamine, b. 157-9°, n23D 1.5005. Diazotization of C6F5NH2 required concd. acids since the salts of the amine hydrolyzed very readily in dil. solns. In 48% HBr, diazotization of C6F5NH2 gave C6F5N:NHCl6F5, probably owing to slow diazotization. The reaction was temp.-dependent, the diazoaminobenzene being formed much faster at 10° than at -10°. The product decompd. in warm HBr to give 5.4% C6F5Br and a mixt. of o- and p-dibromotetrafluorobenzenes. In concd. H2SO4 the reaction was very slow even at 25°. Addition of HQAc hastened it. Deamination with hypophosphorus acid gave a mixt. probably of C6F5H and C6H2F4. Better diazotization could be carried out in liquid HF and the diazo product underwent successful Sandmeyer reaction. A soln. of 20 g. C6F5NH2 in 75 ml. anhyd. HF at -20° was treated with 7.27 g. NaN02 during 30 min. After stirring for 1 hr. at -10° the mixt. was treated with 17.6 g. KI during 30 min. and allowed to warm to 25° in 1 hr. to give 16.5 g. C6F5I, b35 77-9°. Use of 12 g. KBr and 15 g. Cu2Br2 instead of KI gave 35% C6F5Br. The reaction of diazotized amine with C6F5OLi gave C6F5N:N(O)C6F5, which decompd. on removal of solvent. The Sandmeyer nitrile synthesis was not successful. A soln. of 10 g. C6F5NH2 in 100 ml. HOAc was oxidized with 25 ml. 30% H2O2 at 25° for 24 hrs. to give decafluoroazoxybenzene (IV), m. 53-4°. A mixt. of 5 g. IV, 15 g. Zn powder, 5 g. NH4Cl, 10 ml. H2O, and 75 ml. 95% EtOH refluxed 30 min. gave 2 g. decafluoroazobenzene, m. 57-9°. C6F6 reacted readily with organolithium compds. A soln. of MeLi, prepd. from 4.5 g. Li and 43 g. MeI in 50 ml. ether, was cooled to -10° to -20°, added dropwise to a soln. of 60 g. C6F6 in 250 ml. pentane, and stirred for 17 hrs. at room temp. to give 34 g. C6F5Me, b. 115°. Similarly, reaction of BuLi, prepd. from 1.06 g. Li and 18.3 g. BuBr in 30 ml. ether,



L64 ANSWER 30 OF 30 CAPLUS COPYRIGHT 2005 ACS on STN (Continued)  
 with 25.3 g. C6F6 in 25 ml. ether gave 10.5 g. unreacted C6F6, 7 g.  
 C6F5Bu, b25 86-7°, n20D 1.4229, and 2.5 g. of a compd., b1  
 230°, n20D 1.4683, probably impure tributylidifluorobenzene.  
 Similarly, 32.7 g. C6F6 in 150 ml. ether with 0.18 mole PhLi in 250 ml.  
 ether gave 8.5 g. 2,3,5,6-tetrafluoro-p-terphenyl, m. 220°, and 33  
 g. C6F5Ph, m. 69°. A similar reaction of 18.6 g. C6F6 and  
 isopropenyllithium prepd. from 12.1 g. 2-bromopropene gave 5 g.  
 2,3,4,5,6-pentafluoro- $\alpha$ -methylstyrene, b52 72-4°. With  
 vinylolithium, prepd. from 0.1 mole PhLi and 0.025 mole tetravinyltin,  
 10.6  
 g. C6F6 gave 4 g. unreacted C6F6 and 20% C6F5CH:CH2, b25 34°.  
 LiAlH4 redn. of 21 g. C6F6 in ether gave 17 g. of a mixt. of C6F6 and  
 C6F5H which was sepd. by vapor phase chromatography to give 7.5 g. C6HF5.  
 All the products in all above reactions were studied by infrared and  
 nuclear magnetic resonance spectroscopy. The mechanism of reaction and  
 the directional effects were discussed.

ACCESSION NUMBER: 1964:52434 CAPLUS  
 DOCUMENT NUMBER: 60:52434  
 ORIGINAL REFERENCE NO.: 60:9170b-h, 9171a-e  
 TITLE: Reactions of polyfluorobenzenes with nucleophilic  
 reagents  
 AUTHOR(S): Wall, Leo A.; Pummer, Walter J.; Fearn, James E.;  
 Antonucci, Joseph M.  
 SOURCE: J. Res. Natl. Bur. Std. (1963), 67A(5), 481-97  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 IT 6804-37-1, Benzene, 1,2,4,5-tetrafluoro-3,6-  
 bis(pentafluorophenoxy)-(?)  
 (preparation of)  
 RN 6804-37-1 CAPLUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)- (7CI, 8CI) (CA  
 INDEX NAME)



```
=> fil reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                                ENTRY      SESSION
FULL ESTIMATED COST          150.00      5056.53

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  SINCE FILE      TOTAL
                                                ENTRY      SESSION
CA SUBSCRIBER PRICE          -21.90      -374.49
```

FILE 'REGISTRY' ENTERED AT 20:41:08 ON 12 JAN 2005  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8  
 DICTIONARY FILE UPDATES: 10 JAN 2005 HIGHEST RN 811411-12-8


TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
 information enter HELP PROP at an arrow prompt in the file or refer  
 to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=>
Uploading C:\Program Files\Stnexp\Queries\10718532.str
```



```
chain nodes :
1 13 23 25 26 28 29 30 31 32 33 35
ring nodes :
2 3 4 5 6 7 8 9 10 11 12 16 17 18 19 20 21 22
chain bonds :
1-4 8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31
22-35
ring bonds :
2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22
18-19 19-20 20-21 21-22
exact/norm bonds :
8-29 9-28 10-13 11-25 12-23 16-26 17-32 18-33 19-23 20-30 21-31
exact bonds :
1-4 22-35
```

2-3 2-7 3-4 4-5 5-6 6-7 8-9 8-12 9-10 10-11 11-16 12-16 17-18 17-22  
18-19 19-20 20-21 21-22


G2 : X, Ak, H

```

1:CLASS 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:CLASS 14:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom
21:Atom 22:Atom 23:CLASS 25:CLASS 26:CLASS 28:CLASS 29:CLASS 30:CLASS
31:CLASS 32:CLASS 33:CLASS 35:CLASS

```

L65 STR



G2 X, Ak, H

SAMPLE SEARCH INITIATED 20:41:56 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 51 TO ITERATE

0 ANSWERS

```

FULL FILE PROJECTIONS:  ONLINE  **COMPLETE**
                        BATCH    **COMPLETE**
PROJECTED ITERATIONS:   592 TO      1448
PROJECTED ANSWERS:      0 TO        0

```

Page 481

=> s 165 full  
FULL SEARCH INITIATED 20:42:01 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 885 TO ITERATE

100.0% PROCESSED 885 ITERATIONS 5 ANSWERS  
SEARCH TIME: 00.00.01

L67 5 SEA SSS FUL L65

=> fil caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	161.76	5218.29
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-374.49


FILE 'CAPLUS' ENTERED AT 20:42:04 ON 12 JAN 2005  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 12 Jan 2005 VOL 142 ISS 3  
FILE LAST UPDATED: 11 Jan 2005 (20050111/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 167  
L68 6 L67  
  
=> d 168 1-6 abs. ibib hitstr



AB Benzyl cyclopropanecarboxylates I (R = H, F, Cl, Br; R1 = H, cyano, C.tplbond.CH; R2 = H, Me; R3 = CH:CCl2, CHBrCBrCl2, CHBrCBr3, Me), having insecticidal and acaricidal activities at 500 ppm with low fish toxicity, were prepared e.g. from II. Thus, 6.0 mmol NaCN and 0.88 mmol PhCH2NEt3Cl

in H2O was treated dropwise with 4.0 mmol II and 4.2 mmol dl-cis,trans-2,2-dimethyl-3-(2,2-dichlorovinyl)cyclopropanecarbonyl chloride in PhMe to give 1.93 g Y (R = R2 = H, R1 = cyano, R3 = CH:CCl2).

ACCESSION NUMBER: 1984:610760 CAPIUS  
DOCUMENT NUMBER: 101:210760  
TITLE: Insecticidal cyclopropanecarboxylate esters  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.  
CODEN: JKOQAF


DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59134749	A2	19840802	JP 1983-8536	19830120



PRIORITY APPLN. INFO.: JP 1983-8536 19830120

IT 90928-31-7P 93207-48-8P  
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation and insecticidal activity of)

RN 90928-31-7 CAPIUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2-tetrabromoethyl)-, 1-(4-fluoro-3-(4-(4-fluorophenoxy)phenoxy)phenyl)-2-propynyl ester (9CI) (CA INDEX NAME)



RN 93207-48-8 CAPIUS  
CN Cyclopropanecarboxylic acid, 3-(2,2-dichloroethenyl)-2,2-dimethyl-, cyano[4-(4-fluoro-3-(4-(4-fluorophenoxy)phenoxy)phenyl)methyl ester (9CI) (CA INDEX NAME)



AB Twelve title esters (I; R = H, halo; R1 = H, F; R2 = H, cyano, HC.tplbond.C, R3 = haloethyl, haloethyl), effective insecticides and miticides at 500 ppm, were prepared. Thus, 5.0 mmol II and 5.25 mmol (I)-III in MePh were added to a solution of 7.5 mmol NaCN and 1.1 mmol PhCH2NEt3 Cl- in H2O at room temperature to give 2.07 g (2)-I (R = R1 = H;

R2 = cyano, R3 = Cl2C:CH).

ACCESSION NUMBER: 1984:438160 CAPIUS  
DOCUMENT NUMBER: 101:38160  
TITLE: Cyclopropanecarboxylate esters as insecticides and miticides with low toxicity  
PATENT ASSIGNEE(S): Sumitomo Chemical Co., Ltd., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.  
CODEN: JKOQAF


DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59042342	A2	19840308	JP 1982-153029	19820901

PRIORITY APPLN. INFO.: JP 1982-153029 19820901

IT 90928-31-7P  
RL: SPN (Synthetic preparation); PREP (Preparation) (preparation of)

RN 90928-31-7 CAPIUS  
CN Cyclopropanecarboxylic acid, 2,2-dimethyl-3-(1,2,2,2-tetrabromoethyl)-, 1-(4-fluoro-3-(4-(4-fluorophenoxy)phenoxy)phenyl)-2-propynyl ester (9CI) (CA INDEX NAME)




L68 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB Reaction of phenols with Br<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CF<sub>3</sub> at 170-300° in the presence of Cu and alkali gave mono- and diphenoxylbenzotrifluorides. Thus, 2,5- and 3,4-Br<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CF<sub>3</sub>, powdered Cu, KI, and m-CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>OH at 200-20° gave 10.6 (m-CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>O)C<sub>6</sub>H<sub>4</sub>CF<sub>3</sub>, 25.4 (m-CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>O)C<sub>6</sub>H<sub>3</sub>BrCF<sub>3</sub>, and 42.4% (m-CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>O)C<sub>6</sub>H<sub>3</sub>CF<sub>3</sub> isomers.

ACCESSION NUMBER: 1975:170346 CAPLUS  
 DOCUMENT NUMBER: 82:170346  
 TITLE: Phenoxy derivatives of trifluoromethylbenzene  
 INVENTOR(S): Schlafke, Rolf; Jenkner, Herbert  
 PATENT ASSIGNEE(S): Chemische Fabrik Kalk G.m.b.H.  
 SOURCE: Ger. Offen., 21 pp.  
 CODEN: GWXBX

DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2325878	A1	19741212	DE 1973-2325878	19730522
PRIORITY APPLN. INFO.:			DE 1973-2325878	A 19730522

IT 54846-39-8P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 54846-39-8 CAPLUS  
 CN Benzene, 1,4-bis(4-fluorophenoxy)-2-(trifluoromethyl)- (9CI) (CA INDEX NAME)




L68 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A number of perfluorinated and partially fluorinated phenyl and polyphenyl ethers were synthesized, characterized for thermal stability, fire resistance, and viscosity, and compared with their H analogs to assess the potential use of this class of compds. as functional fluids. Without exception, polyfluorination and perfluorination lower thermal stability; the decrease in stability depends on the position and number of fluorine substituents. The autoignition temperature and fire resistance are not improved over the H analog, and viscosity is degraded. These data coupled with the comparatively high melting points do not suggest a bright future for this class of compds. as useful functional fluids. 6 references.

ACCESSION NUMBER: 1968:77890 CAPLUS  
 DOCUMENT NUMBER: 68:77890  
 TITLE: Synthesis, thermal stability, flammability, and viscosity of some partially fluorinated and perfluorinated aromatic and polyaromatic ethers  
 AUTHOR(S): Richardson, George Albert; Blake, Edward S.  
 CORPORATE SOURCE: Monsanto Res. Corp., Dayton, OH, USA  
 SOURCE: Industrial & Engineering Chemistry Product Research and Development (1968), 7(1), 17-21  
 CODEN: IEPRA6; ISSN: 0196-4321  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 IT 6804-37-1P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 6804-37-1 CAPLUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)- (7CI, 8CI) (CA INDEX NAME)




L68 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2005 ACS on STN  
 GI For diagram(s), see printed CA Issue.  
 AB The PbO<sub>2</sub> oxidation of pentafluorophenol gave the oxocyclohexadienyl phenyl ether (I). Na phenoxides (II and III) reacted with Br to give ethers (IV and V).  
 ACCESSION NUMBER: 1974:520135 CAPLUS  
 DOCUMENT NUMBER: 81:120135  
 TITLE: Polyfluorophenols. I. Mild oxidation of pentafluorophenol  
 AUTHOR(S): Deniville, Leon; Huynh Anh Hoa  
 CORPORATE SOURCE: Lab. Chim. Text. Tinctoriale, Conservatoire Natl. Arts  
 SOURCE: Metiers, Paris, Fr.  
 Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 487-90  
 CODEN: BSCFAS; ISSN: 0037-8968  
 DOCUMENT TYPE: Journal  
 LANGUAGE: French  
 IT 53359-93-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 53359-93-6 CAPLUS  
 CN Phenol, 2,3,5,6-tetrafluoro-4-(2,3,5,6-tetrafluoro-4-(pentafluorophenoxy)phenoxy)- (9CI) (CA INDEX NAME)



L68 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2005 ACS on STN  
 AB A mixture of 26.5 g. C<sub>6</sub>F<sub>6</sub>, 26.5 g. 85% KOH, and 75 ml. H<sub>2</sub>O was heated in a sealed bomb at 175° for 5 hrs. with agitation to give 33.1 g. C<sub>6</sub>F<sub>5</sub>OH, b. 144-5°. A mixture of 67 g. C<sub>6</sub>F<sub>5</sub>H, 21.6 g. KOH, 150 ml. pyridine, and 2 ml. H<sub>2</sub>O was refluxed 1 hr., treated with 21.6 KOH, and refluxed 24 hrs. to give 21 g. 2,3,5,6-tetrafluorophenol, b<sub>20</sub> 47°. Similarly refluxing 36 g. C<sub>6</sub>F<sub>5</sub>Me and 28 g. KOH in 300 ml. tert-BuOH gave 12 g. 2,3,5,6-tetrafluoro-p-cresol, m. 52°. Reaction of 20 g. C<sub>6</sub>F<sub>5</sub>I with 2 g. KOH and 1 ml. H<sub>2</sub>O in 100 ml. pyridine gave only one product, 2,3,5,6-tetrafluoro-4-iodophenol, m. 79-81°; benzoate m. 59-60.2°. Reaction of 66 g. C<sub>6</sub>F<sub>5</sub>Br with 28.5 g. KOH and 1 ml. H<sub>2</sub>O in 150 ml. pyridine, however, gave a mixture of products: 3.5 g. 2-bromo-3,4,5,6-tetrafluorophenol, m. 41-3° (3,5-dinitrobenzoate, m. 104-5°); and 11.5 g. 4-bromo-2,3,5,6-tetrafluorophenol (3,5-dinitrobenzoate m. 131-3°). Similarly, 100 g. 2-chlorotetrafluoro- $\alpha,\alpha,\alpha$ -trifluorotoluene on treatment with 5.6 g. KOH and 1 ml. H<sub>2</sub>O in 100 ml. pyridine gave 2.5 g. 2-chlorotrifluoro- $\alpha,\alpha,\alpha$ -trifluoro-o-cresol, b<sub>15</sub> 92-3°, n<sub>D</sub> 1.4510; and 15 g. 2-chlorotrifluoro- $\alpha,\alpha,\alpha$ -trifluoro-p-cresol, b<sub>15</sub> 102-3°, n<sub>D</sub> 1.4510. The reactions of polyfluorobenzenes were then studied with alkoxides. A solution of 123.5 g. C<sub>6</sub>F<sub>5</sub>Br in 70 ml. pyridine was treated with a solution of 11.5 g. Na in 150 ml. MeOH during 1.5 hrs. and the mixture refluxed 15 hrs. and acidified with 1 l. 10% HCl to give 66 g. 4-bromo-2,3,5,6-tetrafluoroanisole, b<sub>5</sub> 79-81°, n<sub>D</sub> 1.4812. Similarly, a mixture of 10 g. C<sub>6</sub>F<sub>5</sub>I in 50 ml. pyridine and 0.8 g. Na in 15 ml. MeOH on refluxing for 3 hrs. gave 1.5 g. unchanged C<sub>6</sub>F<sub>5</sub>I and 5.5 g. 2,3,5,6-tetrafluoro-4-iodoanisole (I), b<sub>20</sub> 113-15°, n<sub>D</sub> 1.5229. Refluxing 1 g. I with 1 g. activated Cu powder for 12 min. gave 0.2 g. octafluoro-4,4'-dimethoxybiphenyl, m. 90-1.2°. To a cold solution of 9 g. Na in 250 ml. PhCH<sub>2</sub>OH was added 75 g. C<sub>6</sub>F<sub>6</sub> and the mixture refluxed 15 hrs. to give 30 g. benzyl pentafluorophenyl ether (II), m. 44°. A better yield was obtained when a solution of 4.6 g. Na and 22 g. PhCH<sub>2</sub>OH in 250 ml. tert-BuOH was refluxed with 40 g. C<sub>6</sub>F<sub>6</sub> for 40 hrs. to give 33 g. II. A solution of 8 g. C<sub>6</sub>F<sub>6</sub> in 30 ml. HCONMe<sub>2</sub> was treated with 5.28 g. PhOK and the mixture refluxed 0.5 hr. to give 1 g. 2,3,5,6-tetrafluoro-1,4-diphenoxybenzene, m. 147-9°, and 3.5 g. 2,3,4,5,6-pentafluorophenyl phenyl ether, (III), m. 29°. III was also obtained by heating a mixture of 11 g. C<sub>6</sub>F<sub>5</sub>OK, 15 g. PhBr, and 1 g. Cu at 210° in a sealed bomb. A solution of 6 g. C<sub>6</sub>F<sub>5</sub>OK and 12.8 g. C<sub>6</sub>F<sub>6</sub> in 30 ml. HCONMe<sub>2</sub> was refluxed 14 hrs. to give 1.5 g. bis(perfluorophenyl) ether, m. 67-9°, and a second product, m. 145-8°, probably p-bis(pentafluorophenoxy)2,3,5,6-tetrafluorobenzene. Similarly, a solution of 0.6 g. Na in 50 ml. EtOH refluxed with 3.3 g. C<sub>6</sub>F<sub>5</sub>NMe<sub>2</sub> 2 hrs. gave 5.1 g. 4-ethoxy-2,3,5,6-tetrafluoro-N,N-dimethylaniline, b. 34°. Reactions with amines were next investigated. A mixture of 280 g. C<sub>6</sub>F<sub>6</sub> and 400 ml. 28% aqueous NH<sub>3</sub> was rocked in a sealed bomb for 2 hrs. at 235° to give 236 g. C<sub>6</sub>F<sub>5</sub>NH<sub>2</sub>, m. 34°, and 28 g. tetrafluorophenylenediamine (sublimed 75°/1 mm.) shown by its nuclear magnetic resonance spectrum to be essentially the meta isomer mixed with a small amount of the para isomer. Similarly, heating a mixture of

168 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
 56 g. C6F6 and 110 ml. 30% aq. MeNH2 at 220° for 3 hrs. gave 59% C6F5NMe, b. 170-2°, and 25% 2,3,5,6-tetrafluoro-N,N'-diphenylphenylenediamine, m. 94°. The reaction product obtained by heating 50 g. C6F6 and 110 ml. 25% aq. Me2NH at 235° for 1 hr. was distd. at 1 mm. pressure and five fractions were collected. The first fraction (65%), b. 88°, was C6F5NMe2. Fraction 2, b. 88-126°, was shown by vapor phase chromatography to be C6F5NMe2 with 3 other compds. Fraction 3, b. 126-134°, consisted of 3 isomers of bis(dimethylamino)tetrafluorobenzene with the meta-isomer predominating. Fraction 4, b. 134-40°, contained equal amts. of the meta and para isomers. Fraction 5, b. 140-8°, was pure para isomer. The meta and para isomers could be sepd. by vapor phase chromatography. Similarly, heating a mixt. of 30 g. C6F5Br and 70 ml. NH4OH at 200° for 2 hrs. gave 22 g. p-bromotetrafluoroaniline, m. 61°. Heating 16 g. C6F5I and 30 ml. 8% NH4OH at 165° for 2 hrs. gave 7.6 g. tetrafluoro-p-iodoaniline, m. 77°. Benzyl pentafluorophenyl ether (20 g.) was heated with large excess of 28% NH4OH to give 3 g. p-(benzyloxy)tetrafluoroaniline, m. 97°. Similarly, heating 50 g. 2-chlorotetrafluoro-*o,o',o''*-trifluorotoluene and 120 ml. 28% NH4OH at 21° for 2 hrs. gave 22 g. 2-chlorotrifluoro-*o,o',o''*-trifluoro-p-toluidine, which decompd. readily at room temp. in the presence of air. To 100 ml. anhyd. NH3 at -70° were added 0.1 g. Fe(NO3)3 and 2.99 g. Na and, after disappearance of the blue color, 25 g. C6F5OMe during 45 min. After 5 hrs. at -70° the reaction mixt. was worked up to give 7 g. unreacted C6F5OMe, 2.8 g. tetrafluoro-p-anisidine, m. 75-6.5°, 1.2 g. 4,4'-dimethoxyoctafluorodiphenylamine, m. 78-9°, and 2.2 g. 4,4',4''-trimethoxydecafluorotriphenylamine, b. 157-9°, n23D 1.5005. Diazotization of C6F5NH2 required concd. acids since the salts of the amine hydrolyzed very readily in dil. solns. In 48% HBr, diazotization of C6F5NH2 gave C6F5N:NNHC6F5, probably owing to slow diazotization. The reaction was temp.-dependent, the diazoaminobenzene being formed much faster at 10° than at -10°. The product decompd. in warm HBr to give 5.4% C6F5Br and a mixt. of *o*- and *p*-dibromotetrafluorobenzenes. In concd. H2SO4 the reaction was very slow even at 25°. Addition of HOAc hastened it. Deamination with hypophosphorus acid gave a mixt. probably of C6F5H and C6H2F4. Better diazotization could be carried out in liquid HF and the diazo product underwent successful Sandmeyer reaction. A soln. of 20 g. C6F5NH2 in 75 ml. anhyd. HF at -20° was treated with 7.27 g. NaNO2 during 30 min. After stirring for 1 hr. at -10° the mixt. was treated with 17.6 g. KI during 30 min. and allowed to warm to 25° in 1 hr. to give 16.5 g. C6F5I, b35 77-9°. Use of 12 g. KBr and 15 g. Cu2Br2 instead of KI gave 35% C6F5Br. The reaction of diazotized amine with C6F5OLi gave C6F5N:N(O)C6F5, which decompd. on removal of solvent. The Sandmeyer nitrile synthesis was not successful. A soln. of 10 g. C6F5NH2 in 100 ml. HOAc was oxidized with 25 ml. 30% H2O2 at 25° for 24 hrs. to give decafluoroazoxybenzene (IV), m. 53-4°. A mixt. of 5 g. IV, 15 g. Zn powder, 5 g. NH4Cl, 10 ml. H2O, and 75 ml. 95% EtOH refluxed 30 min. gave 2 g. decafluoroazobenzene, m. 57-9°. C6F6 reacted readily with organolithium compds. A soln. of MeLi, prepd. from 4.5 g. Li and 43 g. MeI in 50 ml. ether, was cooled to -10° to -20°, added dropwise to a soln. of 60 g. C6F6 in 250 ml. pentane, and stirred for 17 hrs. at room temp. to give 34 g. C6F5Me, b. 113°. Similarly, reaction of BuLi, prepd. from 1.86 g. Li and 18.3 g. BuBr in 30 ml. ether,

168 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2005 ACS ON STN (Continued)  
 with 25.3 g. C6F6 in 25 ml. ether gave 10.5 g. unreacted C6F6, 7 g. C6F5Bu, b25 86-7°, n20D 1.4229, and 2.5 g. of a compd., b. 230°, n20D 1.4683, probably impure tributylidifluorobenzene. Similarly, 32.7 g. C6F6 in 150 ml. ether with 0.18 mole PhLi in 250 ml. ether gave 8.5 g. 2,3,5,6-tetrafluoro-*p*-terphenyl, m. 220°, and 33 g. C6F5Ph, m. 69°. A similar reaction of 18.6 g. C6F6 and isopropenyllithium prepd. from 12.1 g. 2-bromopropene gave 5 g. 2,3,4,5,6-pentafluoro-*o*-methylstyrene, b52 72-4°. With vinylolithium, prepd. from 0.1 mole PhLi and 0.025 mole tetravinyltin, 18.6 g. C6F6 gave 4 g. unreacted C6F6 and 20% C6F5CH:CH2, b25 34°. LiAlH4 redn. of 21 g. C6F6 in ether gave 17 g. of a mixt. of C6F6 and C6F5H which was sepd. by vapor phase chromatography to give 7.5 g. C6H5F. All the products in all above reactions were studied by infrared and nuclear magnetic resonance spectroscopy. The mechanism of reaction and the directional effects were discussed.  
 ACCESSION NUMBER: 1964:52434 CAPLUS  
 DOCUMENT NUMBER: 60:52434  
 ORIGINAL REFERENCE NO.: 60:9170b-h, 9171a-e  
 TITLE: Reactions of polyfluorobenzenes with nucleophilic reagents  
 AUTHOR(S): Wall, Leo A.; Pummer, Walter J.; Fearn, James E.; Antonucci, Joseph M.  
 SOURCE: J. Res. Natl. Bur. Std. (1963), 67A(5), 481-97  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 IT 6804-37-1, Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)-(?)  
 (preparation of)  
 RN 6804-37-1 CAPLUS  
 CN Benzene, 1,2,4,5-tetrafluoro-3,6-bis(pentafluorophenoxy)- (7CI, 8CI) (CA INDEX NAME)



=> logoff y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

30.09

5248.38

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-4.38

-378.87

STN INTERNATIONAL LOGOFF AT 20:42:45 ON 12 JAN 2005